Caroline Even

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11955086/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Nivolumab for Recurrent Squamous-Cell Carcinoma of the Head and Neck. New England Journal of Medicine, 2016, 375, 1856-1867.	27.0	3,845
2	Nivolumab vs investigator's choice in recurrent or metastatic squamous cell carcinoma of the head and neck: 2-year long-term survival update of CheckMate 141 with analyses by tumor PD-L1 expression. Oral Oncology, 2018, 81, 45-51.	1.5	589
3	High-Throughput Genomics and Clinical Outcome in Hard-to-Treat Advanced Cancers: Results of the MOSCATO 01 Trial. Cancer Discovery, 2017, 7, 586-595.	9.4	554
4	Safety and Antitumor Activity of Pembrolizumab in Patients With Programmed Death-Ligand 1–Positive Nasopharyngeal Carcinoma: Results of the KEYNOTE-028 Study. Journal of Clinical Oncology, 2017, 35, 4050-4056.	1.6	335
5	Nivolumab versus standard, single-agent therapy of investigator's choice in recurrent or metastatic squamous cell carcinoma of the head and neck (CheckMate 141): health-related quality-of-life results from a randomised, phase 3 trial. Lancet Oncology, The, 2017, 18, 1104-1115.	10.7	325
6	Safety and Efficacy of Durvalumab With or Without Tremelimumab in Patients With PD-L1–Low/Negative Recurrent or Metastatic HNSCC. JAMA Oncology, 2019, 5, 195.	7.1	235
7	Durvalumab for recurrent or metastatic head and neck squamous cell carcinoma: Results from a single-arm, phase II study in patients with ≥25% tumour cell PD-L1 expression who have progressed on platinum-based chemotherapy. European Journal of Cancer, 2019, 107, 142-152.	2.8	208
8	Nivolumab in Patients with Recurrent or Metastatic Squamous Cell Carcinoma of the Head and Neck: Efficacy and Safety in CheckMate 141 by Prior Cetuximab Use. Clinical Cancer Research, 2019, 25, 5221-5230.	7.0	115
9	Response to salvage chemotherapy after progression on immune checkpoint inhibitors in patients with recurrent and/or metastatic squamous cell carcinoma of the head and neck. European Journal of Cancer, 2019, 121, 123-129.	2.8	115
10	Cetuximab, docetaxel, and cisplatin versus platinum, fluorouracil, and cetuximab as first-line treatment in patients with recurrent or metastatic head and neck squamous-cell carcinoma (GORTEC) Tj ETQq0	0 0 rgBT // 10.7	Overlock 10 1
11	Trabectedin in patients with advanced soft tissue sarcoma: A retrospective national analysis of the French Sarcoma Group. European Journal of Cancer, 2015, 51, 742-750.	2.8	86
12	Impact of invasive fungal disease on the chemotherapy schedule and event-free survival in acute leukemia patients who survived fungal disease: a case-control study. Haematologica, 2011, 96, 337-341.	3.5	72
13	CheckMate 141: 1‥ear Update and Subgroup Analysis of Nivolumab as Firstâ€Line Therapy in Patients with Recurrent/Metastatic Head and Neck Cancer. Oncologist, 2018, 23, 1079-1082.	3.7	70
14	TPF induction chemotherapy increases PD-L1 expression in tumour cells and immune cells in head and neck squamous cell carcinoma. ESMO Open, 2018, 3, e000257.	4.5	62
15	Avelumab–cetuximab–radiotherapy versus standards of care in locally advanced squamous-cell carcinoma of the head and neck: The safety phase of a randomised phase III trial GORTEC 2017-01 (REACH). European Journal of Cancer, 2020, 141, 21-29.	2.8	48
16	Induction chemotherapy with docetaxel, cisplatin and fluorouracil followed by concurrent chemoradiotherapy or chemoradiotherapy alone in locally advanced non-endemic nasopharyngeal carcinoma. Oral Oncology, 2016, 62, 114-121.	1.5	43
17	Phase II, Randomized Study of Spartalizumab (PDR001), an Anti–PD-1 Antibody, versus Chemotherapy in Patients with Recurrent/Metastatic Nasopharyngeal Cancer. Clinical Cancer Research, 2021, 27, 6413-6423.	7.0	37
18	A phase II study of monalizumab in patients with recurrent/metastatic squamous cell carcinoma of the head and neck: The I1 cohort of the EORTC-HNCG-1559 UPSTREAM trial. European Journal of Cancer, 2021, 158, 17-26.	2.8	33

CAROLINE EVEN

#	Article	IF	CITATIONS
19	Does smoking alter the mutation profile of human papillomavirus–driven head and neck cancers?. European Journal of Cancer, 2018, 94, 61-69.	2.8	29
20	Randomized trial comparing two methods of re-irradiation after salvage surgery in head and neck squamous cell carcinoma: Once daily split-course radiotherapy with concomitant chemotherapy or twice daily radiotherapy with cetuximab. Radiotherapy and Oncology, 2018, 128, 467-471.	0.6	18
21	PACSA: Phase II study of pazopanib in patients with progressive recurrent or metastatic (R/M) salivary gland carcinoma (SGC) Journal of Clinical Oncology, 2016, 34, 6086-6086.	1.6	18
22	Long-term Outcomes with Nivolumab as First-line Treatment in Recurrent or Metastatic Head and Neck Cancer: Subgroup Analysis of CheckMate 141. Oncologist, 2022, 27, e194-e198.	3.7	18
23	Anemia and neutrophilâ€toâ€lymphocyte ratio in laryngeal cancer treated with induction chemotherapy. Laryngoscope, 2020, 130, E144-E150.	2.0	15
24	Impact of previous nivolumab treatment on the response to taxanes in patients with recurrent/metastatic head and neck squamous cell carcinoma. European Journal of Cancer, 2021, 159, 125-132.	2.8	11
25	Relationship between the time to locoregional recurrence and survival in laryngeal squamous-cell carcinoma. European Archives of Oto-Rhino-Laryngology, 2017, 274, 2267-2271.	1.6	10
26	Laryngo-esophageal Dysfunction-free Survival in a Preservation Protocol for T3 Laryngeal Squamous-cell Carcinoma. Anticancer Research, 2016, 36, 6625-6630.	1.1	10
27	Patterns of disease events and causes of death in patients with HPV-positive versus HPV-negative oropharyngeal carcinoma. Radiotherapy and Oncology, 2022, 168, 40-45.	0.6	10
28	High incidence of cetuximab-related infusion reactions in head and neck patients. ESMO Open, 2018, 3, e000346.	4.5	8
29	New approaches in salivary gland carcinoma. Current Opinion in Oncology, 2019, 31, 169-174.	2.4	8
30	Notch pathway inhibition with LY3039478 in adenoid cystic carcinoma (ACC) Journal of Clinical Oncology, 2017, 35, 6024-6024.	1.6	8
31	Outcomes following laryngectomy refusal after insufficient response to induction chemotherapy. Laryngoscope, 2017, 127, 1791-1796.	2.0	7
32	Can radiation-recall predict long lasting response to immune checkpoint inhibitors?. Radiotherapy and Oncology, 2021, 154, 125-127.	0.6	7
33	Efficacy and safety of immune checkpoint inhibitors in elderly patients (≥70 years) with squamous cell carcinoma of the head and neck. European Journal of Cancer, 2021, 157, 190-197.	2.8	6
34	Association of PD-L1 Expression on Tumor and Immune Cells with Survival in Recurrent or Metastatic Head and Neck Squamous Cell Carcinoma and Assay Validation. Cancer Research Communications, 2022, 2, 39-48.	1.7	4
35	Computed tomography evaluation after induction chemotherapy for T3 laryngeal cancer: Does response correlate with vocal cord mobility?. Oral Oncology, 2019, 90, 13-16.	1.5	2
36	Complete response upon salvage chemotherapy after anti-PD1 failure: Watch and wait. European Journal of Cancer, 2021, 145, 155-157.	2.8	2

#	Article	IF	CITATIONS
37	Cabazitaxel in recurrent/metastatic squamous cell carcinoma of the head and neck: phase II UNICANCER trial ORL03. Oncotarget, 2017, 8, 51830-51839.	1.8	2
38	Panitumumab as an effective maintenance treatment in metastatic squamous cell carcinoma of the head and neck. Oral Oncology, 2021, 112, 104984.	1.5	1