
## Eleazar Soto

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11965159/publications.pdf Version: 2024-02-01



FLEAZAD SOTO

#	Article	IF	CITATIONS
1	Normal pregnancy is characterized by systemic activation of the complement system. Journal of Maternal-Fetal and Neonatal Medicine, 2005, 17, 239-245.	0.7	172
2	Late-onset preeclampsia is associated with an imbalance of angiogenic and anti-angiogenic factors in patients with and without placental lesions consistent with maternal underperfusion. Journal of Maternal-Fetal and Neonatal Medicine, 2012, 25, 498-507.	0.7	136
3	Maternal plasma concentrations of angiogenic/antiangiogenic factors in the third trimester of pregnancy to identify the patient at risk for stillbirth at or near term and severe late preeclampsia. American Journal of Obstetrics and Gynecology, 2013, 208, 287.e1-287.e15.	0.7	122
4	Maternal plasma concentrations of angiogenic/anti-angiogenic factors are of prognostic value in patients presenting to the obstetrical triage area with the suspicion of preeclampsia. Journal of Maternal-Fetal and Neonatal Medicine, 2011, 24, 1187-1207.	0.7	118
5	A rapid MMP-8 bedside test for the detection of intra-amniotic inflammation identifies patients atÂriskÂforÂimminent preterm delivery. American Journal of Obstetrics and Gynecology, 2006, 195, 1025-1030.	0.7	87
6	Human β-defensin-2: A natural antimicrobial peptide present in amniotic fluid participates in the host response to microbial invasion of the amniotic cavity. Journal of Maternal-Fetal and Neonatal Medicine, 2007, 20, 15-22.	0.7	80
7	Evidence for complement activation in the amniotic fluid of women with spontaneous preterm labor and intra-amniotic infection. Journal of Maternal-Fetal and Neonatal Medicine, 2009, 22, 983-992.	0.7	53
8	Twin-to-twin transfusion syndrome: an antiangiogenic state?. American Journal of Obstetrics and Gynecology, 2008, 198, 382.e1-382.e8.	0.7	51
9	Preeclampsia and pregnancies with small-for-gestational age neonates have different profiles of complement split products. Journal of Maternal-Fetal and Neonatal Medicine, 2010, 23, 646-657.	0.7	48
10	What Does 2â€Dimensional Imaging Add to 3―and 4â€Dimensional Obstetric Ultrasonography?. Journal of Ultrasound in Medicine, 2006, 25, 691-699.	0.8	44
11	The diagnostic performance of the Mass Restricted (MR) score in the identification of microbial invasion of the amniotic cavity or intra-amniotic inflammation is not superior to amniotic fluid interleukin-6. Journal of Maternal-Fetal and Neonatal Medicine, 2014, 27, 757-769.	0.7	44
12	Anaphylatoxins in preterm and term labor. Journal of Perinatal Medicine, 2005, 33, 306-13.	0.6	36
13	Fetal Anomalies and Long-Term Effects Associated with Substance Abuse in Pregnancy: A Literature Review. American Journal of Perinatology, 2015, 32, 405-416.	0.6	35
14	Maternal serum soluble CD30 is increased in normal pregnancy, but decreased in preeclampsia and small for gestational age pregnancies. Journal of Maternal-Fetal and Neonatal Medicine, 2007, 20, 867-878.	0.7	34
15	Fragment Bb in amniotic fluid: evidence for complement activation by the alternative pathway in women with intra-amniotic infection/inflammation. Journal of Maternal-Fetal and Neonatal Medicine, 2009, 22, 905-916.	0.7	33
16	ORIGINAL ARTICLE: Activation of the Alternative Pathway of Complement is a Feature of Preâ€Term Parturition but not of Spontaneous Labor at Term. American Journal of Reproductive Immunology, 2010, 63, 318-330.	1.2	32
17	Objective Evaluation of Sylvian Fissure Development by Multiplanar 3-Dimensional Ultrasonography. Journal of Ultrasound in Medicine, 2007, 26, 347-353.	0.8	28
18	Plasma protein Z concentrations in pregnant women with idiopathic intrauterine bleeding and in women with spontaneous preterm labor. Journal of Maternal-Fetal and Neonatal Medicine, 2007, 20, 453-463.	0.7	27

ELEAZAR SOTO

#	Article	IF	CITATIONS
19	Unexplained intrauterine fetal death is accompanied by activation of complement. Journal of Perinatal Medicine, 2005, 33, 296-305.	0.6	26
20	Increased concentration of the complement split product C5a in acute pyelonephritis during pregnancy. Journal of Maternal-Fetal and Neonatal Medicine, 2005, 17, 247-252.	0.7	24
21	Blood pH and gases in fetuses in preterm labor with and without systemic inflammatory response syndrome. Journal of Maternal-Fetal and Neonatal Medicine, 2012, 25, 1160-1170.	0.7	21
22	Maternal serum soluble CD30 is increased in pregnancies complicated with acute pyelonephritis. Journal of Maternal-Fetal and Neonatal Medicine, 2007, 20, 803-811.	0.7	17
23	Fragment Bb: evidence for activation of the alternative pathway of the complement system in pregnant women with acute pyelonephritis. Journal of Maternal-Fetal and Neonatal Medicine, 2010, 23, 1085-1090.	0.7	17
24	Pyelonephritis during pregnancy: A cause for an acquired deficiency of protein Z. Journal of Maternal-Fetal and Neonatal Medicine, 2008, 21, 629-637.	0.7	13
25	von Hippel-Lindau Disease and Pregnancy. Obstetrical and Gynecological Survey, 2013, 68, 655-662.	0.2	11
26	Fetal Abnormal Growth Associated With Substance Abuse. Clinical Obstetrics and Gynecology, 2013, 56, 142-153.	0.6	11
27	Normal pregnancy is characterized by systemic activation of the complement system. American Journal of Obstetrics and Gynecology, 2004, 191, S139.	0.7	3
28	Decreased MRNA expression of complement regulatory proteins inÂchorioamnionitis. American Journal of Obstetrics and Gynecology, 2006, 195, S71.	0.7	2
29	Evidence for complement activation in premature labor associated with intra-amniotic infection. American Journal of Obstetrics and Gynecology, 2006, 195, S74.	0.7	2