

Eleazar Soto

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11965159/publications.pdf>

Version: 2024-02-01

29
papers

1,327
citations

331259

21
h-index

476904

29
g-index

29
all docs

29
docs citations

29
times ranked

1267
citing authors

#	ARTICLE	IF	CITATIONS
1	Fetal Anomalies and Long-Term Effects Associated with Substance Abuse in Pregnancy: A Literature Review. <i>American Journal of Perinatology</i> , 2015, 32, 405-416.	0.6	35
2	The diagnostic performance of the Mass Restricted (MR) score in the identification of microbial invasion of the amniotic cavity or intra-amniotic inflammation is not superior to amniotic fluid interleukin-6. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2014, 27, 757-769.	0.7	44
3	Maternal plasma concentrations of angiogenic/antiangiogenic factors in the third trimester of pregnancy to identify the patient at risk for stillbirth at or near term and severe late preeclampsia. <i>American Journal of Obstetrics and Gynecology</i> , 2013, 208, 287.e1-287.e15.	0.7	122
4	von Hippel-Lindau Disease and Pregnancy. <i>Obstetrical and Gynecological Survey</i> , 2013, 68, 655-662.	0.2	11
5	Fetal Abnormal Growth Associated With Substance Abuse. <i>Clinical Obstetrics and Gynecology</i> , 2013, 56, 142-153.	0.6	11
6	Blood pH and gases in fetuses in preterm labor with and without systemic inflammatory response syndrome. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2012, 25, 1160-1170.	0.7	21
7	Late-onset preeclampsia is associated with an imbalance of angiogenic and anti-angiogenic factors in patients with and without placental lesions consistent with maternal underperfusion. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2012, 25, 498-507.	0.7	136
8	Maternal plasma concentrations of angiogenic/anti-angiogenic factors are of prognostic value in patients presenting to the obstetrical triage area with the suspicion of preeclampsia. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2011, 24, 1187-1207.	0.7	118
9	ORIGINAL ARTICLE: Activation of the Alternative Pathway of Complement is a Feature of Preterm Parturition but not of Spontaneous Labor at Term. <i>American Journal of Reproductive Immunology</i> , 2010, 63, 318-330.	1.2	32
10	Fragment Bb: evidence for activation of the alternative pathway of the complement system in pregnant women with acute pyelonephritis. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2010, 23, 1085-1090.	0.7	17
11	Preeclampsia and pregnancies with small-for-gestational age neonates have different profiles of complement split products. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2010, 23, 646-657.	0.7	48
12	Fragment Bb in amniotic fluid: evidence for complement activation by the alternative pathway in women with intra-amniotic infection/inflammation. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2009, 22, 905-916.	0.7	33
13	Evidence for complement activation in the amniotic fluid of women with spontaneous preterm labor and intra-amniotic infection. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2009, 22, 983-992.	0.7	53
14	Twin-to-twin transfusion syndrome: an antiangiogenic state?. <i>American Journal of Obstetrics and Gynecology</i> , 2008, 198, 382.e1-382.e8.	0.7	51
15	Pyelonephritis during pregnancy: A cause for an acquired deficiency of protein Z. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2008, 21, 629-637.	0.7	13
16	Plasma protein Z concentrations in pregnant women with idiopathic intrauterine bleeding and in women with spontaneous preterm labor. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2007, 20, 453-463.	0.7	27
17	Human β 2-defensin-2: A natural antimicrobial peptide present in amniotic fluid participates in the host response to microbial invasion of the amniotic cavity. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2007, 20, 15-22.	0.7	80
18	Maternal serum soluble CD30 is increased in normal pregnancy, but decreased in preeclampsia and small for gestational age pregnancies. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2007, 20, 867-878.	0.7	34

#	ARTICLE	IF	CITATIONS
19	Maternal serum soluble CD30 is increased in pregnancies complicated with acute pyelonephritis. Journal of Maternal-Fetal and Neonatal Medicine, 2007, 20, 803-811.	0.7	17
20	Objective Evaluation of Sylvian Fissure Development by Multiplanar 3-Dimensional Ultrasonography. Journal of Ultrasound in Medicine, 2007, 26, 347-353.	0.8	28
21	What Does 2â€­Dimensional Imaging Add to 3â€­and 4â€­Dimensional Obstetric Ultrasonography?. Journal of Ultrasound in Medicine, 2006, 25, 691-699.	0.8	44
22	A rapid MMP-8 bedside test for the detection of intra-amniotic inflammation identifies patients at risk for imminent preterm delivery. American Journal of Obstetrics and Gynecology, 2006, 195, 1025-1030.	0.7	87
23	Decreased mRNA expression of complement regulatory proteins in chorioamnionitis. American Journal of Obstetrics and Gynecology, 2006, 195, S71.	0.7	2
24	Evidence for complement activation in premature labor associated with intra-amniotic infection. American Journal of Obstetrics and Gynecology, 2006, 195, S74.	0.7	2
25	Unexplained intrauterine fetal death is accompanied by activation of complement. Journal of Perinatal Medicine, 2005, 33, 296-305.	0.6	26
26	Anaphylatoxins in preterm and term labor. Journal of Perinatal Medicine, 2005, 33, 306-13.	0.6	36
27	Normal pregnancy is characterized by systemic activation of the complement system. Journal of Maternal-Fetal and Neonatal Medicine, 2005, 17, 239-245.	0.7	172
28	Increased concentration of the complement split product C5a in acute pyelonephritis during pregnancy. Journal of Maternal-Fetal and Neonatal Medicine, 2005, 17, 247-252.	0.7	24
29	Normal pregnancy is characterized by systemic activation of the complement system. American Journal of Obstetrics and Gynecology, 2004, 191, S139.	0.7	3