

# Premini Mahendra

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11972098/publications.pdf>

Version: 2024-02-01

13  
papers

2,679  
citations

759233

12  
h-index

1199594

12  
g-index

13  
all docs

13  
docs citations

13  
times ranked

2208  
citing authors

#	ARTICLE	IF	CITATIONS
1	CD8+ T-cell immunity against cancer-testis antigens develops following allogeneic stem cell transplantation and reveals a potential mechanism for the graft-versus-leukemia effect. <i>Haematologica</i> , 2010, 95, 1572-1578.	3.5	16
2	Reduced-intensity conditioning for allogeneic haematopoietic stem cell transplantation in relapsed and refractory Hodgkin lymphoma: impact of alemtuzumab and donor lymphocyte infusions on long-term outcomes. <i>British Journal of Haematology</i> , 2007, 139, 70-80.	2.5	104
3	Donor KIR genotype has a major influence on the rate of cytomegalovirus reactivation following T-cell replete stem cell transplantation. <i>Blood</i> , 2005, 107, 1230-1232.	1.4	155
4	Allogeneic Stem-Cell Transplantation Using a Reduced-Intensity Conditioning Regimen Has the Capacity to Produce Durable Remissions and Long-Term Disease-Free Survival in Patients With High-Risk Acute Myeloid Leukemia and Myelodysplasia. <i>Journal of Clinical Oncology</i> , 2005, 23, 9387-9393.	1.6	223
5	Adoptive transfer of cytomegalovirus-specific CTL to stem cell transplant patients after selection by HLA-peptide tetramers. <i>Journal of Experimental Medicine</i> , 2005, 202, 379-386.	8.5	466
6	Clinical evidence of a graft-versus-Hodgkin's-lymphoma effect after reduced-intensity allogeneic transplantation. <i>Lancet</i> , The, 2005, 365, 1934-1941.	13.7	273
7	Donor KIR Genotype Does Not Affect VZV Reactivation after Allogeneic Haematopoietic Stem Cell Transplantation.. <i>Blood</i> , 2005, 106, 3236-3236.	1.4	0
8	Chronic graft-versus-host disease is associated with increased numbers of peripheral blood CD4+CD25high regulatory T cells. <i>Blood</i> , 2004, 103, 2410-2416.	1.4	196
9	Origin and subset distribution of peripheral blood dendritic cells in patients with chronic graft-versus-host disease1. <i>Transplantation</i> , 2003, 75, 221-225.	1.0	49
10	High incidence of cytomegalovirus infection after nonmyeloablative stem cell transplantation: potential role of Campath-1H in delaying immune reconstitution. <i>Blood</i> , 2002, 99, 4357-4363.	1.4	349
11	Limiting transplantation-related mortality following unrelated donor stem cell transplantation by using a nonmyeloablative conditioning regimen. <i>Blood</i> , 2002, 99, 1071-1078.	1.4	333
12	In vivo CAMPATH-1H prevents graft-versus-host disease following nonmyeloablative stem cell transplantation. <i>Blood</i> , 2000, 96, 2419-2425.	1.4	483
13	An Unusual Presentation of Acute Myeloid Leukemia with Pericardial and Pleural Effusions Due to Granulocytic Sarcoma. <i>Leukemia and Lymphoma</i> , 1993, 11, 305-307.	1.3	32