

Solange M Gennari

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1197233/publications.pdf>

Version: 2024-02-01

252
papers

8,021
citations

70961

41
h-index

74018

75
g-index

261
all docs

261
docs citations

261
times ranked

4969
citing authors

#	ARTICLE	IF	CITATIONS
1	Rickettsia Species Infecting Amblyomma cooperi Ticks from an Area in the State of São Paulo, Brazil, Where Brazilian Spotted Fever Is Endemic. Journal of Clinical Microbiology, 2004, 42, 90-98.	1.8	522
2	Toxoplasmosis in humans and animals in Brazil: high prevalence, high burden of disease, and epidemiology. Parasitology, 2012, 139, 1375-1424.	0.7	399
3	Geographical patterns of <i>Toxoplasma gondii</i> genetic diversity revealed by multilocus PCR-RFLP genotyping. Parasitology, 2014, 141, 453-461.	0.7	346
4	Population structure and mouse-virulence of <i>Toxoplasma gondii</i> in Brazil. International Journal for Parasitology, 2008, 38, 561-569.	1.3	310
5	Rickettsial Infection in Animals and Brazilian Spotted Fever Endemicity. Emerging Infectious Diseases, 2005, 11, 265-270.	2.0	309
6	Biological and genetic characterisation of <i>Toxoplasma gondii</i> isolates from chickens (Gallus Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 547 T 32, 99-105.	1.3	248
7	Local admixture of amplified and diversified secreted pathogenesis determinants shapes mosaic <i>Toxoplasma gondii</i> genomes. Nature Communications, 2016, 7, 10147.	5.8	243
8	Prevalence of <i>Rickettsia</i> Infection in Dogs from the Urban and Rural Areas of Monte Negro Municipality, Western Amazon, Brazil. Vector-Borne and Zoonotic Diseases, 2007, 7, 249-255.	0.6	138
9	Seasonal dynamics of ticks (Acari: Ixodidae) on horses in the state of São Paulo, Brazil. Veterinary Parasitology, 2002, 105, 65-77.	0.7	136
10	PREVALENCE OF ANTIBODIES TO SPOTTED FEVER GROUP RICKETTSIAE IN HUMANS AND DOMESTIC ANIMALS IN A BRAZILIAN SPOTTED FEVER-ENDEMIC AREA IN THE STATE OF SÃO PAULO, BRAZIL: SEROLOGIC EVIDENCE FOR INFECTION BY RICKETTSIA RICKETTSII AND ANOTHER SPOTTED FEVER GROUP RICKETTSIA. American Journal of Tropical Medicine and Hygiene, 2004, 71, 93-97.	0.6	131
11	Risk factors to tick infestations and their occurrence on horses in the state of São Paulo, Brazil. Veterinary Parasitology, 2001, 97, 1-14.	0.7	127
12	Molecular identification of <i>Giardia duodenalis</i> isolates from humans, dogs, cats and cattle from the state of São Paulo, Brazil, by sequence analysis of fragments of glutamate dehydrogenase (gdh) coding gene. Veterinary Parasitology, 2007, 149, 258-264.	0.7	109
13	The ROP18 and ROP5 gene allele types are highly predictive of virulence in mice across globally distributed strains of <i>Toxoplasma gondii</i> . International Journal for Parasitology, 2016, 46, 141-146.	1.3	103
14	Ticks and rickettsial infection in the wildlife of two regions of the Brazilian Amazon. Experimental and Applied Acarology, 2015, 65, 125-140.	0.7	100
15	<i>Toxoplasma gondii</i> infection in cats from São Paulo state, Brazil: Seroprevalence, oocyst shedding, isolation in mice, and biologic and molecular characterization. Research in Veterinary Science, 2006, 81, 58-67.	0.9	98
16	<i>Toxoplasma gondii</i> : Detection by mouse bioassay, histopathology, and polymerase chain reaction in tissues from experimentally infected pigs. Experimental Parasitology, 2006, 113, 267-271.	0.5	75
17	Clinical Toxoplasmosis in Dogs and Cats: An Update. Frontiers in Veterinary Science, 2019, 6, 54.	0.9	73
18	Prevalence of <i>Ehrlichia canis</i> (Rickettsiales: Anaplasmataceae) in Dogs and <i>Rhipicephalus sanguineus</i> (Acari: Ixodidae) Ticks from Brazil. Journal of Medical Entomology, 2007, 44, 126-132.	0.9	72

#	ARTICLE	IF	CITATIONS
19	Wild canids, domestic dogs and their pathogens in Southeast Brazil: disease threats for canid conservation. <i>Biodiversity and Conservation</i> , 2010, 19, 3513-3524.	1.2	65
20	Study of the Seasonal Dynamics, Life Cycle, and Host Specificity of <i>Amblyomma aureolatum</i> (Acari: Ixodidae). <i>Journal of Medical Entomology</i> , 2004, 41, 324-332.	0.9	64
21	CHARACTERIZATION OF TOXOPLASMA GONDII ISOLATES IN FREE-RANGE CHICKENS FROM AMAZON, BRAZIL. <i>Journal of Parasitology</i> , 2006, 92, 36-40.	0.3	64
22	Partial protection against tissue cysts formation in pigs vaccinated with crude rhoptry proteins of <i>Toxoplasma gondii</i> . <i>Veterinary Parasitology</i> , 2005, 129, 209-217.	0.7	63
23	Prevalence of <i>Sarcocystis neurona</i> and <i>Neospora</i> spp. infection in horses from Brazil based on presence of serum antibodies to parasite surface antigen. <i>Veterinary Parasitology</i> , 2006, 136, 155-159.	0.7	62
24	Phylogenetic relationships of <i>Leishmania</i> species based on trypanosomatid barcode (SSU rDNA) and gGAPDH genes: Taxonomic revision of <i>Leishmania</i> (L.) <i>infantum chagasi</i> in South America. <i>Infection, Genetics and Evolution</i> , 2014, 25, 44-51.	1.0	61
25	Prevalence of anti- <i>Toxoplasma gondii</i> and anti- <i>Neospora caninum</i> antibodies in goat from São Paulo State, Brazil. <i>Small Ruminant Research</i> , 2004, 55, 29-32.	0.6	58
26	Seroepidemiology of <i>Toxoplasma gondii</i> and <i>Neospora caninum</i> in dogs from the state of Paraíba, Northeast region of Brazil. <i>Research in Veterinary Science</i> , 2005, 79, 51-56.	0.9	56
27	<i>Toxoplasma gondii</i> : Comparison of a rhoptry-ELISA with IFAT and MAT for antibody detection in sera of experimentally infected pigs. <i>Experimental Parasitology</i> , 2006, 113, 100-105.	0.5	55
28	Prevalence of anti- <i>Toxoplasma gondii</i> and anti- <i>Neospora caninum</i> antibodies in sheep from Mossoró, Rio Grande do Norte, Brazil. <i>Veterinary Parasitology</i> , 2009, 160, 211-214.	0.7	55
29	Genetic diversity among capybara (<i>Hydrochaeris hydrochaeris</i>) isolates of <i>Toxoplasma gondii</i> from Brazil. <i>Veterinary Parasitology</i> , 2009, 162, 332-337.	0.7	55
30	Analysis of marine bivalve shellfish from the fish market in Santos city, São Paulo state, Brazil, for <i>Toxoplasma gondii</i> . <i>Veterinary Parasitology</i> , 2010, 170, 8-13.	0.7	54
31	Genetic characterization of <i>Toxoplasma gondii</i> from Brazilian wildlife revealed abundant new genotypes. <i>International Journal for Parasitology: Parasites and Wildlife</i> , 2014, 3, 276-283.	0.6	53
32	Prevalence of <i>Ehrlichia canis</i> (Rickettsiales: Anaplasmataceae) in Dogs and <i>Rhipicephalus sanguineus</i> (Acari: Ixodidae) Ticks from Brazil. <i>Journal of Medical Entomology</i> , 2007, 44, 126-132.	0.9	51
33	BIOLOGICAL STUDIES AND MOLECULAR CHARACTERIZATION OF A CRYPTOSPORIDIUM ISOLATE FROM OSTRICHES (<i>STRUTHIO CAMELUS</i>). <i>Journal of Parasitology</i> , 2006, 92, 623-626.	0.3	50
34	Seroprevalence of <i>Toxoplasma gondii</i> in captive neotropical felids from Brazil. <i>Veterinary Parasitology</i> , 2001, 102, 217-224.	0.7	49
35	Prevalence of anti- <i>Toxoplasma gondii</i> and anti- <i>Neospora caninum</i> antibodies in goats slaughtered in the public slaughterhouse of Patos city, Paraíba State, Northeast region of Brazil. <i>Veterinary Parasitology</i> , 2007, 149, 126-129.	0.7	48
36	Worm burden and immunological responses in Corriedale and Crioula Lanada sheep following natural infection with <i>Haemonchus contortus</i> . <i>Small Ruminant Research</i> , 2004, 51, 75-83.	0.6	47

#	ARTICLE	IF	CITATIONS
37	Prevalence of anti-Neospora caninum antibodies in cattle and dogs from Western Amazon, Brazil, in association with some possible risk factors. <i>Veterinary Parasitology</i> , 2006, 142, 71-77.	0.7	46
38	Natural infection with zoonotic subtype of <i>Cryptosporidium parvum</i> in Capybara (<i>Hydrochoerus</i>) Tj ETQq0 0 0 rgBT/Overlock, 10 Tf 50 7	0.7	46
39	Prevalence of <i>Toxoplasma gondii</i> and <i>Neospora caninum</i> infections in sheep from Federal District, central region of Brazil. <i>Tropical Animal Health and Production</i> , 2009, 41, 547-552.	0.5	45
40	Genetic diversity among <i>Toxoplasma gondii</i> isolates of small ruminants from Brazil: Novel genotypes revealed. <i>Veterinary Parasitology</i> , 2010, 170, 307-312.	0.7	45
41	All about neosporosis in Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2017, 26, 253-279.	0.2	45
42	Ocorrência de protozoários e helmintos em amostras de fezes de cães e gatos da cidade de São Paulo. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 1999, 36, 0-0.	0.2	44
43	Prevalence of equine Piroplasmosis and its association with tick infestation in the State of São Paulo, Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2009, 18, 1-8.	0.2	44
44	<i>Candidatus Rickettsia andeanae</i> , a spotted fever group agent infecting <i>Amblyomma parvum</i> ticks in two Brazilian biomes. <i>Memorias Do Instituto Oswaldo Cruz</i> , 2014, 109, 259-261.	0.8	43
45	Seroprevalence of <i>Neospora caninum</i> in female water buffaloes (<i>Bubalus bubalis</i>) from the southeastern region of Brazil. <i>Veterinary Parasitology</i> , 2001, 99, 331-334.	0.7	42
46	First isolation and molecular characterization of <i>Toxoplasma gondii</i> from finishing pigs from São Paulo State, Brazil. <i>Veterinary Parasitology</i> , 2005, 131, 207-211.	0.7	39
47	Occurrence of <i>Neospora caninum</i> antibodies in sera from dogs of the city of São Paulo, Brazil. <i>Veterinary Parasitology</i> , 2002, 106, 177-179.	0.7	37
48	Life-cycle and host specificity of <i>Amblyomma tigrinum</i> (Acari: Ixodidae) under laboratory conditions. <i>Experimental and Applied Acarology</i> , 2002, 26, 115-125.	0.7	35
49	Occurrence of anti- <i>Neospora caninum</i> antibodies in water buffaloes (<i>Bubalus bubalis</i>) from the Northern region of Brazil. <i>Veterinary Parasitology</i> , 2005, 134, 169-171.	0.7	35
50	Ticks, rickettsial and erlichial infection in small mammals from Atlantic forest remnants in northeastern Brazil. <i>International Journal for Parasitology: Parasites and Wildlife</i> , 2018, 7, 380-385.	0.6	35
51	Freqüência de parasitos gastrintestinais em cães e gatos atendidos em hospital-escola veterinário da cidade de São Paulo. <i>Arquivo Brasileiro De Medicina Veterinaria E Zootecnia</i> , 2007, 59, 1338-1340.	0.1	34
52	Effect of <i>Acacia molissima</i> tannin extract on the control of gastrointestinal parasites in sheep. <i>Animal Feed Science and Technology</i> , 2008, 147, 172-181.	1.1	34
53	Prevalência de anticorpos contra agentes virais e bacterianos em eqüídeos do Município de Monte Negro, Rondônia, Amazônia Ocidental Brasileira: Brazilian Western Amazon. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2008, 45, 269.	0.2	34
54	Epidemiological aspects of Visceral Larva Migrants in children living at São Remo Community, São Paulo (SP), Brazil. <i>Veterinary Parasitology</i> , 2005, 134, 93-97.	0.7	33

#	ARTICLE	IF	CITATIONS
55	Prevalence and risk factors for anti-Toxoplasma gondii antibodies in goats of the Serid�3 Oriental microregion, Rio Grande do Norte state, Northeast region of Brazil. <i>Veterinary Parasitology</i> , 2008, 156, 329-332.	0.7	33
56	Genotyping of potentially zoonotic Giardia duodenalis from exotic and wild animals kept in captivity in Brazil. <i>Veterinary Parasitology</i> , 2011, 180, 344-348.	0.7	33
57	<i>Bartonella</i> infection in urban and rural dogs from the tropics: Brazil, Colombia, Sri Lanka and Vietnam. <i>Epidemiology and Infection</i> , 2013, 141, 54-61.	1.0	33
58	Prevalence of antibodies to spotted fever group rickettsiae in humans and domestic animals in a Brazilian spotted fever-endemic area in the state of S�o Paulo, Brazil: serologic evidence for infection by Rickettsia rickettsii and another spotted fever group Rickettsia. <i>American Journal of Tropical Medicine and Hygiene</i> , 2004, 71, 93-7.	0.6	33
59	Occurrence of Anti-Toxoplasma gondii Antibodies in Dogs in the Urban Area of Monte Negro, Rond�nia, Brazil. <i>Veterinary Research Communications</i> , 2004, 28, 113-118.	0.6	32
60	Novel piroplasmid and Hepatozoon organisms infecting the wildlife of two regions of the Brazilian Amazon. <i>International Journal for Parasitology: Parasites and Wildlife</i> , 2017, 6, 115-121.	0.6	32
61	Antibodies to Neospora caninum and Toxoplasma gondii in domestic cats from Brazil. <i>Parasitology Research</i> , 2007, 100, 281-285.	0.6	31
62	Seroprevalence of Toxoplasma gondii Antibodies in Captive Wild Mammals and Birds in Brazil. <i>Journal of Zoo and Wildlife Medicine</i> , 2010, 41, 572-574.	0.3	31
63	Genotyping of Toxoplasma gondii isolates from free range chickens in the Pantanal area of Brazil. <i>Veterinary Parasitology</i> , 2011, 178, 29-34.	0.7	31
64	Ocorr�ncia de parasitos gastrintestinais em fezes de gatos das cidades de S�o Paulo e Guarulhos. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2002, 39, 244-246.	0.2	30
65	Association between the presence of serum antibodies against Neospora spp. and fetal loss in equines. <i>Veterinary Parasitology</i> , 2006, 142, 372-375.	0.7	30
66	Molecular identification of Cryptosporidium spp. from fecal samples of felines, canines and bovines in the state of S�o Paulo, Brazil. <i>Veterinary Parasitology</i> , 2007, 150, 291-296.	0.7	30
67	Occurrence of antibodies anti-Neospora caninum, anti-Toxoplasma gondii, and anti-Leishmania chagasi in serum of dogs from Par� State, Amazon, Brazil. <i>Parasitology Research</i> , 2010, 107, 453-457.	0.6	30
68	Prevalence of anti-Toxoplasma gondii and anti-Neospora caninum antibodies in swine from Northeastern Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2010, 19, 80-84.	0.2	30
69	Molecular phylogenetic analysis in Hammondia-like organisms based on partial Hsp70 coding sequences. <i>Parasitology</i> , 2007, 134, 1195-1203.	0.7	29
70	Detection of antibodies to Neospora caninum in two species of wild canids, Lycalopex gymnocercus and Cerdocyon thous from Brazil. <i>Veterinary Parasitology</i> , 2004, 123, 275-277.	0.7	28
71	Occurrence of antibodies against Toxoplasma gondii and its isolation and genotyping in donkeys, mules, and horses in Brazil. <i>Veterinary Parasitology</i> , 2015, 209, 129-132.	0.7	28
72	Preval�ncia de carrapatos em c�es de �reas rurais da regi�o norte do Estado do Paran�. <i>Arquivo Brasileiro De Medicina Veterinaria E Zootecnia</i> , 2001, 53, 553-556.	0.1	28

#	ARTICLE	IF	CITATIONS
73	Rickettsial spotted fever in capoeirã Village, Itabira, Minas Gerais, Brazil. <i>Revista Do Instituto De Medicina Tropical De Sao Paulo</i> , 2008, 50, 297-301.	0.5	27
74	Prevalence of Antibodies to <i>Trypanosoma cruzi</i> , <i>Leishmania infantum</i> , <i>Encephalitozoon cuniculi</i> , <i>Sarcocystis neurona</i> , and <i>Neospora caninum</i> in Capybara, <i>Hydrochoerus hydrochaeris</i> , from Sã Paulo State, Brazil. <i>Journal of Parasitology</i> , 2010, 96, 521-524.	0.3	27
75	Isolation of <i>Toxoplasma gondii</i> from Capybaras (<i>Hydrochaeris hydrochaeris</i>) from Sã Paulo State, Brazil. <i>Journal of Parasitology</i> , 2008, 94, 1060-1063.	0.3	26
76	Prevalence of <i>Neospora caninum</i> antibodies in cattle from Santarãm, Parã, Brazil. <i>Research in Veterinary Science</i> , 2008, 84, 254-256.	0.9	25
77	Inquãrito sorolãgico para toxoplasmose e leptospirose em mamãferos selvagens neotropicais do Zoolãgico de Aracaju, Sergipe. <i>Pesquisa Veterinaria Brasileira</i> , 2009, 29, 1009-1014.	0.5	25
78	A Survey of <i>Neospora caninum</i> and <i>Toxoplasma gondii</i> Infection in Urban Rodents from Brazil. <i>Journal of Parasitology</i> , 2012, 98, 128-134.	0.3	25
79	Occurrences of gastrointestinal parasites in fecal samples from domestic dogs in Sã Paulo, SP, Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2016, 25, 435-440.	0.2	25
80	<i>Toxoplasma gondii</i> : isolation of tachyzoites rhoptries and incorporation into Iscom. <i>Experimental Parasitology</i> , 2004, 108, 40-46.	0.5	24
81	Transplacental transmission of <i>Toxoplasma gondii</i> in reinfected pregnant female canines. <i>Parasitology Research</i> , 2009, 104, 1213-1217.	0.6	24
82	Shedding of <i>Neospora caninum</i> oocysts by dogs fed different tissues from naturally infected cattle. <i>Veterinary Parasitology</i> , 2011, 179, 220-223.	0.7	24
83	Occurrence of Antibodies to <i>Toxoplasma gondii</i> and <i>Lepstospira</i> spp. in Manatees (<i>Trichechus inunguis</i>) of the Brazilian Amazon. <i>Journal of Zoo and Wildlife Medicine</i> , 2012, 43, 85-88.	0.3	24
84	Isolation and Phylogenetic Relationships of Bat Trypanosomes from Different Biomes in Mato Grosso, Brazil. <i>Journal of Parasitology</i> , 2013, 99, 1071-1076.	0.3	24
85	<i>Trypanosoma cruzi</i> and <i>Leishmania infantum chagasi</i> Infection in Wild Mammals from Maranhão State, Brazil. <i>Vector-Borne and Zoonotic Diseases</i> , 2015, 15, 656-666.	0.6	24
86	<i>Rickettsia felis</i> (Rickettsiales: Rickettsiaceae) in <i>Ctenocephalides felis felis</i> (Siphonaptera: Pulicidae) in the State of Sã Paulo, Brazil. <i>Arquivo Brasileiro De Medicina Veterinaria E Zootecnia</i> , 2005, 57, 321-325.	0.1	23
87	Fatores de risco associados à ocorrãncia de anticorpos anti- <i>Leptospira</i> spp. em cães do municãpio de Monte Negro, Rondãnia, Amazãnia Ocidental Brasileira. <i>Arquivo Brasileiro De Medicina Veterinaria E Zootecnia</i> , 2007, 59, 70-76.	0.1	23
88	Evidence of congenital transmission of <i>Neospora caninum</i> in naturally infected water buffalo (<i>Bubalus bubalis</i>) fetus from Brazil. <i>Parasitology Research</i> , 2011, 108, 741-743.	0.6	23
89	Evaluation of IFA, MAT, ELISAs and immunoblotting for the detection of anti- <i>Toxoplasma gondii</i> antibodies in paired serum and aqueous humour samples from experimentally infected pigs. <i>Research in Veterinary Science</i> , 2008, 84, 237-242.	0.9	22
90	Serosurvey for tick-borne diseases in dogs from the Eastern Amazon, Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2013, 22, 214-219.	0.2	22

#	ARTICLE	IF	CITATIONS
91	Evaluation of experimental <i>Toxoplasma gondii</i> (Nicolle and Manceaux, 1909) infection in pigs by bioassay in mice and polymerase chain reaction. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2003, 40, .	0.2	22
92	Seroprevalence of toxoplasmosis in a low-income community in the São Paulo municipality, SP, Brazil. <i>Revista Do Instituto De Medicina Tropical De Sao Paulo</i> , 2006, 48, 167-170.	0.5	21
93	Seroprevalence of <i>Leptospira</i> spp in cattle from Monte Negro municipality, western Amazon. <i>Pesquisa Veterinaria Brasileira</i> , 2006, 26, 102-104.	0.5	21
94	Prevalência de anticorpos anti- <i>Toxoplasma gondii</i> e anti- <i>Neospora caninum</i> em rebanhos caprinos do município de Mossoró ³ , Rio Grande do Norte. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2008, 45, 81.	0.2	21
95	<i>Toxoplasma gondii</i> in <i>Capybara</i> (<i>Hydrochaeris hydrochaeris</i>) antibodies and DNA detected by IFAT and PCR. <i>Parasitology Research</i> , 2010, 107, 141-146.	0.6	21
96	Pathogenicity of Nc-Bahia and Nc-1 strains of <i>Neospora caninum</i> in experimentally infected cows and buffaloes in early pregnancy. <i>Parasitology Research</i> , 2014, 113, 1521-1528.	0.6	21
97	Novel <i>Anaplasma</i> and <i>Ehrlichia</i> organisms infecting the wildlife of two regions of the Brazilian Amazon. <i>Acta Tropica</i> , 2017, 174, 82-87.	0.9	21
98	High genetic diversity in <i>Toxoplasma gondii</i> isolates from pigs at slaughterhouses in Paráiba state, northeastern Brazil: Circulation of new genotypes and Brazilian clonal lineages. <i>Veterinary Parasitology</i> , 2017, 244, 76-80.	0.7	21
99	Serosurvey of Smooth <i>Brucella</i> , <i>Leptospira</i> spp. and <i>Toxoplasma gondii</i> in Free-Ranging Jaguars (<i>Panthera onca</i>) and Domestic Animals from Brazil. <i>PLoS ONE</i> , 2015, 10, e0143816.	1.1	21
100	Seroprevalence of <i>Toxoplasma gondii</i> antibodies from wild canids from Brazil. <i>Veterinary Parasitology</i> , 2004, 121, 337-340.	0.7	20
101	Presença de anticorpos anti- <i>Neospora caninum</i> e <i>Toxoplasma gondii</i> em cães com leishmaniose visceral da região de Araçatuba, São Paulo, Brasil. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2006, 43, 613.	0.2	20
102	Differential diagnosis of oocysts of Hammondia-like organisms of dogs and cats by PCR-RFLP analysis of 70-kilodalton heat shock protein (HSP70) gene. <i>Parasitology Research</i> , 2008, 103, 235-238.	0.6	20
103	Survey of <i>Trypanosoma</i> and <i>Leishmania</i> in Wild and Domestic Animals in an Atlantic Rainforest Fragment and Surroundings in the State of Espírito Santo, Brazil. <i>Journal of Medical Entomology</i> , 2014, 51, 686-693.	0.9	20
104	Presence of antibodies against <i>Toxoplasma gondii</i> , <i>Neospora caninum</i> and <i>Leishmania infantum</i> in dogs from Piauí: <i>Brazilian Journal of Veterinary Parasitology</i> , 2011, 20, 111-114.	0.2	19
105	A longitudinal study of <i>Neospora caninum</i> infection on three dairy farms in Brazil. <i>Veterinary Parasitology</i> , 2012, 187, 553-557.	0.7	19
106	Seroprevalence rates of antibodies against <i>Leishmania infantum</i> and other protozoan and rickettsial parasites in dogs. <i>Brazilian Journal of Veterinary Parasitology</i> , 2013, 22, 162-166.	0.2	19
107	Seroepidemiological survey for brucellosis, leptospirosis, and toxoplasmosis in free-ranging <i>A. louatta caraya</i> and <i>C. allithrix penicillata</i> from São Paulo State, Brazil. <i>Journal of Medical Primatology</i> , 2014, 43, 197-201.	0.3	19
108	Controle estratégico do carrapato <i>Amblyomma cajennense</i> em eqüinos. <i>Ciencia Rural</i> , 2004, 34, 195-200.	0.3	18

#	ARTICLE	IF	CITATIONS
109	Hammondia heydorni: evidence of genetic diversity among isolates from dogs. <i>Experimental Parasitology</i> , 2004, 107, 65-71.	0.5	18
110	Crab-eating fox (<i>Cerdocyon thous</i>), a South American canid, as a definitive host for <i>Hammondia heydorni</i> . <i>Veterinary Parasitology</i> , 2009, 162, 46-50.	0.7	18
111	Experimental infection of dogs (<i>Canis familiaris</i>) with sporulated oocysts of <i>Neospora caninum</i> . <i>Veterinary Parasitology</i> , 2011, 176, 151-156.	0.7	18
112	Survey of canine tick-borne diseases in Lıbrea, Brazilian Amazon: â€™accidentalâ€™™ findings of <i>Dirofilaria immitis</i> infection. <i>Brazilian Journal of Veterinary Parasitology</i> , 2014, 23, 473-480.	0.2	18
113	Presence of neutralizing antibodies to Orthopoxvirus in Capybaras (<i>Hydrochoerus hydrochaeris</i>) in Brazil. <i>Journal of Infection in Developing Countries</i> , 2014, 8, 1646-1649.	0.5	18
114	Abortion and foetal lesions induced by <i>Neospora caninum</i> in experimentally infected water buffalos (<i>Bubalus bubalis</i>). <i>Parasitology Research</i> , 2015, 114, 193-199.	0.6	18
115	Seroprevalence of <i>Toxoplasma gondii</i> in seabirds from Abrolhos Archipelago, Brazil. <i>Veterinary Parasitology</i> , 2016, 226, 50-52.	0.7	18
116	Diversity of bats trypanosomes in hydroelectric area of Belo Monte in Brazilian Amazonia. <i>Acta Tropica</i> , 2016, 164, 185-193.	0.9	18
117	Infectious Diseases in Free-Ranging Blonde Capuchins, <i>Sapajus flavius</i> , in Brazil. <i>International Journal of Primatology</i> , 2017, 38, 1017-1031.	0.9	18
118	<i>Toxoplasma gondii</i> in domestic and wild animals from forest fragments of the municipality of Natal, northeastern Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2014, 23, 501-508.	0.2	17
119	SWINE INFECTIOUS AGENTS IN <i>TAYASSU PECARI</i> AND <i>PECARI TAJACU</i> TISSUE SAMPLES FROM BRAZIL. <i>Journal of Wildlife Diseases</i> , 2014, 50, 205-209.	0.3	17
120	Prevalence and risk factors associated with ectoparasite infestation of buffaloes in an Amazonian ecosystem. <i>Parasites and Vectors</i> , 2018, 11, 335.	1.0	17
121	Vaccination of pigs with <i>Toxoplasma gondii</i> antigens incorporated in immunostimulating complexes (iscoms). <i>Arquivo Brasileiro De Medicina Veterinaria E Zootecnia</i> , 2003, 55, 388-396.	0.1	17
122	Response of Corriedale and Crioula Lanada sheep to artificial primary infection with <i>Haemonchus contortus</i> . <i>Veterinary Research Communications</i> , 2002, 26, 447-457.	0.6	16
123	Occurrence of anti- <i>Neospora caninum</i> antibodies in Brazilian cervids kept in captivity. <i>Veterinary Parasitology</i> , 2005, 129, 341-343.	0.7	16
124	Prevalence of antibodies to <i>Encephalitozoon cuniculi</i> in horses from Brazil. <i>Veterinary Parasitology</i> , 2006, 142, 380-382.	0.7	16
125	Detection of <i>Toxoplasma gondii</i> by PCR and mouse bioassay in commercial cuts of pork from experimentally infected pigs. <i>Arquivo Brasileiro De Medicina Veterinaria E Zootecnia</i> , 2007, 59, 30-34.	0.1	16
126	Seroprevalence of <i>Toxoplasma gondii</i> in Wild Marsupials and Rodents from the Atlantic Forest of Pernambuco State, Northeastern Region, Brazil. <i>Journal of Parasitology</i> , 2013, 99, 1140-1143.	0.3	16

#	ARTICLE	IF	CITATIONS
127	Toxoplasma gondii antibodies in wild rodents and marsupials from the Atlantic Forest, state of São Paulo, Brazil. Brazilian Journal of Veterinary Parasitology, 2015, 24, 379-382.	0.2	16
128	Prevalence of antibodies against Toxoplasma gondii and Neospora spp. in equids of Western Pará, Brazil. Acta Tropica, 2019, 189, 39-45.	0.9	16
129	Constraints to milk production in grazing dairy cows in Brazil and management strategies for improving their productivity. Preventive Veterinary Medicine, 1999, 38, 217-230.	0.7	15
130	Environmental Factors and Ecosystems Associated with Canine Visceral Leishmaniasis in Northeastern Brazil. Vector-Borne and Zoonotic Diseases, 2015, 15, 765-774.	0.6	15
131	Prevalence of antibodies against Neospora spp. and Sarcocystis neurona in donkeys from northeastern Brazil. Brazilian Journal of Veterinary Parasitology, 2016, 25, 109-111.	0.2	15
132	First report of typical Brazilian Toxoplasma gondii genotypes from isolates of free-range chickens (Gallus gallus domesticus) circulating in the state of Paraíba, Northeast Brazil. Parasitology Research, 2017, 116, 2265-2270.	0.6	15
133	Fatal toxoplasmosis in a southern muriqui (<i>Brachyteles arachnoides</i>) from São Paulo state, Brazil: Pathological, immunohistochemical, and molecular characterization. Journal of Medical Primatology, 2018, 47, 124-127.	0.3	15
134	Occurrence of tissue cyst forming coccidia in Magellanic penguins (<i>Spheniscus magellanicus</i>) rescued on the coast of Brazil. PLoS ONE, 2018, 13, e0209007.	1.1	15
135	Toxoplasma gondii in cattle in Brazil: a review. Brazilian Journal of Veterinary Parasitology, 2020, 29, e015719.	0.2	15
136	Environmental effect on the occurrence of anti-Neospora caninum antibodies in pampas-deer (<i>Ozotoceros bezoarticus</i>). Veterinary Parasitology, 2005, 134, 73-76.	0.7	14
137	Experimental infection of pregnant queens with two major Brazilian clonal lineages of Toxoplasma gondii. Parasitology Research, 2009, 105, 1311-6.	0.6	14
138	Toxoplasma gondii: diagnosis of experimental and natural infection in pigeons (<i>Columba livia</i>) by serological, biological and molecular techniques. Brazilian Journal of Veterinary Parasitology, 2010, 19, 237-243.	0.2	14
139	Anuran trypanosomes: phylogenetic evidence for new clades in Brazil. Systematic Parasitology, 2015, 91, 63-70.	0.5	14
140	Frequency of gastrointestinal parasites in cats seen at the University of São Paulo Veterinary Hospital, Brazil. Brazilian Journal of Veterinary Parasitology, 2016, 25, 423-428.	0.2	14
141	Fatal toxoplasmosis in an immunosuppressed domestic cat from Brazil caused by Toxoplasma gondii clonal type I. Brazilian Journal of Veterinary Parasitology, 2017, 26, 177-184.	0.2	14
142	Cat-rodent Toxoplasma gondii Type II-variant circulation and limited genetic diversity on the Island of Fernando de Noronha, Brazil. Parasites and Vectors, 2017, 10, 220.	1.0	14
143	Novel Ehrlichia sp. detected in Magellanic penguins (<i>Spheniscus magellanicus</i>) and in the seabird tick Ixodes uriae from Magdalena Island, southern Chile. Ticks and Tick-borne Diseases, 2019, 10, 101256.	1.1	14
144	Outbreak of toxoplasmosis in a flock of domestic chickens (<i>Gallus gallus domesticus</i>) and guinea fowl (<i>Numida meleagris</i>). Parasitology Research, 2019, 118, 991-997.	0.6	14

#	ARTICLE	IF	CITATIONS
145	Prevalence of Anti- <i>Neospora caninum</i> and Anti- <i>Toxoplasma gondii</i> Antibodies in Dogs From Two Different Indigenous Communities in the Brazilian Amazon Region. <i>Journal of Parasitology</i> , 2012, 98, 1276-1278.	0.3	13
146	Paracoccidioides brasiliensis infection in dogs from Western Brazilian Amazon. <i>Pesquisa Veterinaria Brasileira</i> , 2012, 32, 649-652.	0.5	13
147	Experimentally Induced Clinical <i>Cystoisospora canis</i> Coccidiosis in Dogs with Prior Natural Patent <i>Cystoisospora ohioensis</i> -like or <i>C. canis</i> Infections. <i>Journal of Parasitology</i> , 2013, 99, 892-895.	0.3	13
148	Epidemiological aspects of <i>Toxoplasma gondii</i> infection in riverside communities in the Southern Brazilian Amazon. <i>Revista Da Sociedade Brasileira De Medicina Tropical</i> , 2015, 48, 301-306.	0.4	13
149	Occurrence of <i>Neospora caninum</i> and <i>Toxoplasma gondii</i> antibodies in dogs from rural properties surrounding a biological reserve, Espirito Santo, Brasil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2016, 25, 536-539.	0.2	13
150	Acute toxoplasmosis in pigs in Brazil caused by <i>Toxoplasma gondii</i> genotype Chinese 1. <i>Parasitology Research</i> , 2016, 115, 2561-2566.	0.6	13
151	Rare case of acute toxoplasmosis in a domestic rabbit (<i>Oryctolagus cuniculus</i>) in Brazil associated with the type Br/II Brazilian clonal lineage of <i>Toxoplasma gondii</i> . <i>Parasitology Research</i> , 2017, 116, 2873-2876.	0.6	13
152	Isolation and genotyping of <i>Toxoplasma gondii</i> in the Midwestern Brazil revealed high genetic diversity and new genotypes. <i>Acta Tropica</i> , 2020, 212, 105681.	0.9	13
153	Multilocus Genotyping of <i>Giardia duodenalis</i> in Mostly Asymptomatic Indigenous People from the Tapirapã Tribe, Brazilian Amazon. <i>Pathogens</i> , 2021, 10, 206.	1.2	13
154	Freqüência de anticorpos anti- <i>Neospora caninum</i> em soros de caprinos do estado de São Paulo e sua relação com o manejo dos animais. <i>Pesquisa Veterinaria Brasileira</i> , 2008, 28, 597-600.	0.5	12
155	Occurrence of <i>Toxoplasma gondii</i> antibodies in birds from the Atlantic Forest, state of São Paulo, Brazil. <i>Veterinary Parasitology</i> , 2014, 200, 193-197.	0.7	12
156	SEROLOGIC EVIDENCE OF <i>TOXOPLASMA GONDII</i> INFECTION IN WILD BIRDS AND MAMMALS FROM SOUTHEAST BRAZIL. <i>Journal of Zoo and Wildlife Medicine</i> , 2014, 45, 197-199.	0.3	12
157	Prevalence and risk factors for <i>Toxoplasma gondii</i> in sheep in the State of Paraíba, Northeastern Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2015, 24, 383-386.	0.2	12
158	Serodiagnosis of visceral and cutaneous leishmaniasis in human and canine populations living in Indigenous Reserves in the Brazilian Amazon Region. <i>Revista Da Sociedade Brasileira De Medicina Tropical</i> , 2017, 50, 61-66.	0.4	12
159	Free-range chickens from Santa Catarina state, southern Brazil, as asymptomatic intermediate hosts for <i>Toxoplasma gondii</i> clonal type I and typical Brazilian genotypes. <i>Veterinary Parasitology: Regional Studies and Reports</i> , 2018, 13, 55-59.	0.3	12
160	Isolation of viable <i>Toxoplasma gondii</i> from organs and Brazilian commercial meat cuts of experimentally infected pigs. <i>Parasitology Research</i> , 2019, 118, 1331-1335.	0.6	12
161	Viability of Sporulated Oocysts of <i>Neospora caninum</i> After Exposure to Different Physical and Chemical Treatments. <i>Journal of Parasitology</i> , 2011, 97, 135-139.	0.3	11
162	Canine distemper virus and <i>Toxoplasma gondii</i> co-infection in dogs with neurological signs. <i>Arquivo Brasileiro De Medicina Veterinaria E Zootecnia</i> , 2012, 64, 221-224.	0.1	11

#	ARTICLE	IF	CITATIONS
163	Morphological and molecular characterization and phylogenetic relationships of a new species of trypanosome in <i>Tapirus terrestris</i> (lowland tapir), <i>Trypanosoma terrestris</i> sp. nov., from Atlantic Rainforest of southeastern Brazil. <i>Parasites and Vectors</i> , 2013, 6, 349.	1.0	11
164	<i>Toxoplasma gondii</i> em pequenos felinos silvestres neotropicais. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2013, 50, 50.	0.2	11
165	Seroepidemiology of <i>Toxoplasma gondii</i> infection in bats from São Paulo city, Brazil. <i>Veterinary Parasitology</i> , 2014, 206, 293-296.	0.7	11
166	Characterization of <i>Toxoplasma gondii</i> isolates from herds of sheep in southern Brazil reveals the archetypal type II genotype and new non-archetypal genotypes. <i>Parasitology International</i> , 2018, 67, 59-63.	0.6	11
167	<i>Amblyomma</i> ticks infesting amphibians and Squamata reptiles from the lower Amazon region, Brazil. <i>Experimental and Applied Acarology</i> , 2018, 75, 399-407.	0.7	11
168	Seroprevalence and risk factors associated with <i>Chlamydophila abortus</i> infection in dairy goats in the Northeast of Brazil. <i>Pesquisa Veterinaria Brasileira</i> , 2012, 32, 1082-1086.	0.5	11
169	Ocorrência de anticorpos anti- <i>Brucella abortus</i> e anti- <i>Brucella canis</i> em cães rurais e urbanos do Município de Monte Negro, Rondônia, Brasil. <i>Ciencia Rural</i> , 2005, 35, 1216-1219.	0.3	11
170	Prevalence of anti- <i>Toxoplasma gondii</i> and anti- <i>Neospora caninum</i> antibodies in swine from Northeastern Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2010, 19, 80-4.	0.2	11
171	Bat trypanosomes from Tapajás-Arapiuns Extractive Reserve in Brazilian Amazon. <i>Brazilian Journal of Veterinary Parasitology</i> , 2017, 26, 152-158.	0.2	10
172	Isolation and genetic characterization of <i>Toxoplasma gondii</i> from free-ranging and captive birds and mammals in Pernambuco state, Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2018, 27, 481-487.	0.2	10
173	Congenital Transmission of <i>Toxoplasma gondii</i> After Experimental Reinfection With Brazilian Typical Strains in Chronically Infected Sheep. <i>Frontiers in Veterinary Science</i> , 2019, 6, 93.	0.9	10
174	Factors associated with the prevalence of antibodies against <i>Brucella abortus</i> in water buffaloes from Santarém, Lower Amazon region, Brazil. <i>Transboundary and Emerging Diseases</i> , 2020, 67, 44-48.	1.3	10
175	Perfil sorológico dos anticorpos colostrais para <i>Neospora caninum</i> em bezerros livres da infecção. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2008, 45, 379.	0.2	9
176	Experimental infection with <i>Neospora caninum</i> in pregnant bitches. <i>Brazilian Journal of Veterinary Parasitology</i> , 2012, 21, 232-236.	0.2	9
177	Occurrence of antibodies against <i>Neospora caninum</i> in wild pigs (<i>Sus scrofa</i>) in the Pantanal, Mato Grosso do Sul, Brazil. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2016, 53, 112.	0.2	9
178	Phenotypic and genotypic characterization of two <i>Toxoplasma gondii</i> isolates in free-range chickens from Uberlândia, Brazil. <i>Epidemiology and Infection</i> , 2016, 144, 1865-1875.	1.0	9
179	Detection of <i>Toxoplasma gondii</i> antibodies in captive non-human primates in the Amazon region, Brazil. <i>Journal of Medical Primatology</i> , 2017, 46, 343-346.	0.3	9
180	Occurrences of antibodies against <i>Toxoplasma gondii</i> , <i>Neospora</i> spp., and <i>Sarcocystis neurona</i> in horses and dogs in the municipality of Pauliceia, São Paulo, Brazil. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2017, 54, 277.	0.2	9

#	ARTICLE	IF	CITATIONS
181	Zoonotic parasites infecting free-living armadillos from Brazil. <i>Transboundary and Emerging Diseases</i> , 2021, 68, 1639-1651.	1.3	9
182	Serologic and molecular diagnostic and bioassay in mice for detection of <i>Toxoplasma gondii</i> in free ranges chickens from Pantanal of Mato Grosso do Sul. <i>Pesquisa Veterinaria Brasileira</i> , 2012, 32, 721-726.	0.5	9
183	Histopathological changes during experimental infections of calves with <i>Cooperia punctata</i> . <i>Journal of Helminthology</i> , 2004, 78, 167-171.	0.4	8
184	Avaliação da ocorrência de anticorpos anti- <i>Toxoplasma gondii</i> , em soros de caprinos do estado de São Paulo, e associação com variáveis epidemiológicas, problemas reprodutivos e riscos à saúde pública. <i>Pesquisa Veterinaria Brasileira</i> , 2008, 28, 606-610.	0.5	8
185	Toxoplasmose canina: aspectos clínicos e patológicos. <i>Seminário Ciências Agrárias</i> , 2009, 29, 189.	0.1	8
186	Evaluation of a Rapid Immunochromatographic Dipstick Test for Detection of Antibodies to <i>Trypanosoma cruzi</i> in Dogs Experimentally Infected with Isolates Obtained from Opossums (<i>Didelphis</i>). <i>Journal of Parasitology</i> , 2011, 97, 140-143.	0.5	8
187	Risk factors associated with <i>Toxoplasma gondii</i> seroprevalence in goats in the State of Paraíba, Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2012, 21, 399-404.	0.2	8
188	Seroprevalence of Tick-Borne Pathogens and Tick Infestation in Dogs from Tapirapó and Karajá Indigenous Communities, Brazil. <i>Vector-Borne and Zoonotic Diseases</i> , 2015, 15, 412-418.	0.6	8
189	SURVEY OF <i>TOXOPLASMA GONDII</i> ANTIBODIES IN MAGELLANIC PENGUINS (<i>SPHENISCUS</i>)	0.3	8
190	Seroprevalence of anti- <i>Toxoplasma gondii</i> and anti- <i>Neospora caninum</i> antibodies in domestic mammals from two distinct regions in the semi-arid region of Northeastern Brazil. <i>Veterinary Parasitology: Regional Studies and Reports</i> , 2016, 5, 14-18.	0.3	8
191	Occurrence of <i>Ehrlichia canis</i> and <i>Hepatozoon canis</i> and probable exposure to <i>Rickettsia amblyommatis</i> in dogs and cats in Natal, RN. <i>Brazilian Journal of Veterinary Parasitology</i> , 2019, 28, 151-156.	0.2	8
192	Occurrence of <i>Toxoplasma gondii</i> Antibodies in <i>Dasyprocta aguti</i> from Brazil: Comparison of Diagnostic Techniques. <i>Journal of Zoo and Wildlife Medicine</i> , 2011, 42, 763-765.	0.3	7
193	SEROLOGIC SURVEY OF INFECTIOUS DISEASES IN POPULATIONS OF MANED WOLF (<i>CHRYSOCYON</i>) ECOLOGICAL STATION, BRAZIL. <i>Journal of Zoo and Wildlife Medicine</i> , 2013, 44, 152-155.	0.3	7
194	Seroprevalence and risk factors for toxoplasmosis and neosporosis in the dog population of Ibiúna, São Paulo, Brazil. <i>Seminário Ciências Agrárias</i> , 2015, 36, 3777.	0.1	7
195	Isolation and biological and molecular characterization of <i>Neospora caninum</i> (NC-SP1) from a naturally infected adult asymptomatic cattle (<i>Bos taurus</i>) in the state of São Paulo, Brazil. <i>Parasitology</i> , 2017, 144, 707-711.	0.7	7
196	Experimental <i>Neospora caninum</i> infection in chickens (<i>Gallus gallus domesticus</i>) with oocysts and tachyzoites of two recent isolates reveals resistance to infection. <i>International Journal for Parasitology</i> , 2018, 48, 117-123.	1.3	7
197	The impact of dry ageing vacuum-packed pork on the viability of <i>Toxoplasma gondii</i> tissue cysts. <i>Food Microbiology</i> , 2020, 86, 103331.	2.1	7
198	Vertical transmission and kinetic of antibodies anti- <i>Neospora caninum</i> in naturally infected lambs in the semiarid region of Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2021, 30, e010621.	0.2	7

#	ARTICLE	IF	CITATIONS
199	Falhas reprodutivas associadas com a presença de do vírus da artrite-encefalite caprina, <i>Toxoplasma gondii</i> e <i>Neospora caninum</i> em caprinos no estado de São Paulo, Brasil. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2012, 49, 67.	0.2	7
200	Ocorrência de anticorpos anti- <i>Toxoplasma gondii</i> em emas (<i>Rhea americana</i>) do Centro de Multiplicação de Animais Silvestres de Mossoró ³ , Rio Grande do Norte. <i>Arquivo Brasileiro De Medicina Veterinária E Zootecnia</i> , 2010, 62, 489-491.	0.1	7
201	<i>Toxoplasma gondii</i> Antibodies in Wild White-Lipped Peccary (<i>Tayassu pecari</i>) From Peru. <i>Journal of Parasitology</i> , 2010, 96, 1232-1232.	0.3	6
202	Presence of anti- <i>Toxoplasma gondii</i> , - <i>Neospora caninum</i> , - <i>Leishmania</i> spp. and - <i>Ehrlichia canis</i> antibodies in free-ranging maned wolves (<i>Chrysocyon brachyurus</i>) in the northeastern region of the state of São Paulo, Brazil. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2016, 53, 243.	0.2	6
203	Ocorrência de anticorpos anti- <i>Toxoplasma gondii</i> , anti- <i>Neospora caninum</i> e anti- <i>Leptospira</i> spp. em gambús (<i>Didelphis</i> spp.) no estado de São Paulo, Brasil. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2016, 53, 1.	0.2	6
204	Pesquisa de anticorpos IgG para <i>Neospora caninum</i> e avaliação dos fatores de risco em ovinos do Estado de Sergipe. <i>Pesquisa Veterinária Brasileira</i> , 2017, 37, 813-819.	0.5	6
205	Detection of anti- <i>Toxoplasma gondii</i> antibodies in small wild mammals from preserved and non-preserved areas in the Caatinga biome, a semi-arid region of Northeast Brazil. <i>Veterinary Parasitology: Regional Studies and Reports</i> , 2018, 14, 75-78.	0.3	6
206	Typical Brazilian genotype of <i>Toxoplasma gondii</i> isolated from a horse destined for human consumption in Europe from a slaughterhouse. <i>Parasitology Research</i> , 2018, 117, 3305-3308.	0.6	6
207	ANTIBODIES AGAINST <i>BRUCELLA ABORTUS</i> AND <i>LEPTOSPIRA</i> SPP. IN CAPTIVE MAMMALS IN THE STATES OF PARÁ AND RIO GRANDE DO NORTE, BRAZIL. <i>Journal of Zoo and Wildlife Medicine</i> , 2018, 49, 355-360.	0.3	6
208	Evaluation of antibodies against <i>Toxoplasma gondii</i> and <i>Leptospira</i> spp. in Magellanic penguins (<i>Spheniscus magellanicus</i>) on Magdalena Island, Chile. <i>Veterinary Parasitology: Regional Studies and Reports</i> , 2019, 16, 100282.	0.3	6
209	A simple method to generate PCR-RFLP typing profiles from DNA sequences in <i>Toxoplasma gondii</i> . <i>Infection, Genetics and Evolution</i> , 2020, 85, 104590.	1.0	6
210	Molecular Characterization of New Haplotype of Genus <i>Sarcocystis</i> in Seabirds from Magdalena Island, Southern Chile. <i>Animals</i> , 2021, 11, 245.	1.0	6
211	Richness of hard ticks (Acari: Ixodidae) from Eastern Brazilian Amazonia, state of Pará, Brazil. <i>International Journal of Acarology</i> , 2021, 47, 159-169.	0.3	6
212	Occurrence and diversity of <i>Sarcocystidae</i> protozoa in muscle and brain tissues of bats from São Paulo state, Brazil. <i>International Journal for Parasitology: Parasites and Wildlife</i> , 2021, 14, 91-96.	0.6	6
213	Evaluation of a PCR Based on Primers to Nc5 Gene for the Detection of <i>Neospora caninum</i> in Brain Tissues of Bovine Aborted Fetuses. <i>Veterinary Research Communications</i> , 2004, 28, 581-585.	0.6	5
214	Flock-level risk factors associated with <i>Neospora caninum</i> seroprevalence in dairy goats in a semiarid region of Northeastern Brazil. <i>Small Ruminant Research</i> , 2013, 112, 239-242.	0.6	5
215	<i>Toxoplasma gondii</i> , <i>Neospora caninum</i> and <i>Leishmania amazonensis</i> antibodies in domestic dogs in the western Brazilian Amazon region. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2016, 53, 1.	0.2	5
216	SEROPREVALENCE OF <i>TOXOPLASMA GONDII</i> IN CAPTIVE ANTILLEAN MANATEE (<i>TRICHECHUS MANATUS</i>)	0.0	5

#	ARTICLE	IF	CITATIONS
217	Longitudinal study of <i>Toxoplasma gondii</i> antibodies in female lambs from Para�ba State, Brazil. <i>Small Ruminant Research</i> , 2020, 188, 106125.	0.6	5
218	Vertical transmission of <i>Toxoplasma gondii</i> in naturally infected ewes in the semiarid region of Brazil. <i>Comparative Immunology, Microbiology and Infectious Diseases</i> , 2021, 74, 101595.	0.7	5
219	<i>Toxoplasma gondii</i> isolated from a Brazilian patient with rare pulmonary toxoplasmosis has a novel genotype and is closely related to Amazonian isolates. <i>Parasitology Research</i> , 2021, 120, 1109-1113.	0.6	5
220	Evaluation of the indirect fluorescent antibody test and modified agglutination test for detection of antibodies against <i>Toxoplasma gondii</i> in experimentally infected pigs. <i>Pesquisa Veterinaria Brasileira</i> , 2004, 24, 199-202.	0.5	5
221	Heartworm (<i>Dirofilaria immitis</i>) disease in a Brazilian oncilla (<i>Leopardus tigrinus</i>). <i>Pesquisa Veterinaria Brasileira</i> , 2009, 29, 474-478.	0.5	5
222	Seroprevalence of anti- <i>Leishmania</i> spp. antibodies in rural dogs from the city of Monte Negro, State of Rond�nia, Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2010, 19, 71-72.	0.2	5
223	Possible role of bovine trophoblast giant cells in transplacental transmission of <i>Neospora caninum</i> in cattle. <i>Brazilian Journal of Veterinary Parasitology</i> , 2007, 16, 21-5.	0.2	5
224	Occurrence of anti- <i>Neospora caninum</i> and anti- <i>Toxoplasma gondii</i> antibodies in dogs with visceral leishmaniasis. <i>Pesquisa Veterinaria Brasileira</i> , 2011, 31, 527-532.	0.5	4
225	Occurrence of antibodies anti- <i>Toxoplasma gondii</i> and anti- <i>Neospora caninum</i> in dogs from Natal, RN, Brazil. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2015, 52, 120.	0.2	4
226	Occurrence of antibodies to <i>Toxoplasma gondii</i> in scavenging black vultures (<i>Coragyps atratus</i>) in Brazil. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2017, 54, 197.	0.2	4
227	Occurrence of anti- <i>Toxoplasma gondii</i> and anti- <i>Neospora caninum</i> antibodies in pigs in the State of Par�, Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2021, 30, e017520.	0.2	4
228	Risk factors for <i>Toxoplasma gondii</i> infection in sheep in the northeastern region of Brazil. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2017, 54, 139.	0.2	4
229	Effect of dietary protein intake on calf resilience to <i>Haemonchus placei</i> infection. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2002, 39, 227-232.	0.2	4
230	Efic�cia anti-helm�ntica de tr�s doses de uma associa�o de pamoato de pyrantel, pamoato de oxantel e praziquantel em gatos naturalmente infectados. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 1997, 34, 284.	0.2	3
231	Anthelmintic effects of condensed tannins on <i>Trichostrongylus colubriformis</i> in experimentally infected sheep. <i>Semina:Ciencias Agrarias</i> , 2010, 31, 1009.	0.1	3
232	Molecular and morphologic characterization of <i>Sarcocystis felis</i> (Apicomplexa: Sarcocystidae) in South American wild felids from Brazil. <i>Veterinary Parasitology</i> , 2016, 217, 15-20.	0.7	3
233	Factors associated with the prevalence of antibodies against <i>Theileria equi</i> in equids of Western Par�, Brazil. <i>Transboundary and Emerging Diseases</i> , 2020, 67, 100-105.	1.3	3
234	From weaning to breeding age: Presence of antibodies to <i>Toxoplasma gondii</i> in sheep in the state of Esp�rito Santo, southeastern Brazil. <i>Veterinary Parasitology: Regional Studies and Reports</i> , 2021, 24, 100531.	0.3	3

#	ARTICLE	IF	CITATIONS
235	Dinâmica de anticorpos durante a gestação em vacas naturalmente infectadas com <i>Neospora caninum</i> de quatro rebanhos leiteiros no Brasil. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2009, 46, 395.	0.2	3
236	IgM e IgG como marcadores da infecção transplacentária por <i>Neospora caninum</i> em fetos bovinos. <i>Pesquisa Veterinária Brasileira</i> , 2010, 30, 551-553.	0.5	3
237	First report on the isolation and genotyping of <i>Toxoplasma gondii</i> strains from free-range chickens in the state of Mato Grosso, Midwestern Brazil. <i>Comparative Immunology, Microbiology and Infectious Diseases</i> , 2022, 80, 101725.	0.7	3
238	NEOSPOORA CANINUM SPECIFIC ANTIBODIES IN FREE-RANGING WHITE-LIPPED PECCARIES (<i>TAYASSU PECARI</i>) FROM THE PERUVIAN AMAZON: DETECTION OF ANTIBODIES IN SERUM AND EVALUATION OF INDIRECT FLUORESCENT ANTIBODY TEST WITH HETEROLOGOUS SECONDARY ANTIBODY. <i>Journal of Zoo and Wildlife Medicine</i> , 2018, 49, 656-661.	0.3	2
239	First isolation and genotyping of <i>Toxoplasma gondii</i> in a free-living giant anteater (<i>Myrmecophaga</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 0.9	0.9	2
240	Níveis de IgG séricos em bezerros experimentalmente infectados pelo <i>Haemonchus placei</i> . <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2002, 39, 107-110.	0.2	2
241	Eficácia e persistência da associação de cipermetrina 4,5 % + DDVP 50% (Ectofarma®) no controle do <i>Boophilus microplus</i> e de larvas de <i>Dermatobia hominis</i> em bovinos leiteiros, mestiços, naturalmente infestados. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 1997, 34, 44.	0.2	2
242	<i>Toxoplasma gondii</i> : avaliação da reação de imunofluorescência indireta usando anticorpo secundário heterólogo em roedores silvestres experimentalmente infectados. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2014, 50, 353.	0.2	1
243	Consumption of animal products and frauds: DNA-based methods for the investigation of authenticity and traceability in dairy and meat-derived products – a review. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2015, 52, 183.	0.2	1
244	Molecular Diagnosis and Prevalence of <i>Trypanosoma vivax</i> (Trypanosomatida: Trypanosomatidae) in Buffaloes and Ectoparasites in the Brazilian Amazon Region. <i>Journal of Medical Entomology</i> , 2021, 58, 403-407.	0.9	1
245	First molecular-based detection study of <i>Leishmania infantum</i> in the Tapirapó indigenous population in the Brazilian Amazon. <i>Brazilian Journal of Medical and Biological Research</i> , 2022, 55, e11654.	0.7	1
246	Serological survey of Toxoplasmosis in South American coatis (<i>Nasua nasua</i>) in Tietê Ecological Park, São Paulo, SP, Brazil. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2016, 53, 1.	0.2	0
247	Drug-induced panniculitis due to deworming in a dog – case report. <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 2017, 54, 434-438.	0.2	0
248	Alguns aspectos nutricionais e metabólicos de bezerros mantidos com dois níveis diferentes de proteína na dieta e infectados experimentalmente com <i>Haemonchus placei</i> . <i>Brazilian Journal of Veterinary Research and Animal Science</i> , 1996, 33, 295.	0.2	0
249	Occurrence of <i>Toxoplasma gondii</i> antibodies in lowland tapirs maintained ex situ in Brazil and Paraguay. <i>Ciencia Rural</i> , 2017, 47, .	0.3	0
250	Serum anti- <i>Toxoplasma gondii</i> antibodies in <i>Passer domesticus</i> (Linnaeus, 1758) (Passeriformes:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 Animal Science, 2020, 57, e164867.	0.2	0
251	Occurrence of <i>Toxoplasma gondii</i> and <i>Neospora caninum</i> antibodies and risk factors in domiciliated dogs of Manaus, Amazonas, Brazil. <i>Brazilian Journal of Veterinary Parasitology</i> , 2022, 31, e020321.	0.2	0
252	Negative seroprevalence for <i>Toxoplasma gondii</i> in free-living primates from Central Amazonia. <i>Journal of Medical Primatology</i> , 2022, , .	0.3	0