
## Seng H Cheng

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11991345/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Defective intracellular transport and processing of CFTR is the molecular basis of most cystic fibrosis. Cell, 1990, 63, 827-834.	13.5	1,697
2	Phosphorylation of the R domain by cAMP-dependent protein kinase regulates the CFTR chloride channel. Cell, 1991, 66, 1027-1036.	13.5	651
3	Expression of cystic fibrosis transmembrane conductance regulator corrects defective chloride channel regulation in cystic fibrosis airway epithelial cells. Nature, 1990, 347, 358-363.	13.7	649
4	Sustained Therapeutic Reversal of Huntington's Disease by Transient Repression of Huntingtin Synthesis. Neuron, 2012, 74, 1031-1044.	3.8	635
5	Detailed Analysis of Structures and Formulations of Cationic Lipids for Efficient Gene Transfer to the Lung. Human Gene Therapy, 1996, 7, 1701-1717.	1.4	453
6	Antisense Oligonucleotides Delivered to the Mouse CNS Ameliorate Symptoms of Severe Spinal Muscular Atrophy. Science Translational Medicine, 2011, 3, 72ra18.	5.8	437
7	Targeting nuclear RNA for in vivo correction of myotonic dystrophy. Nature, 2012, 488, 111-115.	13.7	435
8	Repeated nebulisation of non-viral CFTR gene therapy in patients with cystic fibrosis: a randomised, double-blind, placebo-controlled, phase 2b trial. Lancet Respiratory Medicine,the, 2015, 3, 684-691.	5.2	344
9	Expression and characterization of the cystic fibrosis transmembrane conductance regulator. Nature, 1990, 347, 382-386.	13.7	337
10	Hypoxia-Inducible Factor-1 Mediates Activation of Cultured Vascular Endothelial Cells by Inducing Multiple Angiogenic Factors. Circulation Research, 2003, 93, 664-673.	2.0	292
11	CNS expression of glucocerebrosidase corrects α-synuclein pathology and memory in a mouse model of Gaucher-related synucleinopathy. Proceedings of the National Academy of Sciences of the United States of America, 2011, 108, 12101-12106.	3.3	282
12	Acid βâ€glucosidase mutants linked to gaucher disease, parkinson disease, and lewy body dementia alter αâ€synuclein processing. Annals of Neurology, 2011, 69, 940-953.	2.8	276
13	CpG-free plasmids confer reduced inflammation and sustained pulmonary gene expression. Nature Biotechnology, 2008, 26, 549-551.	9.4	269
14	CNS-targeted gene therapy improves survival and motor function in a mouse model of spinal muscular atrophy. Journal of Clinical Investigation, 2010, 120, 1253-1264.	3.9	252
15	Improved Cationic Lipid Formulations for In Vivo Gene Therapy. Annals of the New York Academy of Sciences, 1995, 772, 126-139.	1.8	249
16	Basis of Pulmonary Toxicity Associated with Cationic Lipid-Mediated Gene Transfer to the Mammalian Lung. Human Gene Therapy, 1997, 8, 689-707.	1.4	239
17	Cystic fibrosis transmembrane conductance regulator: A chloride channel with novel regulation. Neuron, 1992, 8, 821-829.	3.8	226
18	Comprehensive Analysis of the Acute Toxicities Induced by Systemic Administration of Cationic Lipid Plasmid DNA Complexes in Mice, Human Gene Therapy, 2000, 11, 2493-2513	1.4	218

#	Article	IF	CITATIONS
19	Generation and characterization of a ΔF508 cystic fibrosis mouse model. Nature Genetics, 1995, 10, 445-452.	9.4	215
20	Reduced Inflammatory Response to Plasmid DNA Vectors by Elimination and Inhibition of Immunostimulatory CpG Motifs. Molecular Therapy, 2000, 1, 255-262.	3.7	211
21	CpG-Depleted Plasmid DNA Vectors with Enhanced Safety and Long-Term Gene Expression in Vivo. Molecular Therapy, 2002, 5, 731-738.	3.7	203
22	Augmenting CNS glucocerebrosidase activity as a therapeutic strategy for parkinsonism and other Gaucher-related synucleinopathies. Proceedings of the National Academy of Sciences of the United States of America, 2013, 110, 3537-3542.	3.3	203
23	Contribution of Plasmid DNA to Inflammation in the Lung after Administration of Cationic Lipid:pDNA Complexes. Human Gene Therapy, 1999, 10, 223-234.	1.4	199
24	Dimorphic Effects of Transforming Growth Factor-Î <sup>2</sup> Signaling During Aortic Aneurysm Progression in Mice Suggest a Combinatorial Therapy for Marfan Syndrome. Arteriosclerosis, Thrombosis, and Vascular Biology, 2015, 35, 911-917.	1.1	189
25	Glucosylceramide synthase inhibition alleviates aberrations in synucleinopathy models. Proceedings of the United States of America, 2017, 114, 2699-2704.	3.3	165
26	Silencing Mutant Huntingtin by Adeno-Associated Virus-Mediated RNA Interference Ameliorates Disease Manifestations in the YAC128 Mouse Model of Huntington's Disease. Human Gene Therapy, 2014, 25, 461-474.	1.4	135
27	High and Sustained Transgene Expression in Vivo from Plasmid Vectors Containing a Hybrid Ubiquitin Promoter. Molecular Therapy, 2001, 4, 75-82.	3.7	129
28	Hypoxia Up-regulates Expression of Peroxisome Proliferator-activated Receptor Î <sup>3</sup> Angiopoietin-related Gene (PGAR) in Cardiomyocytes: Role of Hypoxia Inducible Factor 1α. Journal of Molecular and Cellular Cardiology, 2002, 34, 765-774.	0.9	128
29	Stabilization of Vascular Endothelial Growth Factor mRNA by Hypoxia-Inducible Factor 1. Biochemical and Biophysical Research Communications, 2002, 291, 908-914.	1.0	125
30	Fetal gene therapy for neurodegenerative disease of infants. Nature Medicine, 2018, 24, 1317-1323.	15.2	117
31	Optimization of Plasmid Vectors for High-Level Expression in Lung Epithelial Cells. Human Gene Therapy, 1997, 8, 575-584.	1.4	116
32	Intracellular protein trafficking defects in human disease. Trends in Cell Biology, 1992, 2, 145-149.	3.6	99
33	A Concentrated and Stable Aerosol Formulation of Cationic Lipid:DNA Complexes Giving High-Level Gene Expression in Mouse Lung. Human Gene Therapy, 1997, 8, 765-773.	1.4	93
34	Expression of constitutively stable hybrid hypoxia-inducible factor-1α protects cultured rat cardiomyocytes against simulated ischemia-reperfusion injury. American Journal of Physiology - Cell Physiology, 2005, 288, C314-C320.	2.1	89
35	Optimization of Formulations and Conditions for the Aerosol Delivery of Functional Cationic Lipid:DNA Complexes. Human Gene Therapy, 1997, 8, 313-322.	1.4	85
36	Partial restoration of cAMP-stimulated CFTR chloride channel activity in ΔF508 cells by deoxyspergualin. American Journal of Physiology - Cell Physiology, 1998, 275, C171-C178.	2.1	82

#	Article	IF	CITATIONS
37	Efficiency of Cationic Lipid-Mediated Transfection of Polarized and Differentiated Airway Epithelial Cells <i>In Vitro</i> and <i>In Vivo</i> . Human Gene Therapy, 1998, 9, 1531-1542.	1.4	81
38	Contribution of Toll-like Receptor 9 Signaling to the Acute Inflammatory Response to Nonviral Vectors. Molecular Therapy, 2004, 9, 241-248.	3.7	81
39	Translational Fidelity of Intrathecal Delivery of Self-Complementary AAV9–Survival Motor Neuron 1 for Spinal Muscular Atrophy. Human Gene Therapy, 2014, 25, 619-630.	1.4	79
40	Hypoxia-inducible factor 1 mediates hypoxia-induced cardiomyocyte lipid accumulation by reducing the DNA binding activity of peroxisome proliferator-activated receptor α/retinoid X receptor. Biochemical and Biophysical Research Communications, 2007, 364, 567-572.	1.0	67
41	A randomised, double-blind, placebo-controlled phase IIB clinical trial of repeated application of gene therapy in patients with cystic fibrosis: TableÂ1. Thorax, 2013, 68, 1075-1077.	2.7	66
42	Gaucher-related synucleinopathies: The examination of sporadic neurodegeneration from a rare (disease) angle. Progress in Neurobiology, 2015, 125, 47-62.	2.8	63
43	Inefficient cationic lipid-mediated siRNA and antisense oligonucleotide transfer to airway epithelial cells in vivo. Respiratory Research, 2006, 7, 26.	1.4	59
44	Antisense Oligonucleotide-Mediated Correction of Transcriptional Dysregulation is Correlated with Behavioral Benefits in the YAC128 Mouse Model of Huntington's Disease. Journal of Huntington's Disease, 2013, 2, 217-228.	0.9	58
45	Expression of angiopoietins in renal epithelial and clear cell carcinoma cells: regulation by hypoxia and participation in angiogenesis. American Journal of Physiology - Renal Physiology, 2004, 287, F649-F657.	1.3	55
46	Gene expression profiles in human cardiac cells subjected to hypoxia or expressing a hybrid form of HIF-11±. Physiological Genomics, 2002, 8, 23-32.	1.0	54
47	Systemic Delivery of a Peptide-Linked Morpholino Oligonucleotide Neutralizes Mutant RNA Toxicity in a Mouse Model of Myotonic Dystrophy. Nucleic Acid Therapeutics, 2013, 23, 109-117.	2.0	52
48	Glucocerebrosidase modulates cognitive and motor activities in murine models of Parkinson's disease. Human Molecular Genetics, 2016, 25, ddw124.	1.4	44
49	Correction of the cystic fibrosis defect by gene complementation in human intrahepatic biliary epithelial cell lines. Gastroenterology, 1995, 108, 584-592.	0.6	42
50	Electroporation enhances reporter gene expression following delivery of naked plasmid DNA to the lung. Journal of Gene Medicine, 2007, 9, 369-380.	1.4	42
51	IGF-1 delivery to CNS attenuates motor neuron cell death but does not improve motor function in type III SMA mice. Neurobiology of Disease, 2012, 45, 272-279.	2.1	41
52	Production of Cystic Fibrosis Transmembrane Conductance Regulator in the Milk of Transgenic Mice. Nature Biotechnology, 1992, 10, 74-77.	9.4	39
53	Adenovirus-Transduced Lung as a Portal for Delivering α-Galactosidase A into Systemic Circulation for Fabry Disease. Molecular Therapy, 2002, 5, 745-754.	3.7	37
54	DNA Sequences in Cationic Lipid:pDNA-Mediated Systemic Toxicities. Human Gene Therapy, 2003, 14, 203-214.	1.4	36

#	Article	IF	CITATIONS
55	Cationic Lipid Structure and Formulation Considerations for Optimal Gene Transfection of the Lung. Journal of Drug Targeting, 2000, 7, 453-469.	2.1	35
56	Inhibition of osteoclastogenesis by prolyl hydroxylase inhibitor dimethyloxallyl glycine. Journal of Bone and Mineral Metabolism, 2010, 28, 510-519.	1.3	33
57	Purification and Characterization of Recombinant Cystic Fibrosis Transmembrane Conductance Regulator from Chinese Hamster Ovary and Insect Cells. Journal of Biological Chemistry, 1995, 270, 17033-17043.	1.6	32
58	Aerosolization of Cationic Lipid:pDNA Complexes—In Vitro Optimization of Nebulizer Parameters for Human Clinical Studies. Human Gene Therapy, 1998, 9, 43-52.	1.4	31
59	Characterization of the oligosaccharide structures associated with the cystic fibrosis transmembrane conductance regulator. Glycobiology, 2000, 10, 1225-1233.	1.3	31
60	The safety profile of a cationic lipid-mediated cystic fibrosis gene transfer agent following repeated monthly aerosol administration toÂsheep. Biomaterials, 2013, 34, 10267-10277.	5.7	30
61	Prospects for the gene therapy of spinal muscular atrophy. Trends in Molecular Medicine, 2011, 17, 259-265.	3.5	29
62	Secreted Gaussia luciferase as a sensitive reporter gene for in vivo and ex vivo studies of airway gene transfer. Biomaterials, 2011, 32, 2614-2624.	5.7	29
63	The use of carboxymethylcellulose gel to increase non-viral gene transfer in mouse airways. Biomaterials, 2010, 31, 2665-2672.	5.7	27
64	A hypoxic inducible factorâ€1α hybrid enhances collateral development and reduces vascular leakage in diabetic rats. Journal of Gene Medicine, 2009, 11, 390-400.	1.4	26
65	Mutant <i>GBA1</i> Expression and Synucleinopathy Risk: First Insights from Cellular and Mouse Models. Neurodegenerative Diseases, 2012, 10, 195-202.	0.8	26
66	Partial correction of theα-galactosidase A deficiency and reduction of glycolipid storage in Fabry mice using synthetic vectors. Journal of Gene Medicine, 2004, 6, 85-92.	1.4	22
67	Gene Transfer to the CNS Is Efficacious in Immune-primed Mice Harboring Physiologically Relevant Titers of Anti-AAV Antibodies. Molecular Therapy, 2012, 20, 1713-1723.	3.7	22
68	A randomised, double-blind, placebo-controlled trial of repeated nebulisation of non-viral cystic fibrosis transmembrane conductance regulator (CFTR) gene therapy in patients with cystic fibrosis. Efficacy and Mechanism Evaluation, 2016, 3, 1-210.	0.9	22
69	Nitric oxide inhibits heterologous CFTR expression in polarized epithelial cells. American Journal of Physiology - Lung Cellular and Molecular Physiology, 1999, 277, L89-L96.	1.3	20
70	Increased Duration of Transgene Expression in the Lung with Plasmid DNA Vectors Harboring Adenovirus E4 Open Reading Frame 3. Human Gene Therapy, 1999, 10, 1833-1843.	1.4	20
71	Identification of transfected cell types following non-viral gene transfer to the murine lung. Journal of Gene Medicine, 2007, 9, 184-196.	1.4	18
72	Limitations of the Murine Nose in the Development of Nonviral Airway Gene Transfer. American Journal of Respiratory Cell and Molecular Biology, 2010, 43, 46-54.	1.4	18

#	Article	IF	CITATIONS
73	Cationic Lipid-Mediated Gene Delivery to the Airways. , 1999, , 39-68.		16
74	In vivo imaging of gene transfer to the respiratory tract. Biomaterials, 2008, 29, 1533-1540.	5.7	13
75	The role of doxorubicin in non-viral gene transfer in the lung. Biomaterials, 2009, 30, 1971-1977.	5.7	12
76	Low-frequency ultrasound increases non-viral gene transfer to the mouse lung. Acta Biochimica Et Biophysica Sinica, 2010, 42, 45-51.	0.9	11
77	Rapid identification of novel functional promoters for gene therapy. Journal of Molecular Medicine, 2012, 90, 1487-1496.	1.7	9
78	Restoration of Cyclic Adenosine Monophosphate–Stimulated Chloride Channel Activity in Human Cystic Fibrosis Tracheobronchial Submucosal Gland Cells by Adenovirus-Mediated and Cationic Lipid–Mediated Gene Transfer. American Journal of Respiratory Cell and Molecular Biology, 1999, 20, 1107-1115.	1.4	7
79	Assessment of the nuclear pore dilating agent transâ€cyclohexaneâ€1,2â€diol in differentiated airway epithelium. Journal of Gene Medicine, 2012, 14, 491-500.	1.4	6
80	Formulation of Synthetic Vectors for Cystic Fibrosis Gene Therapy. , 2002, 70, 585-598.		0
81	Formulation of Synthetic Gene Delivery Vectors for Transduction of the Airway Epithelium. , 2004, 245, 95-114.		0