## Eun Jung Choy

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/12042304/publications.pdf

Version: 2024-02-01

933447 996975 16 327 10 15 citations h-index g-index papers 16 16 16 479 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Food web structure of a restored macroalgal bed in the eastern Korean peninsula determined by C and N stable isotope analyses. Marine Biology, 2008, 153, 1181-1198.	1.5	69
2	Pathways of organic matter through food webs of diverse habitats in the regulated Nakdong River estuary (Korea). Estuarine, Coastal and Shelf Science, 2008, 78, 215-226.	2.1	58
3	Reproductive cycle and gross biochemical composition of the ark shell Scapharca subcrenata (Lischke, 1869) reared on subtidal mudflats in a temperate bay of Korea. Aquaculture, 2011, 322-323, 149-157.	3.5	37
4	Linking Intertidal and Subtidal Food Webs: Consumer-Mediated Transport of Intertidal Benthic Microalgal Carbon. PLoS ONE, 2015, 10, e0139802.	2.5	31
5	Quantifying the trophic base for benthic secondary production in the Nakdong River estuary of Korea using stable C and N isotopes. Journal of Experimental Marine Biology and Ecology, 2009, 382, 18-26.	1.5	24
6	The role of the adductor muscle as an energy storage organ in the pen shell <i>Atrina japonica</i> (Reeve, 1858). Journal of Molluscan Studies, 2015, 81, 502-511.	1.2	19
7	CONDITION, REPRODUCTIVE ACTIVITY, AND GROSS BIOCHEMICAL COMPOSITION OF THE MANILA CLAM, TAPES PHILIPPINARUM IN NATURAL AND NEWLY CREATED SANDY HABITATS OF THE SOUTHERN COAST OF KOREA. Journal of Shellfish Research, 2007, 26, 401-412.	0.9	15
8	Time-course uptake and elimination of benzo(a)pyrene and its damage to reproduction and ensuing reproductive outputs of Pacific oyster, Crassostrea gigas. Marine Biology, 2007, 151, 157-165.	1.5	15
9	Trophic Structures of Two Contrasting Estuarine Ecosystems With and without a Dike on the Temperate Coast of Korea as Determined by Stable Isotopes. Estuaries and Coasts, 2020, 43, 560-577.	2.2	12
10	Spatial distribution of small phytoplankton composition in the Chukchi Sea. Polar Biology, 2014, 37, 99-109.	1.2	11
11	Sterol dynamics of heterotrophic Tetraselmis suecica and its nutritional implication in the bivalve aquaculture. Aquaculture Research, 2004, 35, 371-377.	1.8	10
12	Spatial and Temporal Variations of Microphytobenthos on the Common Reed Phragmites australis bed in a Marine Protected Area of Yeoja Bay, Korea. Wetlands, 2013, 33, 737-745.	1.5	9
13	Stable isotope analysis of a newly established macrofaunal food web 1.5 years after the Hebei Spirit oil spill. Marine Pollution Bulletin, 2015, 90, 167-180.	5.0	9
14	Incubation capacity limits clutch size in south polar skuas. Antarctic Science, 2010, 22, 19.	0.9	6
15	Effect of the Hebei Spirit Oil Spill on the Condition, Reproduction, and Energy Storage Cycle of the Manila Clam Ruditapes philippinarum on the West Coast of Korea. Estuaries and Coasts, 2020, 43, 602-614.	2.2	2
16	The Sterolic Properties of Heterotrophic Tetraselmis suecica. Journal of Fisheries Science and Technology, 2004, 7, 34-38.	0.2	0