
## Swa Himaya

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/12064359/publications.pdf Version: 2024-02-01



ςνα Ηιμανά

7

#	Article	IF	CITATIONS
1	An active peptide purified from gastrointestinal enzyme hydrolysate of Pacific cod skin gelatin attenuates angiotensin-1 converting enzyme (ACE) activity and cellular oxidative stress. Food Chemistry, 2012, 132, 1872-1882.	8.2	165
2	Free radical scavenging and angiotensin-I converting enzyme inhibitory peptides from Pacific cod (Gadus macrocephalus) skin gelatin. International Journal of Biological Macromolecules, 2011, 49, 1110-1116.	7.5	102
3	Fumigaclavine C from a Marine-Derived Fungus Aspergillus Fumigatus Induces Apoptosis in MCF-7 Breast Cancer Cells. Marine Drugs, 2013, 11, 5063-5086.	4.6	61
4	Tyrosol exerts a protective effect against dopaminergic neuronal cell death in in vitro model of Parkinson's disease. Food Chemistry, 2013, 141, 1147-1157.	8.2	48
5	Neoechinulin A suppresses amyloid-β oligomer-induced microglia activation and thereby protects PC-12 cells from inflammation-mediated toxicity. NeuroToxicology, 2013, 35, 30-40.	3.0	44
6	Sea cucumber, Stichopus japonicus ethyl acetate fraction modulates the lipopolysaccharide induced iNOS and COX-2 via MAPK signaling pathway in murine macrophages. Environmental Toxicology and Pharmacology, 2010, 30, 68-75.	4.0	40
7	1-(5-bromo-2-hydroxy-4-methoxyphenyl)ethanone [SE1] suppresses pro-inflammatory responses by blocking NF-κB and MAPK signaling pathways in activated microglia. European Journal of Pharmacology, 2011, 670, 608-616.	3.5	29
8	Anti-proliferative effects of isosclerone isolated from marine fungus Aspergillus fumigatus in MCF-7 human breast cancer cells. Process Biochemistry, 2014, 49, 2292-2298.	3.7	20
9	Isolation and characterization of marine-derived Mucor sp. for the fermentative production of tyrosol. Process Biochemistry, 2014, 49, 1402-1408.	3.7	8

10 Marine Nutraceuticals. , 2015, , 995-1014.