

# Swa Himaya

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/12064359/publications.pdf>

Version: 2024-02-01

10  
papers

524  
citations

1163117

8  
h-index

1474206

9  
g-index

10  
all docs

10  
docs citations

10  
times ranked

864  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Marine Nutraceuticals. , 2015, , 995-1014.  |     | 7         |
| 2  | Anti-proliferative effects of isosclerone isolated from marine fungus <i>Aspergillus fumigatus</i> in MCF-7 human breast cancer cells. <i>Process Biochemistry</i> , 2014, 49, 2292-2298.   | 3.7 | 20        |
| 3  | Isolation and characterization of marine-derived <i>Mucor</i> sp. for the fermentative production of tyrosol. <i>Process Biochemistry</i> , 2014, 49, 1402-1408.  | 3.7 | 8         |
| 4  | Neoechinulin A suppresses amyloid- $\beta^2$ oligomer-induced microglia activation and thereby protects PC-12 cells from inflammation-mediated toxicity. <i>NeuroToxicology</i> , 2013, 35, 30-40.  | 3.0 | 44        |
| 5  | Tyrosol exerts a protective effect against dopaminergic neuronal cell death in in vitro model of Parkinson's disease. <i>Food Chemistry</i> , 2013, 141, 1147-1157.   | 8.2 | 48        |
| 6  | Fumigaclavine C from a Marine-Derived Fungus <i>Aspergillus Fumigatus</i> Induces Apoptosis in MCF-7 Breast Cancer Cells. <i>Marine Drugs</i> , 2013, 11, 5063-5086.  | 4.6 | 61        |
| 7  | An active peptide purified from gastrointestinal enzyme hydrolysate of Pacific cod skin gelatin attenuates angiotensin-1 converting enzyme (ACE) activity and cellular oxidative stress. <i>Food Chemistry</i> , 2012, 132, 1872-1882.        | 8.2 | 165       |
| 8  | Free radical scavenging and angiotensin-I converting enzyme inhibitory peptides from Pacific cod ( <i>Gadus macrocephalus</i> ) skin gelatin. <i>International Journal of Biological Macromolecules</i> , 2011, 49, 1110-1116.                | 7.5 | 102       |
| 9  | 1-(5-bromo-2-hydroxy-4-methoxyphenyl)ethanone [SE1] suppresses pro-inflammatory responses by blocking NF- $\kappa$ B and MAPK signaling pathways in activated microglia. <i>European Journal of Pharmacology</i> , 2011, 670, 608-616.        | 3.5 | 29        |
| 10 | Sea cucumber, <i>Stichopus japonicus</i> ethyl acetate fraction modulates the lipopolysaccharide induced iNOS and COX-2 via MAPK signaling pathway in murine macrophages. <i>Environmental Toxicology and Pharmacology</i> , 2010, 30, 68-75. | 4.0 | 40        |