

Svetlana Skurtveit

List of Publications by Citations

Source: <https://exaly.com/author-pdf/1209355/svetlana-skurtveit-publications-by-citations.pdf>
Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

222 papers	4,653 citations	36 h-index	55 g-index
239 ext. papers	5,387 ext. citations	3.7 avg, IF	5.63 L-index

#	Paper	IF	Citations
222	Environmental assessment of Norwegian priority pharmaceuticals based on the EMEA guideline. <i>Ecotoxicology and Environmental Safety</i> , 2008 , 71, 328-40	7	155
221	Coffee intake and mortality from liver cirrhosis. <i>Annals of Epidemiology</i> , 2003 , 13, 419-23	6.4	145
220	Prescription drug use among fathers and mothers before and during pregnancy. A population-based cohort study of 106,000 pregnancies in Norway 2004-2006. <i>British Journal of Clinical Pharmacology</i> , 2008 , 65, 653-60	3.8	138
219	Risk of road traffic accidents associated with the prescription of drugs: a registry-based cohort study. <i>Annals of Epidemiology</i> , 2007 , 17, 597-602	6.4	135
218	International trends in clozapine use: a study in 17 countries. <i>Acta Psychiatrica Scandinavica</i> , 2017 , 136, 37-51	6.5	120
217	Road traffic accident risk related to prescriptions of the hypnotics zopiclone, zolpidem, flunitrazepam and nitrazepam. <i>Sleep Medicine</i> , 2008 , 9, 818-22	4.6	116
216	International trends in antipsychotic use: A study in 16 countries, 2005-2014. <i>European Neuropsychopharmacology</i> , 2017 , 27, 1064-1076	1.2	112
215	Clinical impairment of benzodiazepines--relation between benzodiazepine concentrations and impairment in apprehended drivers. <i>Drug and Alcohol Dependence</i> , 2002 , 68, 131-41	4.9	84
214	Engagement with opioid maintenance treatment and reductions in crime: a longitudinal national cohort study. <i>Addiction</i> , 2012 , 107, 393-9	4.6	78
213	Neonatal outcomes following in utero exposure to methadone or buprenorphine: a National Cohort Study of opioid-agonist treatment of Pregnant Women in Norway from 1996 to 2009. <i>Drug and Alcohol Dependence</i> , 2013 , 127, 200-6	4.9	78
212	Increasing use of opioids from 2004 to 2007 - pharmacoepidemiological data from a complete national prescription database in Norway. <i>European Journal of Pain</i> , 2010 , 14, 289-94	3.7	72
211	Switching statins in Norway after new reimbursement policy: a nationwide prescription study. <i>British Journal of Clinical Pharmacology</i> , 2007 , 64, 476-81	3.8	63
210	High risk of overdose death following release from prison: variations in mortality during a 15-year observation period. <i>Addiction</i> , 2017 , 112, 1432-1439	4.6	61
209	Pharmacokinetic differences of morphine and morphine-glucuronides are reflected in locomotor activity. <i>Pharmacology Biochemistry and Behavior</i> , 2002 , 73, 883-92	3.9	58
208	Drinking pattern among adolescents with immigrant and Norwegian backgrounds: a two-way influence?. <i>Addiction</i> , 2005 , 100, 1453-63	4.6	57
207	Undertreatment and overtreatment with statins: the Oslo Health Study 2000-2001. <i>Journal of Internal Medicine</i> , 2004 , 255, 494-502	10.8	55
206	Risk of diabetes after gestational diabetes and preeclampsia. A registry-based study of 230,000 women in Norway. <i>European Journal of Epidemiology</i> , 2011 , 26, 157-63	12.1	52

205	Use of anti-asthmatic medications as a proxy for prevalence of asthma in children and adolescents in Norway: a nationwide prescription database analysis. <i>European Journal of Clinical Pharmacology</i> , 2007 , 63, 693-8	2.8	52
204	Rearrest rates among Norwegian drugged drivers compared with drunken drivers. <i>Drug and Alcohol Dependence</i> , 2002 , 66, 85-92	4.9	52
203	Chronic pain and use of opioids: a population-based pharmacoepidemiological study from the Norwegian prescription database and the Nord-Trøndelag health study. <i>Pain</i> , 2014 , 155, 1213-1221	8	51
202	Breastfeeding reduces the need for withdrawal treatment in opioid-exposed infants. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2013 , 102, 1060-6	3.1	47
201	Prenatal exposure to antidepressants and language competence at age three: results from a large population-based pregnancy cohort in Norway. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2014 , 121, 1621-31	3.7	46
200	Testing for benzodiazepine inebriation--relationship between benzodiazepine concentration and simple clinical tests for impairment in a sample of drugged drivers. <i>European Journal of Clinical Pharmacology</i> , 2003 , 59, 593-601	2.8	46
199	Minor increase in risk of road traffic accidents after prescriptions of antidepressants: a study of population registry data in Norway. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, 1099-103	4.6	45
198	Prevalence of use of non-prescription analgesics in the Norwegian HUNT3 population: Impact of gender, age, exercise and prescription of opioids. <i>BMC Public Health</i> , 2015 , 15, 461	4.1	44
197	Starting insomnia treatment: the use of benzodiazepines versus z-hypnotics. A prescription database study of predictors. <i>European Journal of Clinical Pharmacology</i> , 2009 , 65, 295-301	2.8	44
196	Carisoprodol use and abuse in Norway: a pharmacoepidemiological study. <i>British Journal of Clinical Pharmacology</i> , 2007 , 64, 210-8	3.8	44
195	What is the point of guidelines? Benzodiazepine and z-hypnotic use by an elderly population. <i>Sleep Medicine</i> , 2012 , 13, 893-7	4.6	43
194	Use of anxiolytic or hypnotic drugs and total mortality in a general middle-aged population. <i>Pharmacoepidemiology and Drug Safety</i> , 2007 , 16, 913-8	2.6	42
193	Many correlates of poor quality of life among substance users entering treatment are not addiction-specific. <i>Health and Quality of Life Outcomes</i> , 2016 , 14, 39	3	41
192	Criminal convictions among dependent heroin users during a 3-year period prior to opioid maintenance treatment: a longitudinal national cohort study. <i>Journal of Substance Abuse Treatment</i> , 2011 , 41, 407-14	4.2	41
191	Benzodiazepines predict use of opioids--a follow-up study of 17,074 men and women. <i>Pain Medicine</i> , 2010 , 11, 805-14	2.8	41
190	High validity of mother-reported use of antiasthmatics among children: a comparison with a population-based prescription database. <i>Journal of Clinical Epidemiology</i> , 2011 , 64, 878-84	5.7	39
189	The risk of motor vehicle accidents involving drivers with prescriptions for codeine or tramadol. <i>Clinical Pharmacology and Therapeutics</i> , 2009 , 85, 596-9	6.1	39
188	Impairment due to intake of carisoprodol. <i>Drug and Alcohol Dependence</i> , 2004 , 74, 311-8	4.9	38

187	Nicotine dependence predicts repeated use of prescribed opioids. Prospective population-based cohort study. <i>Annals of Epidemiology</i> , 2010 , 20, 890-7	6.4	36
186	Increased mortality among previously apprehended drunken and drugged drivers. <i>Drug and Alcohol Dependence</i> , 2002 , 68, 143-50	4.9	36
185	Hypnotic drug use among 0-17 year olds during 2004-2011: a nationwide prescription database study. <i>Scandinavian Journal of Public Health</i> , 2012 , 40, 704-11	3	34
184	Effect of Time-Dependent Selective Serotonin Reuptake Inhibitor Antidepressants During Pregnancy on Behavioral, Emotional, and Social Development in Preschool-Aged Children. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2018 , 57, 200-208	7.2	33
183	Persistent opioid use and socio-economic factors: a population-based study in Norway. <i>Acta Anaesthesiologica Scandinavica</i> , 2014 , 58, 437-45	1.9	33
182	Prescribed opioid analgesic use developments in three Nordic countries, 2006-2017. <i>Scandinavian Journal of Pain</i> , 2019 , 19, 345-353	1.9	32
181	Differential patterns of opioid use: defining persistent opioid use in a prescription database. <i>European Journal of Pain</i> , 2012 , 16, 359-69	3.7	32
180	Adherence, persistence and switch rates for anticholinergic drugs used for overactive bladder in women: data from the Norwegian Prescription Database. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2013 , 92, 1208-15	3.8	31
179	Pregnancy-Related Risk Factors Are Associated With a Significant Burden of Treated Hypertension Within 10 Years of Delivery: Findings From a Population-Based Norwegian Cohort. <i>Journal of the American Heart Association</i> , 2018 , 7,	6	31
178	To what extent does a cohort of new users of weak opioids develop persistent or probable problematic opioid use?. <i>Pain</i> , 2011 , 152, 1555-1561	8	30
177	Drug exposure: inclusion of dispensed drugs before pregnancy may lead to underestimation of risk associations. <i>Journal of Clinical Epidemiology</i> , 2013 , 66, 964-72	5.7	29
176	Concomitant medication among persistent opioid users with chronic non-malignant pain. <i>Acta Anaesthesiologica Scandinavica</i> , 2012 , 56, 1267-76	1.9	29
175	Use of prescribed opioid analgesics and co-medication with benzodiazepines in women before, during, and after pregnancy: a population-based cohort study. <i>European Journal of Clinical Pharmacology</i> , 2011 , 67, 953-60	2.8	29
174	Prescription pattern of codeine for non-malignant pain: a pharmacoepidemiological study from the Norwegian Prescription Database. <i>Acta Anaesthesiologica Scandinavica</i> , 2009 , 53, 627-33	1.9	29
173	Road traffic accident risk in patients with diabetes mellitus receiving blood glucose-lowering drugs. Prospective follow-up study. <i>Diabetic Medicine</i> , 2009 , 26, 404-8	3.5	29
172	A pharmacoepidemiological cohort study of subjects starting strong opioids for nonmalignant pain: a study from the Norwegian Prescription Database. <i>Pain</i> , 2013 , 154, 2487-2493	8	28
171	Effects of preconceptional paternal drug exposure on birth outcomes: cohort study of 340 000 pregnancies using Norwegian population-based databases. <i>British Journal of Clinical Pharmacology</i> , 2013 , 75, 1134-41	3.8	28
170	The validity of self-reported prescription medication use among adolescents varied by therapeutic class. <i>Journal of Clinical Epidemiology</i> , 2008 , 61, 714-7	5.7	28

169	The risk of traffic accidents after prescriptions of carisoprodol. <i>Accident Analysis and Prevention</i> , 2007 , 39, 1050-5	6.1	28
168	Drugged drivers in Norway with benzodiazepine detections. <i>Forensic Science International</i> , 2002 , 125, 75-82	2.6	28
167	Codeine and clinical impairment in samples in which morphine is not detected. <i>European Journal of Clinical Pharmacology</i> , 2003 , 58, 785-9	2.8	28
166	Female drivers suspected for drunken or drugged driving. <i>Forensic Science International</i> , 1995 , 75, 139-48	4.6	28
165	Psychotropic drug use among 0-17 year olds during 2004-2014: a nationwide prescription database study. <i>BMC Psychiatry</i> , 2016 , 16, 12	4.2	28
164	Hepatitis C Treatment Uptake among Patients Who Have Received Opioid Substitution Treatment: A Population-Based Study. <i>PLoS ONE</i> , 2016 , 11, e0166451	3.7	27
163	Association of Maternal Use of Benzodiazepines and Z-Hypnotics During Pregnancy With Motor and Communication Skills and Attention-Deficit/Hyperactivity Disorder Symptoms in Preschoolers. <i>JAMA Network Open</i> , 2019 , 2, e191435	10.4	26
162	Flunitrazepam: psychomotor impairment, agitation and paradoxical reactions. <i>Forensic Science International</i> , 2006 , 159, 83-91	2.6	26
161	Prenatal exposure to anxiolytics and hypnotics and language competence at 3 years of age. <i>European Journal of Clinical Pharmacology</i> , 2015 , 71, 283-91	2.8	25
160	Trends in prescription drug use during pregnancy and postpartum in Norway, 2005 to 2015. <i>Pharmacoepidemiology and Drug Safety</i> , 2018 , 27, 995-1004	2.6	25
159	Influence of socioeconomic factors on the adherence of alendronate treatment in incident users in Norway. <i>Pharmacoepidemiology and Drug Safety</i> , 2012 , 21, 297-304	2.6	24
158	Neonatal outcomes after fetal exposure to methadone and buprenorphine: national registry studies from the Czech Republic and Norway. <i>Addiction</i> , 2018 , 113, 1286-1294	4.6	23
157	Methadone-Related Overdose Deaths in a Liberal Opioid Maintenance Treatment Programme. <i>European Addiction Research</i> , 2016 , 22, 249-58	4.6	23
156	. <i>Pharmacogenetics and Genomics</i> , 2003 , 13, 383-388		23
155	Fast quantification of ethanol in whole blood specimens by the enzymatic alcohol dehydrogenase method. Optimization by experimental design. <i>Journal of Analytical Toxicology</i> , 2005 , 29, 66-70	2.9	23
154	Heroin-using drivers: importance of morphine and morphine-6-glucuronide on late clinical impairment. <i>European Journal of Clinical Pharmacology</i> , 2006 , 62, 905-12	2.8	22
153	Association of prenatal exposure to benzodiazepines and child internalizing problems: A sibling-controlled cohort study. <i>PLoS ONE</i> , 2017 , 12, e0181042	3.7	22
152	Factors associated with dropout among patients in opioid maintenance treatment (OMT) and predictors of re-entry. A national registry-based study. <i>Addictive Behaviors</i> , 2014 , 39, 1504-9	4.2	21

151	Does Pregabalin (Lyrica®) help patients reduce their use of benzodiazepines? A comparison with gabapentin using the Norwegian Prescription Database. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2010 , 107, 883-6	3.1	21
150	Statin treatment in a cohort of 20 212 men and women in Norway according to cardiovascular risk factors and level of education. <i>British Journal of Clinical Pharmacology</i> , 2009 , 67, 355-62	3.8	21
149	Prevalence, incidence and persistence of anti-asthma medication use in 2- to 29-year-olds: a nationwide prescription study. <i>European Journal of Clinical Pharmacology</i> , 2010 , 66, 399-406	2.8	21
148	Prescription drug use among pregnant women in opioid Maintenance Treatment. <i>Addiction</i> , 2013 , 108, 367-76	4.6	20
147	Validation of FRAX and the impact of self-reported falls among elderly in a general population: the HUNT study, Norway. <i>Osteoporosis International</i> , 2017 , 28, 2935-2944	5.3	20
146	Increase in psychotropic drug use between 2006 and 2010 among adolescents in Norway: a nationwide prescription database study. <i>Clinical Epidemiology</i> , 2012 , 4, 225-31	5.9	20
145	Substance use pattern, self-control and social network are associated with crime in a substance-using population. <i>Drug and Alcohol Review</i> , 2017 , 36, 245-252	3.2	19
144	Comorbidities in an asthma population 8-29 years old: a study from the Norwegian Prescription Database. <i>Pharmacoepidemiology and Drug Safety</i> , 2012 , 21, 1045-52	2.6	19
143	An increased risk of motor vehicle accidents after prescription of methadone. <i>Addiction</i> , 2012 , 107, 967-72	4.6	19
142	Long-term Use of Z-Hypnotics and Co-medication with Benzodiazepines and Opioids. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2017 , 120, 292-298	3.1	18
141	Use of prescribed opioids by children and adolescents: Differences between Denmark, Norway and Sweden. <i>European Journal of Pain</i> , 2015 , 19, 1095-100	3.7	18
140	Morphine-6-glucuronide-induced locomotor stimulation in mice: role of opioid receptors. <i>Basic and Clinical Pharmacology and Toxicology</i> , 1998 , 82, 3-10		18
139	Aspects of statin prescribing in Norwegian counties with high, average and low statin consumption - an individual-level prescription database study. <i>BMC Clinical Pharmacology</i> , 2007 , 7, 14		18
138	Tapering from Methadone or Buprenorphine during Pregnancy: Maternal and Neonatal Outcomes in Norway 1996-2009. <i>European Addiction Research</i> , 2015 , 21, 253-61	4.6	17
137	Dispensing of benzodiazepines and benzodiazepine-related drugs to pregnant women: a population-based cohort study. <i>European Journal of Clinical Pharmacology</i> , 2014 , 70, 1367-74	2.8	17
136	The association between smoking and subsequent repeated use of prescribed opioids among adolescents and young adults--a population-based cohort study. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 90-8	2.6	17
135	Study of rearrests for drunken driving in Norway. <i>Forensic Science International</i> , 1998 , 92, 21-8	2.6	17
134	Mortality among subjects previously apprehended for driving under the influence of traffic-hazardous medicinal drugs. <i>Drug and Alcohol Dependence</i> , 2005 , 79, 423-9	4.9	17

133	Paediatric Off-Label Use of Melatonin--A Register Linkage Study between the Norwegian Prescription Database and Patient Register. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2015 , 117, 267-73	3.1	16
132	Motor development in children prenatally exposed to selective serotonin reuptake inhibitors: a large population-based pregnancy cohort study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2016 , 123, 1908-1917	3.7	16
131	Laxative prescriptions to cancer outpatients receiving opioids: a study from the Norwegian prescription database. <i>Supportive Care in Cancer</i> , 2013 , 21, 67-73	3.9	16
130	Co-medication of statins and CYP3A4 inhibitors before and after introduction of new reimbursement policy. <i>British Journal of Clinical Pharmacology</i> , 2009 , 67, 234-41	3.8	16
129	Integrated treatment of hepatitis C virus infection among people who inject drugs: study protocol for a randomised controlled trial (INTRO-HCV). <i>BMC Infectious Diseases</i> , 2019 , 19, 943	4	15
128	Psychotropic drug use among persons with mental distress symptoms: a population-based study in Norway. <i>Scandinavian Journal of Public Health</i> , 2007 , 35, 356-64	3	15
127	Use of psychotropic drugs in an urban adolescent population: the impact of health-related variables, lifestyle and sociodemographic factors--The Oslo Health Study 2000-2001. <i>Pharmacoepidemiology and Drug Safety</i> , 2005 , 14, 277-83	2.6	15
126	Building abstinent networks is an important resource in improving quality of life. <i>Drug and Alcohol Dependence</i> , 2017 , 180, 431-438	4.9	14
125	A complete national cohort study of prescriptions of analgesics and benzodiazepines to cancer survivors in Norway 10 years after diagnosis. <i>Pain</i> , 2019 , 160, 852-859	8	14
124	Preeclampsia in pregnancy and later use of antihypertensive drugs. <i>European Journal of Epidemiology</i> , 2015 , 30, 501-8	12.1	14
123	Antidepressants to cancer patients during the last year of life--a population-based study. <i>Psycho-Oncology</i> , 2013 , 22, 506-14	3.9	14
122	A day-by-day investigation of changes in criminal convictions before and after entering and leaving opioid maintenance treatment: a national cohort study. <i>BMC Psychiatry</i> , 2013 , 13, 262	4.2	14
121	Introduction of low dose transdermal buprenorphine -- did it influence use of potentially addictive drugs in chronic non-malignant pain patients?. <i>European Journal of Pain</i> , 2009 , 13, 949-53	3.7	14
120	Prescriptions of opioids to children and adolescents; a study from a national prescription database in Norway. <i>Paediatric Anaesthesia</i> , 2010 , 20, 537-44	1.8	14
119	Self-reported data on medicine use in the Norwegian Mother and Child cohort study compared to data from the Norwegian Prescription Database. <i>Norsk Epidemiologi</i> , 2014 , 24,	2.8	14
118	COVID-19 survey among people who use drugs in three cities in Norway. <i>Drug and Alcohol Dependence</i> , 2020 , 217, 108302	4.9	14
117	The duration and course of opioid therapy in patients with chronic non-malignant pain. <i>Acta Anaesthesiologica Scandinavica</i> , 2016 , 60, 128-37	1.9	14
116	Language competence and communication skills in 3-year-old children after prenatal exposure to analgesic opioids. <i>Pharmacoepidemiology and Drug Safety</i> , 2017 , 26, 625-634	2.6	13

115	Antidepressant drug use among adolescents during 2004-2013: a population-based register linkage study. <i>Acta Psychiatrica Scandinavica</i> , 2016 , 134, 420-429	6.5	13
114	Co-morbidity in persistent opioid users with chronic non-malignant pain in Norway. <i>European Journal of Pain</i> , 2014 , 18, 1083-93	3.7	13
113	Benzodiazepine use in all alcohol consumers predicts use of opioids in patients 20 years later--a follow-up study of 13,390 men and women aged 40-42 years. <i>Pharmacoepidemiology and Drug Safety</i> , 2008 , 17, 926-933	2.6	13
112	The CYP2C19 genotype and the use of oral contraceptives influence the pharmacokinetics of carisoprodol in healthy human subjects. <i>European Journal of Clinical Pharmacology</i> , 2005 , 61, 499-506	2.8	13
111	A Unique Opportunity to Study Short and Long Term Consequences in Children Prenatally Exposed to Illicit Drugs and Opioid Maintenance Treatment Using Czech and Scandinavian Registers. <i>Central European Journal of Public Health</i> , 2016 , 24, 248-251	1.2	13
110	Mental disorder diagnoses among children and adolescents who use antipsychotic drugs. <i>European Neuropsychopharmacology</i> , 2016 , 26, 1412-1418	1.2	13
109	An increased risk of road traffic accidents after prescriptions of lithium or valproate?. <i>Pharmacoepidemiology and Drug Safety</i> , 2009 , 18, 492-6	2.6	12
108	Effect of the market withdrawal of carisoprodol on use of other prescribed drugs with abuse potential. <i>Clinical Pharmacology and Therapeutics</i> , 2012 , 91, 438-41	6.1	12
107	Substance use during and after pregnancy among a national cohort of pregnant women in opioid maintenance treatment and their partners. <i>Journal of Substance Use</i> , 2012 , 17, 277-286	0.8	12
106	Determinants of control of high blood pressure. The Oslo Health Study 2000-2001. <i>Blood Pressure</i> , 2004 , 13, 343-9	1.7	12
105	Lack of crosstolerance between morphine and morphine-6-glucuronide as revealed by locomotor activity. <i>Pharmacology Biochemistry and Behavior</i> , 2000 , 66, 205-10	3.9	12
104	Proton pump inhibitors and fracture risk. The HUNT study, Norway. <i>Osteoporosis International</i> , 2020 , 31, 109-118	5.3	12
103	Validating the generic quality of life tool "QOL10" in a substance use disorder treatment cohort exposes a unique social construct. <i>BMC Medical Research Methodology</i> , 2016 , 16, 60	4.7	12
102	Substance use and sociodemographic background as risk factors for lifetime psychotic experiences in a non-clinical sample. <i>Journal of Substance Abuse Treatment</i> , 2017 , 74, 42-47	4.2	11
101	Trends in prescription of strong opioids for 41-80 year old Norwegians, 2005-2010. <i>European Journal of Pain</i> , 2014 , 18, 438-46	3.7	11
100	Prescribing of drugs for attention-deficit hyperactivity disorder in opioid maintenance treatment patients in Norway. <i>European Addiction Research</i> , 2014 , 20, 59-65	4.6	11
99	Prescriptions of Antidepressants to Survivors of Cancer in Childhood, Adolescence, and Young Adulthood: A Population-Based Study. <i>Journal of Adolescent and Young Adult Oncology</i> , 2017 , 6, 120-126	2.2	10
98	Use of anti-osteoporotic drugs in central Norway after a forearm fracture. <i>Archives of Osteoporosis</i> , 2015 , 10, 235	2.9	10

97	Prescription of analgesics to patients in opioid maintenance therapy: a pharmacoepidemiological study. <i>Drug and Alcohol Dependence</i> , 2011 , 116, 158-62	4.9	10
96	Characteristics of drivers testing positive for heroin or ecstasy in Norway. <i>Traffic Injury Prevention</i> , 2004 , 5, 107-11	1.8	10
95	Recidivism among drunken and drugged drivers in Norway. <i>Alcohol and Alcoholism</i> , 1996 , 31, 609-11	3.5	10
94	Increased prescription rates of anxiolytics and hypnotics to survivors of cancer in childhood, adolescence, and young adulthood-A population-based study. <i>Pediatric Blood and Cancer</i> , 2018 , 65, e26830	3.0	10
93	Increase in diagnosis of depressive disorders contributes to the increase in antidepressant use in adolescents. <i>Acta Psychiatrica Scandinavica</i> , 2018 , 137, 413-421	6.5	9
92	The association between prescribed opioid use for mothers and children: a record-linkage study. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 111-8	2.8	9
91	Low Calcium Intake in Midpregnancy Is Associated with Hypertension Development within 10 Years after Pregnancy: The Norwegian Mother and Child Cohort Study. <i>Journal of Nutrition</i> , 2017 , 147, 1757-1763	4.1	9
90	A Cohort Study on Long-Term Adverse Effects of Parental Drinking: Background and Study Design. <i>Substance Abuse: Research and Treatment</i> , 2015 , 9, 77-83	1.6	9
89	Polypharmacy of potentially addictive medication in the older persons--quantifying usage. <i>Pharmacoepidemiology and Drug Safety</i> , 2012 , 21, 199-206	2.6	9
88	Association between prescribing hypnotics for parents and children in Norway. <i>Archives of Disease in Childhood</i> , 2013 , 98, 732-6	2.2	9
87	Sex differences in gamma-glutamyltransferase in people aged 40-42 years in two Norwegian counties. <i>Drug and Alcohol Dependence</i> , 2002 , 67, 95-8	4.9	9
86	Benzodiazepine prescription for patients in treatment for drug use disorders: a nationwide cohort study in Denmark, 2000-2010. <i>BMC Psychiatry</i> , 2016 , 16, 168	4.2	9
85	Association of Constellations of Parental Risk With Children's Subsequent Anxiety and Depression: Findings From a HUNT Survey and Health Registry Study. <i>JAMA Pediatrics</i> , 2019 , 173, 251-259	8.3	9
84	Anti-osteoporosis drug use: too little, too much, or just right? The HUNT study, Norway. <i>Osteoporosis International</i> , 2018 , 29, 1875-1885	5.3	9
83	Road traffic noise and registry based use of sleep medication. <i>Environmental Health</i> , 2017 , 16, 110	6	8
82	Mental distress and subsequent use of psychotropic drugs among adolescents-a prospective register linkage study. <i>Journal of Adolescent Health</i> , 2012 , 50, 578-87	5.8	8
81	Performance of the WHOQOL-BREF among Norwegian substance use disorder patients. <i>BMC Medical Research Methodology</i> , 2019 , 19, 44	4.7	7
80	Hospitalization of children after prenatal exposure to opioid maintenance therapy during pregnancy: a national registry study from the Czech Republic. <i>Addiction</i> , 2019 , 114, 1225-1235	4.6	7

79	Associations of Maternal Use of Benzodiazepines or Benzodiazepine-like Hypnotics During Pregnancy With Immediate Pregnancy Outcomes in Norway. <i>JAMA Network Open</i> , 2020 , 3, e205860	10.4	7
78	Dispensation of attention deficit hyperactivity disorder (ADHD) medications in patients receiving opioid agonist therapy; a national prospective cohort study in Norway from 2015 to 2017. <i>BMC Psychiatry</i> , 2020 , 20, 119	4.2	7
77	Prescriptions of analgesics during complete disease trajectories in patients who are diagnosed with and die from cancer within the five-year period 2005-2009. <i>European Journal of Pain</i> , 2017 , 21, 530-540	3.7	7
76	The Norwegian Offender Mental Health and Addiction Study - Design and Implementation of a National Survey and Prospective Cohort Study. <i>Substance Abuse: Research and Treatment</i> , 2015 , 9, 59-66	1.6	7
75	Opioid switching to methadone: a pharmacoepidemiological study from a national prescription database. <i>Palliative Medicine</i> , 2012 , 26, 804-12	5.5	7
74	Low-dose transdermal buprenorphine - long-term use and co-medication with other potentially addictive drugs. <i>Acta Anaesthesiologica Scandinavica</i> , 2012 , 56, 88-94	1.9	7
73	A 2-year prospective study of psychological distress among a national cohort of pregnant women in opioid maintenance treatment and their partners. <i>Journal of Substance Use</i> , 2013 , 18, 148-160	0.8	7
72	Behavioural sensitization in mice induced by morphine-glucuronide metabolites. <i>Pharmacology Biochemistry and Behavior</i> , 2008 , 90, 578-85	3.9	7
71	Morphine-3-glucuronide inhibits morphine induced, but enhances morphine-6-glucuronide induced locomotor activity in mice. <i>Pharmacology Biochemistry and Behavior</i> , 2007 , 86, 576-86	3.9	7
70	Cannabis Use during Pregnancy and Risk of Adverse Birth Outcomes: A Longitudinal Cohort Study. <i>European Addiction Research</i> , 2021 , 27, 131-141	4.6	7
69	Multiple prescribers in older frequent opioid users--does it mean abuse?. <i>Journal of Population Therapeutics and Clinical Pharmacology</i> , 2013 , 20, e397-405	3.3	7
68	Preterm births and use of medication in early adulthood: a population-based registry study. <i>Pharmacoepidemiology and Drug Safety</i> , 2017 , 26, 742-751	2.6	6
67	Dispensations of benzodiazepines, z-hypnotics, and gabapentinoids to patients receiving opioid agonist therapy; a prospective cohort study in Norway from 2013 to 2017. <i>BMC Health Services Research</i> , 2020 , 20, 352	2.9	6
66	Prenatal Exposure to Folic Acid and Antidepressants and Language Development: A Population-Based Cohort Study. <i>Journal of Clinical Psychopharmacology</i> , 2016 , 36, 333-9	1.7	6
65	Tramadol use in Norway: A register-based population study. <i>Pharmacoepidemiology and Drug Safety</i> , 2019 , 28, 54-61	2.6	6
64	Disability pension is associated with the use of benzodiazepines 20 years later: a prospective study. <i>Scandinavian Journal of Public Health</i> , 2009 , 37, 320-6	3	6
63	Occupational physical activity, overweight, and mortality: a follow-up study of 47,405 Norwegian women and men. <i>Research Quarterly for Exercise and Sport</i> , 2007 , 78, 151-61	1.9	6
62	Coffee and wine consumption is associated with reduced mortality from alcoholic liver disease: follow-up of 219,279 Norwegian men and women aged 30-67 years. <i>Annals of Epidemiology</i> , 2018 , 28, 753-758	6.4	6

61	Prescription of analgesics to long-term survivors of cancer in early adulthood, adolescence, and childhood in Norway: a national cohort study. <i>Pain</i> , 2020 , 161, 1083-1091	8	5
60	The incidence, psychiatric co-morbidity and pharmacological treatment of severe mental disorders in children and adolescents. <i>European Psychiatry</i> , 2018 , 49, 16-22	6	5
59	Consumption of and satisfaction with health care among opioid users with chronic non-malignant pain. <i>Acta Anaesthesiologica Scandinavica</i> , 2015 , 59, 1355-66	1.9	5
58	Prenatal Methamphetamine Exposure and Adverse Neonatal Outcomes: A Nationwide Cohort Study. <i>European Addiction Research</i> , 2021 , 27, 97-106	4.6	5
57	Integrated treatment of hepatitis C virus infection among people who inject drugs: A multicenter randomized controlled trial (INTRO-HCV). <i>PLoS Medicine</i> , 2021 , 18, e1003653	11.6	5
56	Association between blood carisoprodol:meprobamate concentration ratios and CYP2C19 genotype in carisoprodol-drugged drivers: decreased metabolic capacity in heterozygous CYP2C19*1/CYP2C19*2 subjects?. <i>Pharmacogenetics and Genomics</i> , 2003 , 13, 383-8		5
55	Uptake and predictors of direct-acting antiviral treatment for hepatitis C among people receiving opioid agonist therapy in Sweden and Norway: a drug utilization study from 2014 to 2017. <i>Substance Abuse Treatment, Prevention, and Policy</i> , 2020 , 15, 44	3.4	4
54	Similarities and Differences in Victimization Risk Factors for Nonoffending and Offending Substance Users. <i>Victims and Offenders</i> , 2018 , 13, 526-541	1.5	4
53	Prenatal exposure to opioid maintenance treatment and neonatal outcomes: Nationwide registry studies from the Czech Republic and Norway. <i>Pharmacology Research and Perspectives</i> , 2019 , 7, e00501	3.1	4
52	Prevalence, incidence and persistence of etanercept and adalimumab in Norway 2005-2009. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 457-63	2.6	4
51	Mental distress and subsequent use of anxiolytic drugs--a prospective population-based cohort study of 16,000 individuals. <i>Scandinavian Journal of Public Health</i> , 2010 , 38, 465-73	3	4
50	Opioids for outpatients with cancer in their last year of life: A nationwide pharmacoepidemiological study. <i>Journal of Opioid Management</i> , 2016 , 12, 25-36	0.8	4
49	Living a normal life? Follow-up study of women who had been in opioid maintenance treatment during pregnancy. <i>Journal of Substance Abuse Treatment</i> , 2020 , 113, 108004	4.2	3
48	In utero exposure to analgesic opioids and language development in 5-year old children. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 736-744	2.6	3
47	Socioeconomic characteristics of women with substance use disorder during pregnancy and neonatal outcomes in their newborns: A national registry study from the Czech Republic. <i>Drug and Alcohol Dependence</i> , 2020 , 209, 107933	4.9	3
46	Medication use and association with urinary incontinence in women: Data from the Norwegian Prescription Database and the HUNT study. <i>Neurourology and Urodynamics</i> , 2018 , 37, 1448-1457	2.3	3
45	Misuse of prescription drugs and overdose deaths. <i>Journal of Substance Use</i> , 2016 , 21, 515-520	0.8	3
44	Agreement between self-reported and registry-based use of sleep medications and tranquilizers. <i>Pharmacoepidemiology and Drug Safety</i> , 2019 , 28, 1336-1343	2.6	3

43	Emergency service use is common in the year before death among drug users who die from an overdose. <i>Journal of Substance Use</i> , 2017 , 22, 331-336	0.8	3
42	Driving under the influence (DUI) among patients in opioid maintenance treatment (OMT): a registry-based national cohort study. <i>Addiction</i> , 2013 , 108, 1954-61	4.6	3
41	Disability pension as predictor of later use of benzodiazepines among benzodiazepine users. <i>Social Science and Medicine</i> , 2010 , 70, 921-5	5.1	3
40	Erratum to "Sex differences in gamma-glutamyltransferase in people aged 40-42 years in two Norwegian counties". <i>Drug and Alcohol Dependence</i> , 2003 , 69, 103-6	4.9	3
39	Reimbursement of analgesics for chronic pain. <i>Tidsskrift for Den Norske Lægeforening</i> , 2012 , 132, 2489-93	3.5	3
38	Prescription of antidepressants to patients on opioid maintenance therapy: a pharmacoepidemiological study. <i>Norsk Epidemiologi</i> , 2011 , 21,	2.8	3
37	Incidence of diagnosed pediatric anxiety disorders and use of prescription drugs: a nation-wide registry study. <i>European Child and Adolescent Psychiatry</i> , 2020 , 29, 1063-1073	5.5	3
36	Confirming the factor structure of a generic quality of life instrument among pre-treatment substance use disorder patients. <i>Health and Quality of Life Outcomes</i> , 2019 , 17, 84	3	2
35	Dispensing of prescribed analgesics in Norway among young people with foreign-or Norwegian-born parents. <i>Scandinavian Journal of Pain</i> , 2011 , 2, 36-44	1.9	2
34	Why do sales of lipid-lowering drugs vary between counties in Norway? Evidence from the OPPHED Health Study 2000-2001. <i>Scandinavian Journal of Primary Health Care</i> , 2006 , 24, 115-21	2.7	2
33	Carisoprodol should be taken off the market. <i>Southern Medical Journal</i> , 2008 , 101, 1074-5	0.6	2
32	Potentially addictive drugs dispensing to patients receiving opioid agonist therapy: a register-based prospective cohort study in Norway and Sweden from 2015 to 2017. <i>BMJ Open</i> , 2020 , 10, e036860	3	2
31	Too much or too little opioids to patients receiving opioid agonist therapy in Norway (2013-2017): a prospective cohort study. <i>BMC Health Services Research</i> , 2020 , 20, 668	2.9	2
30	On the path towards universal coverage of hepatitis C treatment among people receiving opioid agonist therapy (OAT) in Norway: a prospective cohort study from 2013 to 2017. <i>BMJ Open</i> , 2020 , 10, e036355	3	2
29	Parental risk constellations and future alcohol use disorder (AUD) in offspring: A combined HUNT survey and health registries study. <i>Psychology of Addictive Behaviors</i> , 2021 ,	3.4	2
28	Use of hypnotic drugs among children, adolescents, and young adults in Scandinavia. <i>Acta Psychiatrica Scandinavica</i> , 2021 , 144, 100-112	6.5	2
27	Effect of the market withdrawal of dextropropoxyphene on use of other prescribed analgesics. <i>Scandinavian Journal of Pain</i> , 2018 , 18, 667-674	1.9	2
26	Hypnotics use in children 0-18 months: moderate agreement between mother-reported survey data and prescription registry data. <i>Journal of Pharmaceutical Policy and Practice</i> , 2017 , 10, 28	3.2	1

25	Use of nasal fentanyl for cancer pain: A pharmacoepidemiological study. <i>Palliative Medicine</i> , 2015 , 29, 661-6	5.5	1
24	A prospective study of asthma and subsequent use of hypnotics in young adults. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 370-7	2.6	1
23	The absence of evidence is not the same as evidence for absence!. <i>Addiction</i> , 2008 , 103, 513-4	4.6	1
22	Pre conception use of cannabis and cocaine among men with pregnant partners. <i>NAD Nordic Studies on Alcohol and Drugs</i> , 2020 , 37, 43-53	1.1	1
21	In utero opioid exposure and risk of infections in childhood: A multinational Nordic cohort study. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 1596-1604	2.6	1
20	Nationwide Study of Neuropsychiatric Comorbidity and Medicines Use in Children With Autism Spectrum Disorder in Norway. <i>Frontiers in Psychiatry</i> , 2020 , 11, 596032	5	1
19	Impact of clinical and sociodemographic factors on fatigue among patients with substance use disorder: a cohort study from Norway for the period 2016-2020. <i>Substance Abuse Treatment, Prevention, and Policy</i> , 2020 , 15, 93	3.4	1
18	Opioid maintenance treatment in the Czech Republic, Norway and Denmark: a study protocol of a comparative registry linkage study. <i>BMJ Open</i> , 2021 , 11, e047028	3	1
17	Association of Timing and Duration of Prenatal Analgesic Opioid Exposure With Attention-Deficit/Hyperactivity Disorder in Children. <i>JAMA Network Open</i> , 2021 , 4, e2124324	10.4	1
16	Changes in the use of antiseizure medications in children and adolescents in Norway, 2009-2018.. <i>Epilepsy Research</i> , 2022 , 181, 106872	3	0
15	Desistance from crime following substance use treatment: the role of treatment retention, social network and self-control. <i>BMC Psychiatry</i> , 2021 , 21, 563	4.2	0
14	Opioid maintenance treatment of pregnant women in the Scandinavian countries.. <i>NAD Nordic Studies on Alcohol and Drugs</i> , 2020 , 37, 298-312	1.1	0
13	Commentary on Raitasalo et al. (2015): the great potential in nation-wide registers to study prescription drug use and abuse. <i>Addiction</i> , 2015 , 110, 644-5	4.6	
12	Reduced prescribing of benzodiazepines in Denmark and Norway. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2015 , 116, 457-8	3.1	
11	Mother-reported and prescription registry data on use of hypnotics for children 0-18 months as a risk factor later development of ADHD. <i>European Psychiatry</i> , 2017 , 41, S440-S441	6	
10	Incidence of and factors associated with anticholinergic drug use among Norwegian women with urinary incontinence. <i>International Urogynecology Journal</i> , 2018 , 29, 489-495	2	
9	How are opioids used in Norway? Persistent use, utilization of depot formulation and age profile in non-palliation patients. <i>Scandinavian Journal of Pain</i> , 2010 , 1, 171-171	1.9	
8	Women's experiences of giving birth in Northwest Russia in 2000 and 2002 and in Northern Norway 2000. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2007 , 86, 373-5	3.8	

7	665 ALMOST ALL CHRONIC NON-MALIGNANT PAIN PATIENTS ON LONG-TERM OPIOIDS ARE USING SHORT-ACTING MEDICATION. <i>European Journal of Pain</i> , 2006 , 10, S174-S174	3.7
6	Association Between Prescribed Hypnotics in Infants and Toddlers and Later ADHD: A Large Cohort Study from Norway. <i>Child Psychiatry and Human Development</i> , 2021 , 52, 533-543	3.3
5	Prescriptions of analgesics during chronic cancer disease trajectories: A complete national cohort study. <i>Pharmacoepidemiology and Drug Safety</i> , 2021 , 30, 1504-1513	2.6
4	Pre-injury dispensing of psychoactive prescription drugs in a ten years trauma population: a retrospective registry analysis. <i>Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine</i> , 2021 , 29, 125	3.6
3	Filled prescriptions for opioids among children and adolescents - A nationwide study from 2010 to 2018. <i>Acta Anaesthesiologica Scandinavica</i> , 2021 , 65, 1475-1483	1.9
2	PriSUD-Nordic-Diagnosing and Treating Substance Use Disorders in the Prison Population: Protocol for a Mixed Methods Study.. <i>JMIR Research Protocols</i> , 2022 , 11, e35182	2
1	Prescription opioids among older adults: ten years of data across five countries.. <i>BMC Geriatrics</i> , 2022 , 22, 429	4.1