Svetlana Skurtveit

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/1209355/svetlana-skurtveit-publications-by-year.pdf

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

4,653 36 222 55 h-index g-index citations papers 5,387 5.63 3.7 239 L-index avg, IF ext. citations ext. papers

#	Paper	IF	Citations
222	Changes in the use of antiseizure medications in children and adolescents in Norway, 2009-2018 <i>Epilepsy Research</i> , 2022 , 181, 106872	3	O
221	PriSUD-Nordic-Diagnosing and Treating Substance Use Disorders in the Prison Population: Protocol for a Mixed Methods Study <i>JMIR Research Protocols</i> , 2022 , 11, e35182	2	
220	Prescription opioids among older adults: ten years of data across five countries <i>BMC Geriatrics</i> , 2022 , 22, 429	4.1	
219	Desistance from crime following substance use treatment: the role of treatment retention, social network and self-control. <i>BMC Psychiatry</i> , 2021 , 21, 563	4.2	0
218	Cannabis Use during Pregnancy and Risk of Adverse Birth Outcomes: A Longitudinal Cohort Study. <i>European Addiction Research</i> , 2021 , 27, 131-141	4.6	7
217	Prenatal Methamphetamine Exposure and Adverse Neonatal Outcomes: A Nationwide Cohort Study. <i>European Addiction Research</i> , 2021 , 27, 97-106	4.6	5
216	Association Between Prescribed Hypnotics in Infants and Toddlers and Later ADHD: A Large Cohort Study from Norway. <i>Child Psychiatry and Human Development</i> , 2021 , 52, 533-543	3.3	
215	Parental risk constellations and future alcohol use disorder (AUD) in offspring: A combined HUNT survey and health registries study. <i>Psychology of Addictive Behaviors</i> , 2021 ,	3.4	2
214	Opioid maintenance treatment in the Czech Republic, Norway and Denmark: a study protocol of a comparative registry linkage study. <i>BMJ Open</i> , 2021 , 11, e047028	3	1
213	Use of hypnotic drugs among children, adolescents, and young adults in Scandinavia. <i>Acta Psychiatrica Scandinavica</i> , 2021 , 144, 100-112	6.5	2
212	Integrated treatment of hepatitis C virus infection among people who inject drugs: A multicenter randomized controlled trial (INTRO-HCV). <i>PLoS Medicine</i> , 2021 , 18, e1003653	11.6	5
211	Prescriptions of analgesics during chronic cancer disease trajectories: A complete national cohort study. <i>Pharmacoepidemiology and Drug Safety</i> , 2021 , 30, 1504-1513	2.6	
210	Pre-injury dispensing of psychoactive prescription drugs in a ten years trauma population: a retrospective registry analysis. <i>Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine</i> , 2021 , 29, 125	3.6	
209	Filled prescriptions for opioids among children and adolescents - A nationwide study from 2010 to 2018. <i>Acta Anaesthesiologica Scandinavica</i> , 2021 , 65, 1475-1483	1.9	
208	Association of Timing and Duration of Prenatal Analgesic Opioid Exposure With Attention-Deficit/Hyperactivity Disorder in Children. <i>JAMA Network Open</i> , 2021 , 4, e2124324	10.4	1
207	Associations of Maternal Use of Benzodiazepines or Benzodiazepine-like Hypnotics During Pregnancy With Immediate Pregnancy Outcomes in Norway. <i>JAMA Network Open</i> , 2020 , 3, e205860	10.4	7
206	Living a normal life? Follow-up study of women who had been in opioid maintenance treatment during pregnancy. <i>Journal of Substance Abuse Treatment</i> , 2020 , 113, 108004	4.2	3

205	Prescription of analgesics to long-term survivors of cancer in early adulthood, adolescence, and childhood in Norway: a national cohort study. <i>Pain</i> , 2020 , 161, 1083-1091	8	5
204	In utero exposure to analgesic opioids and language development in 5-year old children. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 736-744	2.6	3
203	Dispensation of attention deficit hyperactivity disorder (ADHD) medications in patients receiving opioid agonist therapy; a national prospective cohort study in Norway from 2015 to 2017. <i>BMC Psychiatry</i> , 2020 , 20, 119	4.2	7
202	Uptake and predictors of direct-acting antiviral treatment for hepatitis C among people receiving opioid agonist therapy in Sweden and Norway: a drug utilization study from 2014 to 2017. <i>Substance Abuse Treatment, Prevention, and Policy,</i> 2020 , 15, 44	3.4	4
201	Socioeconomic characteristics of women with substance use disorder during pregnancy and neonatal outcomes in their newborns: A national registry study from the Czech Republic. <i>Drug and Alcohol Dependence</i> , 2020 , 209, 107933	4.9	3
2 00	Dispensations of benzodiazepines, z-hypnotics, and gabapentinoids to patients receiving opioid agonist therapy; a prospective cohort study in Norway from 2013 to 2017. <i>BMC Health Services Research</i> , 2020 , 20, 352	2.9	6
199	Incidence of diagnosed pediatric anxiety disorders and use of prescription drugs: a nation-wide registry study. <i>European Child and Adolescent Psychiatry</i> , 2020 , 29, 1063-1073	5.5	3
198	Proton pump inhibitors and fracture risk. The HUNT study, Norway. <i>Osteoporosis International</i> , 2020 , 31, 109-118	5.3	12
197	Pre conception use of cannabis and cocaine among men with pregnant partners. <i>NAD Nordic Studies on Alcohol and Drugs</i> , 2020 , 37, 43-53	1.1	1
196	COVID-19 survey among people who use drugs in three cities in Norway. <i>Drug and Alcohol Dependence</i> , 2020 , 217, 108302	4.9	14
195	Potentially addictive drugs dispensing to patients receiving opioid agonist therapy: a register-based prospective cohort study in Norway and Sweden from 2015 to 2017. <i>BMJ Open</i> , 2020 , 10, e036860	3	2
194	Opioid maintenance treatment of pregnant women in the Scandinavian countries <i>NAD Nordic Studies on Alcohol and Drugs</i> , 2020 , 37, 298-312	1.1	O
193	Too much or too little opioids to patients receiving opioid agonist therapy in Norway (2013-2017): a prospective cohort study. <i>BMC Health Services Research</i> , 2020 , 20, 668	2.9	2
192	On the path towards universal coverage of hepatitis C treatment among people receiving opioid agonist therapy (OAT) in Norway: a prospective cohort study from 2013 to 2017. <i>BMJ Open</i> , 2020 , 10, e036355	3	2
191	In utero opioid exposure and risk of infections in childhood: A multinational Nordic cohort study. <i>Pharmacoepidemiology and Drug Safety</i> , 2020 , 29, 1596-1604	2.6	1
190	Nationwide Study of Neuropsychiatric Comorbidity and Medicines Use in Children With Autism Spectrum Disorder in Norway. <i>Frontiers in Psychiatry</i> , 2020 , 11, 596032	5	1
189	Impact of clinical and sociodemographic factors on fatigue among patients with substance use disorder: a cohort study from Norway for the period 2016-2020. <i>Substance Abuse Treatment, Prevention, and Policy,</i> 2020 , 15, 93	3.4	1
188	Prescribed opioid analgesic use developments in three Nordic countries, 2006-2017. <i>Scandinavian Journal of Pain</i> , 2019 , 19, 345-353	1.9	32

187	Confirming the factor structure of a generic quality of life instrument among pre-treatment substance use disorder patients. <i>Health and Quality of Life Outcomes</i> , 2019 , 17, 84	3	2
186	A complete national cohort study of prescriptions of analgesics and benzodiazepines to cancer survivors in Norway 10 years after diagnosis. <i>Pain</i> , 2019 , 160, 852-859	8	14
185	Performance of the WHOQOL-BREF among Norwegian substance use disorder patients. <i>BMC Medical Research Methodology</i> , 2019 , 19, 44	4.7	7
184	Association of Maternal Use of Benzodiazepines and Z-Hypnotics During Pregnancy With Motor and Communication Skills and Attention-Deficit/Hyperactivity Disorder Symptoms in Preschoolers. JAMA Network Open, 2019 , 2, e191435	10.4	26
183	Hospitalization of children after prenatal exposure to opioid maintenance therapy during pregnancy: a national registry study from the Czech Republic. <i>Addiction</i> , 2019 , 114, 1225-1235	4.6	7
182	Tramadol use in Norway: A register-based population study. <i>Pharmacoepidemiology and Drug Safety</i> , 2019 , 28, 54-61	2.6	6
181	Prenatal exposure to opioid maintenance treatment and neonatal outcomes: Nationwide registry studies from the Czech Republic and Norway. <i>Pharmacology Research and Perspectives</i> , 2019 , 7, e00501	3.1	4
180	Integrated treatment of hepatitis C virus infection among people who inject drugs: study protocol for a randomised controlled trial (INTRO-HCV). <i>BMC Infectious Diseases</i> , 2019 , 19, 943	4	15
179	Agreement between self-reported and registry-based use of sleep medications and tranquilizers. <i>Pharmacoepidemiology and Drug Safety</i> , 2019 , 28, 1336-1343	2.6	3
178	Association of Constellations of Parental Risk With Children@Subsequent Anxiety and Depression: Findings From a HUNT Survey and Health Registry Study. <i>JAMA Pediatrics</i> , 2019 , 173, 251-259	8.3	9
177	Neonatal outcomes after fetal exposure to methadone and buprenorphine: national registry studies from the Czech Republic and Norway. <i>Addiction</i> , 2018 , 113, 1286-1294	4.6	23
176	Increase in diagnosis of depressive disorders contributes to the increase in antidepressant use in adolescents. <i>Acta Psychiatrica Scandinavica</i> , 2018 , 137, 413-421	6.5	9
175	The incidence, psychiatric co-morbidity and pharmacological treatment of severe mental disorders in children and adolescents. <i>European Psychiatry</i> , 2018 , 49, 16-22	6	5
174	Medication use and association with urinary incontinence in women: Data from the Norwegian Prescription Database and the HUNT study. <i>Neurourology and Urodynamics</i> , 2018 , 37, 1448-1457	2.3	3
173	Effect of Time-Dependent Selective Serotonin Reuptake Inhibitor Antidepressants During Pregnancy on Behavioral, Emotional, and Social Development in Preschool-Aged Children. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2018 , 57, 200-208	7.2	33
172	Similarities and Differences in Victimization Risk Factors for Nonoffending and Offending Substance Users. <i>Victims and Offenders</i> , 2018 , 13, 526-541	1.5	4
171	Trends in prescription drug use during pregnancy and postpartum in Norway, 2005 to 2015. <i>Pharmacoepidemiology and Drug Safety</i> , 2018 , 27, 995-1004	2.6	25
170	Incidence of and factors associated with anticholinergic drug use among Norwegian women with urinary incontinence. <i>International Urogynecology Journal</i> , 2018 , 29, 489-495	2	

169	Increased prescription rates of anxiolytics and hypnotics to survivors of cancer in childhood, adolescence, and young adulthood-A population-based study. <i>Pediatric Blood and Cancer</i> , 2018 , 65, e26	0588	10
168	Effect of the market withdrawal of dextropropoxyphene on use of other prescribed analgesics. <i>Scandinavian Journal of Pain</i> , 2018 , 18, 667-674	1.9	2
167	Coffee and wine consumption is associated with reduced mortality from alcoholic liver disease: follow-up of 219,279 Norwegian men and women aged 30-67 years. <i>Annals of Epidemiology</i> , 2018 , 28, 753-758	6.4	6
166	Pregnancy-Related Risk Factors Are Associated With a Significant Burden of Treated Hypertension Within 10 Years of Delivery: Findings From a Population-Based Norwegian Cohort. <i>Journal of the American Heart Association</i> , 2018 , 7,	6	31
165	Anti-osteoporosis drug use: too little, too much, or just right? The HUNT study, Norway. <i>Osteoporosis International</i> , 2018 , 29, 1875-1885	5.3	9
164	Substance use pattern, self-control and social network are associated with crime in a substance-using population. <i>Drug and Alcohol Review</i> , 2017 , 36, 245-252	3.2	19
163	Substance use and sociodemographic background as risk factors for lifetime psychotic experiences in a non-clinical sample. <i>Journal of Substance Abuse Treatment</i> , 2017 , 74, 42-47	4.2	11
162	Preterm births and use of medication in early adulthood: a population-based registry study. <i>Pharmacoepidemiology and Drug Safety</i> , 2017 , 26, 742-751	2.6	6
161	International trends in clozapine use: a study in 17 countries. <i>Acta Psychiatrica Scandinavica</i> , 2017 , 136, 37-51	6.5	120
160	High risk of overdose death following release from prison: variations in mortality during a 15-year observation period. <i>Addiction</i> , 2017 , 112, 1432-1439	4.6	61
159	Language competence and communication skills in 3-year-old children after prenatal exposure to analgesic opioids. <i>Pharmacoepidemiology and Drug Safety</i> , 2017 , 26, 625-634	2.6	13
158	Building abstinent networks is an important resource in improving quality of life. <i>Drug and Alcohol Dependence</i> , 2017 , 180, 431-438	4.9	14
157	Hypnotics use in children 0-18[months: moderate agreement between mother-reported survey data and prescription registry data. <i>Journal of Pharmaceutical Policy and Practice</i> , 2017 , 10, 28	3.2	1
156	Road traffic noise and registry based use of sleep medication. <i>Environmental Health</i> , 2017 , 16, 110	6	8
155	Low Calcium Intake in Midpregnancy Is Associated with Hypertension Development within 10 Years after Pregnancy: The Norwegian Mother and Child Cohort Study. <i>Journal of Nutrition</i> , 2017 , 147, 1757-	1763	9
154	Mother-reported and prescription registry data on use of hypnotics for children 018 months as a risk factor later development of ADHD. <i>European Psychiatry</i> , 2017 , 41, S440-S441	6	
153	International trends in antipsychotic use: A study in 16 countries, 2005-2014. <i>European Neuropsychopharmacology</i> , 2017 , 27, 1064-1076	1.2	112
152	Validation of FRAX and the impact of self-reported falls among elderly in a general population: the HUNT study, Norway. <i>Osteoporosis International</i> , 2017 , 28, 2935-2944	5.3	20

151	Long-term Use of Z-Hypnotics and Co-medication with Benzodiazepines and Opioids. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2017 , 120, 292-298	3.1	18
150	Prescriptions of Antidepressants to Survivors of Cancer in Childhood, Adolescence, and Young Adulthood: A Population-Based Study. <i>Journal of Adolescent and Young Adult Oncology</i> , 2017 , 6, 120-12-	6 ^{2.2}	10
149	Emergency service use is common in the year before death among drug users who die from an overdose. <i>Journal of Substance Use</i> , 2017 , 22, 331-336	0.8	3
148	Prescriptions of analgesics during complete disease trajectories in patients who are diagnosed with and die from cancer within the five-year period 2005-2009. <i>European Journal of Pain</i> , 2017 , 21, 530-540	3.7	7
147	Association of prenatal exposure to benzodiazepines and child internalizing problems: A sibling-controlled cohort study. <i>PLoS ONE</i> , 2017 , 12, e0181042	3.7	22
146	Antidepressant drug use among adolescents during 2004-2013: a population-based register linkage study. <i>Acta Psychiatrica Scandinavica</i> , 2016 , 134, 420-429	6.5	13
145	Misuse of prescription drugs and overdose deaths. <i>Journal of Substance Use</i> , 2016 , 21, 515-520	0.8	3
144	Methadone-Related Overdose Deaths in a Liberal Opioid Maintenance Treatment Programme. <i>European Addiction Research</i> , 2016 , 22, 249-58	4.6	23
143	Many correlates of poor quality of life among substance users entering treatment are not addiction-specific. <i>Health and Quality of Life Outcomes</i> , 2016 , 14, 39	3	41
142	Prenatal Exposure to Folic Acid and Antidepressants and Language Development: A Population-Based Cohort Study. <i>Journal of Clinical Psychopharmacology</i> , 2016 , 36, 333-9	1.7	6
141	Motor development in children prenatally exposed to selective serotonin reuptake inhibitors: a large population-based pregnancy cohort study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2016 , 123, 1908-1917	3.7	16
140	A Unique Opportunity to Study Short and Long Term Consequences in Children Prenatally Exposed to Illicit Drugs and Opioid Maintenance Treatment Using Czech and Scandinavian Registers. <i>Central European Journal of Public Health</i> , 2016 , 24, 248-251	1.2	13
139	Opioids for outpatients with cancer in their last year of life: A nationwide pharmacoepidemiological study. <i>Journal of Opioid Management</i> , 2016 , 12, 25-36	0.8	4
138	Hepatitis C Treatment Uptake among Patients Who Have Received Opioid Substitution Treatment: A Population-Based Study. <i>PLoS ONE</i> , 2016 , 11, e0166451	3.7	27
137	Validating the generic quality of life tool "QOL10" in a substance use disorder treatment cohort exposes a unique social construct. <i>BMC Medical Research Methodology</i> , 2016 , 16, 60	4.7	12
136	Benzodiazepine prescription for patients in treatment for drug use disorders: a nationwide cohort study in Denmark, 2000-2010. <i>BMC Psychiatry</i> , 2016 , 16, 168	4.2	9
135	Psychotropic drug use among 0-17 year olds during 2004-2014: a nationwide prescription database study. <i>BMC Psychiatry</i> , 2016 , 16, 12	4.2	28
134	The duration and course of opioid therapy in patients with chronic non-malignant pain. <i>Acta Anaesthesiologica Scandinavica</i> , 2016 , 60, 128-37	1.9	14

133	Mental disorder diagnoses among children and adolescents who use antipsychotic drugs. <i>European Neuropsychopharmacology</i> , 2016 , 26, 1412-1418	1.2	13
132	Paediatric Off-Label Use of MelatoninA Register Linkage Study between the Norwegian Prescription Database and Patient Register. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2015 , 117, 267-73	3.1	16
131	Prenatal exposure to anxiolytics and hypnotics and language competence at 3 years of age. <i>European Journal of Clinical Pharmacology</i> , 2015 , 71, 283-91	2.8	25
130	Preeclampsia in pregnancy and later use of antihypertensive drugs. <i>European Journal of Epidemiology</i> , 2015 , 30, 501-8	12.1	14
129	Tapering from Methadone or Buprenorphine during Pregnancy: Maternal and Neonatal Outcomes in Norway 1996-2009. <i>European Addiction Research</i> , 2015 , 21, 253-61	4.6	17
128	Commentary on Raitasalo et al. (2015): the great potential in nation-wide registers to study prescription drug use and abuse. <i>Addiction</i> , 2015 , 110, 644-5	4.6	
127	Reduced prescribing of benzodiazepines in Denmark and Norway. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2015 , 116, 457-8	3.1	
126	Use of nasal fentanyl for cancer pain: A pharmacoepidemiological study. <i>Palliative Medicine</i> , 2015 , 29, 661-6	5.5	1
125	A Cohort Study on Long-Term Adverse Effects of Parental Drinking: Background and Study Design. <i>Substance Abuse: Research and Treatment</i> , 2015 , 9, 77-83	1.6	9
124	Use of anti-osteoporotic drugs in central Norway after a forearm fracture. <i>Archives of Osteoporosis</i> , 2015 , 10, 235	2.9	10
123	Consumption of and satisfaction with health care among opioid users with chronic non-malignant pain. <i>Acta Anaesthesiologica Scandinavica</i> , 2015 , 59, 1355-66	1.9	5
122	Use of prescribed opioids by children and adolescents: Differences between Denmark, Norway and Sweden. <i>European Journal of Pain</i> , 2015 , 19, 1095-100	3.7	18
121	The Norwegian Offender Mental Health and Addiction Study - Design and Implementation of a National Survey and Prospective Cohort Study. <i>Substance Abuse: Research and Treatment</i> , 2015 , 9, 59-66	1.6	7
120	Prevalence of use of non-prescription analgesics in the Norwegian HUNT3 population: Impact of gender, age, exercise and prescription of opioids. <i>BMC Public Health</i> , 2015 , 15, 461	4.1	44
119	Dispensing of benzodiazepines and benzodiazepine-related drugs to pregnant women: a population-based cohort study. <i>European Journal of Clinical Pharmacology</i> , 2014 , 70, 1367-74	2.8	17
118	Chronic pain and use of opioids: a population-based pharmacoepidemiological study from the Norwegian prescription database and the Nord-Tr\(\text{B}\) delag health study. <i>Pain</i> , 2014 , 155, 1213-1221	8	51
117	Factors associated with dropout among patients in opioid maintenance treatment (OMT) and predictors of re-entry. A national registry-based study. <i>Addictive Behaviors</i> , 2014 , 39, 1504-9	4.2	21
116	Persistent opioid use and socio-economic factors: a population-based study in Norway. <i>Acta Anaesthesiologica Scandinavica</i> , 2014 , 58, 437-45	1.9	33

115	Trends in prescription of strong opioids for 41-80 year old Norwegians, 2005-2010. <i>European Journal of Pain</i> , 2014 , 18, 438-46	3.7	11
114	Co-morbidity in persistent opioid users with chronic non-malignant pain in Norway. <i>European Journal of Pain</i> , 2014 , 18, 1083-93	3.7	13
113	Prenatal exposure to antidepressants and language competence at age three: results from a large population-based pregnancy cohort in Norway. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2014 , 121, 1621-31	3.7	46
112	Prescribing of drugs for attention-deficit hyperactivity disorder in opioid maintenance treatment patients in Norway. <i>European Addiction Research</i> , 2014 , 20, 59-65	4.6	11
111	Self-reported data on medicine use in the Norwegian Mother and Child cohort study compared to data from the Norwegian Prescription Database. <i>Norsk Epidemiologi</i> , 2014 , 24,	2.8	14
110	Antidepressants to cancer patients during the last year of lifea population-based study. <i>Psycho-Oncology</i> , 2013 , 22, 506-14	3.9	14
109	The association between prescribed opioid use for mothers and children: a record-linkage study. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 111-8	2.8	9
108	A day-by-day investigation of changes in criminal convictions before and after entering and leaving opioid maintenance treatment: a national cohort study. <i>BMC Psychiatry</i> , 2013 , 13, 262	4.2	14
107	Adherence, persistence and switch rates for anticholinergic drugs used for overactive bladder in women: data from the Norwegian Prescription Database. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 2013 , 92, 1208-15	3.8	31
106	Laxative prescriptions to cancer outpatients receiving opioids: a study from the Norwegian prescription database. <i>Supportive Care in Cancer</i> , 2013 , 21, 67-73	3.9	16
105	Drug exposure: inclusion of dispensed drugs before pregnancy may lead to underestimation of risk associations. <i>Journal of Clinical Epidemiology</i> , 2013 , 66, 964-72	5.7	29
104	A pharmacoepidemiological cohort study of subjects starting strong opioids for nonmalignant pain: a study from the Norwegian Prescription Database. <i>Pain</i> , 2013 , 154, 2487-2493	8	28
103	Prescription drug use among pregnant women in opioid Maintenance Treatment. <i>Addiction</i> , 2013 , 108, 367-76	4.6	20
102	Effects of preconceptional paternal drug exposure on birth outcomes: cohort study of 340 000 pregnancies using Norwegian population-based databases. <i>British Journal of Clinical Pharmacology</i> , 2013 , 75, 1134-41	3.8	28
101	Neonatal outcomes following in utero exposure to methadone or buprenorphine: a National Cohort Study of opioid-agonist treatment of Pregnant Women in Norway from 1996 to 2009. <i>Drug and Alcohol Dependence</i> , 2013 , 127, 200-6	4.9	78
100	Driving under the influence (DUI) among patients in opioid maintenance treatment (OMT): a registry-based national cohort study. <i>Addiction</i> , 2013 , 108, 1954-61	4.6	3
99	Breastfeeding reduces the need for withdrawal treatment in opioid-exposed infants. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2013 , 102, 1060-6	3.1	47
98	Association between prescribing hypnotics for parents and children in Norway. <i>Archives of Disease in Childhood</i> , 2013 , 98, 732-6	2.2	9

(2012-2013)

97	A 2-year prospective study of psychological distress among a national cohort of pregnant women in opioid maintenance treatment and their partners. <i>Journal of Substance Use</i> , 2013 , 18, 148-160	0.8	7
96	Multiple prescribers in older frequent opioid usersdoes it mean abuse?. <i>Journal of Population Therapeutics and Clinical Pharmacology</i> , 2013 , 20, e397-405	3.3	7
95	Concomitant medication among persistent opioid users with chronic non-malignant pain. <i>Acta Anaesthesiologica Scandinavica</i> , 2012 , 56, 1267-76	1.9	29
94	What is the point of guidelines? Benzodiazepine and z-hypnotic use by an elderly population. <i>Sleep Medicine</i> , 2012 , 13, 893-7	4.6	43
93	Mental distress and subsequent use of psychotropic drugs among adolescents-a prospective register linkage study. <i>Journal of Adolescent Health</i> , 2012 , 50, 578-87	5.8	8
92	Differential patterns of opioid use: defining persistent opioid use in a prescription database. <i>European Journal of Pain</i> , 2012 , 16, 359-69	3.7	32
91	Opioid switching to methadone: a pharmacoepidemiological study from a national prescription database. <i>Palliative Medicine</i> , 2012 , 26, 804-12	5.5	7
90	Increase in psychotropic drug use between 2006 and 2010 among adolescents in Norway: a nationwide prescription database study. <i>Clinical Epidemiology</i> , 2012 , 4, 225-31	5.9	20
89	Polypharmacy of potentially addictive medication in the older personsquantifying usage. <i>Pharmacoepidemiology and Drug Safety</i> , 2012 , 21, 199-206	2.6	9
88	Comorbidities in an asthma population 8-29 years old: a study from the Norwegian Prescription Database. <i>Pharmacoepidemiology and Drug Safety</i> , 2012 , 21, 1045-52	2.6	19
87	Influence of socioeconomic factors on the adherence of alendronate treatment in incident users in Norway. <i>Pharmacoepidemiology and Drug Safety</i> , 2012 , 21, 297-304	2.6	24
86	Low-dose transdermal buprenorphine - long-term use and co-medication with other potentially addictive drugs. <i>Acta Anaesthesiologica Scandinavica</i> , 2012 , 56, 88-94	1.9	7
85	Engagement with opioid maintenance treatment and reductions in crime: a longitudinal national cohort study. <i>Addiction</i> , 2012 , 107, 393-9	4.6	78
84	An increased risk of motor vehicle accidents after prescription of methadone. <i>Addiction</i> , 2012 , 107, 967	-7426	19
83	Hypnotic drug use among 0-17 year olds during 2004-2011: a nationwide prescription database study. <i>Scandinavian Journal of Public Health</i> , 2012 , 40, 704-11	3	34
82	Effect of the market withdrawal of carisoprodol on use of other prescribed drugs with abuse potential. <i>Clinical Pharmacology and Therapeutics</i> , 2012 , 91, 438-41	6.1	12
81	Substance use during and after pregnancy among a national cohort of pregnant women in opioid maintenance treatment and their partners. <i>Journal of Substance Use</i> , 2012 , 17, 277-286	0.8	12
8o	Reimbursement of analgesics for chronic pain. <i>Tidsskrift for Den Norske Laegeforening</i> , 2012 , 132, 2489-	-9335	3

79	Dispensing of prescribed analgesics in Norway among young people with foreign-or Norwegian-born parents. <i>Scandinavian Journal of Pain</i> , 2011 , 2, 36-44	1.9	2
78	Prescription of analgesics to patients in opioid maintenance therapy: a pharmacoepidemiological study. <i>Drug and Alcohol Dependence</i> , 2011 , 116, 158-62	4.9	10
77	High validity of mother-reported use of antiasthmatics among children: a comparison with a population-based prescription database. <i>Journal of Clinical Epidemiology</i> , 2011 , 64, 878-84	5.7	39
76	Criminal convictions among dependent heroin users during a 3-year period prior to opioid maintenance treatment: a longitudinal national cohort study. <i>Journal of Substance Abuse Treatment</i> , 2011 , 41, 407-14	4.2	41
75	To what extent does a cohort of new users of weak opioids develop persistent or probable problematic opioid use?. <i>Pain</i> , 2011 , 152, 1555-1561	8	30
74	Risk of diabetes after gestational diabetes and preeclampsia. A registry-based study of 230,000 women in Norway. <i>European Journal of Epidemiology</i> , 2011 , 26, 157-63	12.1	52
73	Use of prescribed opioid analgesics and co-medication with benzodiazepines in women before, during, and after pregnancy: a population-based cohort study. <i>European Journal of Clinical Pharmacology</i> , 2011 , 67, 953-60	2.8	29
72	The association between smoking and subsequent repeated use of prescribed opioids among adolescents and young adultsa population-based cohort study. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 90-8	2.6	17
71	Prevalence, incidence and persistence of etanercept and adalimumab in Norway 2005-2009. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 457-63	2.6	4
70	A prospective study of asthma and subsequent use of hypnotics in young adults. <i>Pharmacoepidemiology and Drug Safety</i> , 2011 , 20, 370-7	2.6	1
69	Prescription of antidepressants to patients on opioid maintenance therapy D pharmacoepidemiological study. <i>Norsk Epidemiologi</i> , 2011 , 21,	2.8	3
68	Does Pregabalin (Lyrica([])) help patients reduce their use of benzodiazepines? A comparison with gabapentin using the Norwegian Prescription Database. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2010 , 107, 883-6	3.1	21
67	Mental distress and subsequent use of anxiolytic drugs—a prospective population-based cohort study of 16,000 individuals. <i>Scandinavian Journal of Public Health</i> , 2010 , 38, 465-73	3	4
66	How are opioids used in Norway? Persistent use, utilization of depot formulation and age profile in non-palliation patients. <i>Scandinavian Journal of Pain</i> , 2010 , 1, 171-171	1.9	
65	Nicotine dependence predicts repeated use of prescribed opioids. Prospective population-based cohort study. <i>Annals of Epidemiology</i> , 2010 , 20, 890-7	6.4	36
64	Benzodiazepines predict use of opioidsa follow-up study of 17,074 men and women. <i>Pain Medicine</i> , 2010 , 11, 805-14	2.8	41
63	Prescriptions of opioids to children and adolescents; a study from a national prescription database in Norway. <i>Paediatric Anaesthesia</i> , 2010 , 20, 537-44	1.8	14
62	Prevalence, incidence and persistence of anti-asthma medication use in 2- to 29-year-olds: a nationwide prescription study. <i>European Journal of Clinical Pharmacology</i> , 2010 , 66, 399-406	2.8	21

(2008-2010)

61	Disability pension as predictor of later use of benzodiazepines among benzodiazepine users. <i>Social Science and Medicine</i> , 2010 , 70, 921-5	5.1	3	
60	Increasing use of opioids from 2004 to 2007 - pharmacoepidemiological data from a complete national prescription database in Norway. <i>European Journal of Pain</i> , 2010 , 14, 289-94	3.7	72	
59	Disability pension is associated with the use of benzodiazepines 20 years later: a prospective study. <i>Scandinavian Journal of Public Health</i> , 2009 , 37, 320-6	3	6	
58	Starting insomnia treatment: the use of benzodiazepines versus z-hypnotics. A prescription database study of predictors. <i>European Journal of Clinical Pharmacology</i> , 2009 , 65, 295-301	2.8	44	
57	An increased risk of road traffic accidents after prescriptions of lithium or valproate?. <i>Pharmacoepidemiology and Drug Safety</i> , 2009 , 18, 492-6	2.6	12	
56	Co-medication of statins and CYP3A4 inhibitors before and after introduction of new reimbursement policy. <i>British Journal of Clinical Pharmacology</i> , 2009 , 67, 234-41	3.8	16	
55	Statin treatment in a cohort of 20 212 men and women in Norway according to cardiovascular risk factors and level of education. <i>British Journal of Clinical Pharmacology</i> , 2009 , 67, 355-62	3.8	21	
54	The risk of motor vehicle accidents involving drivers with prescriptions for codeine or tramadol. <i>Clinical Pharmacology and Therapeutics</i> , 2009 , 85, 596-9	6.1	39	
53	Prescription pattern of codeine for non-malignant pain: a pharmacoepidemiological study from the Norwegian Prescription Database. <i>Acta Anaesthesiologica Scandinavica</i> , 2009 , 53, 627-33	1.9	29	
52	Road traffic accident risk in patients with diabetes mellitus receiving blood glucose-lowering drugs. Prospective follow-up study. <i>Diabetic Medicine</i> , 2009 , 26, 404-8	3.5	29	
51	Introduction of low dose transdermal buprenorphine did it influence use of potentially addictive drugs in chronic non-malignant pain patients?. <i>European Journal of Pain</i> , 2009 , 13, 949-53	3.7	14	
50	The absence of evidence is not the same as evidence for absence!. <i>Addiction</i> , 2008 , 103, 513-4	4.6	1	
49	Prescription drug use among fathers and mothers before and during pregnancy. A population-based cohort study of 106,000 pregnancies in Norway 2004-2006. <i>British Journal of Clinical Pharmacology</i> , 2008 , 65, 653-60	3.8	138	
48	Behavioural sensitization in mice induced by morphine-glucuronide metabolites. <i>Pharmacology Biochemistry and Behavior</i> , 2008 , 90, 578-85	3.9	7	
47	The validity of self-reported prescription medication use among adolescents varied by therapeutic class. <i>Journal of Clinical Epidemiology</i> , 2008 , 61, 714-7	5.7	28	
46	Road traffic accident risk related to prescriptions of the hypnotics zopiclone, zolpidem, flunitrazepam and nitrazepam. <i>Sleep Medicine</i> , 2008 , 9, 818-22	4.6	116	
45	Environmental assessment of Norwegian priority pharmaceuticals based on the EMEA guideline. <i>Ecotoxicology and Environmental Safety</i> , 2008 , 71, 328-40	7	155	
44	Benzodiazepine use in all alcohol consumers predicts use of opioids in patients 20 years latera follow-up study of 13,390 men and women aged 40-42 years. <i>Pharmacoepidemiology and Drug Safety</i> , 2008 , 17, 926-933	2.6	13	

43	Minor increase in risk of road traffic accidents after prescriptions of antidepressants: a study of population registry data in Norway. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, 1099-103	4.6	45
42	Carisoprodol should be taken off the market. Southern Medical Journal, 2008, 101, 1074-5	0.6	2
41	The risk of traffic accidents after prescriptions of carisoprodol. <i>Accident Analysis and Prevention</i> , 2007 , 39, 1050-5	6.1	28
40	Aspects of statin prescribing in Norwegian counties with high, average and low statin consumption - an individual-level prescription database study. <i>BMC Clinical Pharmacology</i> , 2007 , 7, 14		18
39	Use of anxiolytic or hypnotic drugs and total mortality in a general middle-aged population. <i>Pharmacoepidemiology and Drug Safety</i> , 2007 , 16, 913-8	2.6	42
38	Carisoprodol use and abuse in Norway: a pharmacoepidemiological study. <i>British Journal of Clinical Pharmacology</i> , 2007 , 64, 210-8	3.8	44
37	Switching statins in Norway after new reimbursement policy: a nationwide prescription study. <i>British Journal of Clinical Pharmacology</i> , 2007 , 64, 476-81	3.8	63
36	Morphine-3-glucuronide inhibits morphine induced, but enhances morphine-6-glucuronide induced locomotor activity in mice. <i>Pharmacology Biochemistry and Behavior</i> , 2007 , 86, 576-86	3.9	7
35	Women@ experiences of giving birth in Northwest Russia in 2000 and 2002 and in Northern Norway 2000. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 2007 , 86, 373-5	3.8	
34	Use of anti-asthmatic medications as a proxy for prevalence of asthma in children and adolescents in Norway: a nationwide prescription database analysis. <i>European Journal of Clinical Pharmacology</i> , 2007 , 63, 693-8	2.8	52
33	Occupational physical activity, overweight, and mortality: a follow-up study of 47,405 Norwegian women and men. <i>Research Quarterly for Exercise and Sport</i> , 2007 , 78, 151-61	1.9	6
32	Psychotropic drug use among persons with mental distress symptoms: a population-based study in Norway. <i>Scandinavian Journal of Public Health</i> , 2007 , 35, 356-64	3	15
31	Risk of road traffic accidents associated with the prescription of drugs: a registry-based cohort study. <i>Annals of Epidemiology</i> , 2007 , 17, 597-602	6.4	135
30	Heroin-using drivers: importance of morphine and morphine-6-glucuronide on late clinical impairment. <i>European Journal of Clinical Pharmacology</i> , 2006 , 62, 905-12	2.8	22
29	Why do sales of lipid-lowering drugs vary between counties in Norway? Evidence from the OPPHED Health Study 2000-2001. <i>Scandinavian Journal of Primary Health Care</i> , 2006 , 24, 115-21	2.7	2
28	665 ALMOST ALL CHRONIC NON-MALIGNANT PAIN PATIENTS ON LONG-TERM OPIOIDS ARE USING SHORT-ACTING MEDICATION. <i>European Journal of Pain</i> , 2006 , 10, S174-S174	3.7	
27	Flunitrazepam: psychomotor impairment, agitation and paradoxical reactions. <i>Forensic Science International</i> , 2006 , 159, 83-91	2.6	26
26	Mortality among subjects previously apprehended for driving under the influence of traffic-hazardous medicinal drugs. <i>Drug and Alcohol Dependence</i> , 2005 , 79, 423-9	4.9	17

(2002-2005)

25	Drinking pattern among adolescents with immigrant and Norwegian backgrounds: a two-way influence?. <i>Addiction</i> , 2005 , 100, 1453-63	4.6	57
24	The CYP2C19 genotype and the use of oral contraceptives influence the pharmacokinetics of carisoprodol in healthy human subjects. <i>European Journal of Clinical Pharmacology</i> , 2005 , 61, 499-506	2.8	13
23	Use of psychotropic drugs in an urban adolescent population: the impact of health-related variables, lifestyle and sociodemographic factorsThe Oslo Health Study 2000-2001. <i>Pharmacoepidemiology and Drug Safety</i> , 2005 , 14, 277-83	2.6	15
22	Fast quantification of ethanol in whole blood specimens by the enzymatic alcohol dehydrogenase method. Optimization by experimental design. <i>Journal of Analytical Toxicology</i> , 2005 , 29, 66-70	2.9	23
21	Characteristics of drivers testing positive for heroin or ecstasy in Norway. <i>Traffic Injury Prevention</i> , 2004 , 5, 107-11	1.8	10
20	Determinants of control of high blood pressure. The Oslo Health Study 2000-2001. <i>Blood Pressure</i> , 2004 , 13, 343-9	1.7	12
19	Undertreatment and overtreatment with statins: the Oslo Health Study 2000-2001. <i>Journal of Internal Medicine</i> , 2004 , 255, 494-502	10.8	55
18	Impairment due to intake of carisoprodol. <i>Drug and Alcohol Dependence</i> , 2004 , 74, 311-8	4.9	38
17	. Pharmacogenetics and Genomics, 2003 , 13, 383-388		23
16	Codeine and clinical impairment in samples in which morphine is not detected. <i>European Journal of Clinical Pharmacology</i> , 2003 , 58, 785-9	2.8	28
15	Testing for benzodiazepine inebriationrelationship between benzodiazepine concentration and simple clinical tests for impairment in a sample of drugged drivers. <i>European Journal of Clinical Pharmacology</i> , 2003 , 59, 593-601	2.8	46
14	Erratum to "Sex differences in gamma-glutamyltransferase in people aged 40-42 years in two Norwegian counties". <i>Drug and Alcohol Dependence</i> , 2003 , 69, 103-6	4.9	3
13	Coffee intake and mortality from liver cirrhosis. <i>Annals of Epidemiology</i> , 2003 , 13, 419-23	6.4	145
12	Association between blood carisoprodol:meprobamate concentration ratios and CYP2C19 genotype in carisoprodol-drugged drivers: decreased metabolic capacity in heterozygous CYP2C19*1/CYP2C19*2 subjects?. <i>Pharmacogenetics and Genomics</i> , 2003 , 13, 383-8		5
11	Pharmacokinetic differences of morphine and morphine-glucuronides are reflected in locomotor activity. <i>Pharmacology Biochemistry and Behavior</i> , 2002 , 73, 883-92	3.9	58
10	Drugged drivers in Norway with benzodiazepine detections. <i>Forensic Science International</i> , 2002 , 125, 75-82	2.6	28
9	Rearrest rates among Norwegian drugged drivers compared with drunken drivers. <i>Drug and Alcohol Dependence</i> , 2002 , 66, 85-92	4.9	52
8	Sex differences in gamma-glutamyltransferase in people aged 40-42 years in two Norwegian counties. <i>Drug and Alcohol Dependence</i> , 2002 , 67, 95-8	4.9	9

7	Increased mortality among previously apprehended drunken and drugged drivers. <i>Drug and Alcohol Dependence</i> , 2002 , 68, 143-50	4.9	36	
6	Clinical impairment of benzodiazepinesrelation between benzodiazepine concentrations and impairment in apprehended drivers. <i>Drug and Alcohol Dependence</i> , 2002 , 68, 131-41	4.9	84	
5	Lack of crosstolerance between morphine and morphine-6-glucuronide as revealed by locomotor activity. <i>Pharmacology Biochemistry and Behavior</i> , 2000 , 66, 205-10	3.9	12	
4	Morphine-6-glucuronide-induced locomotor stimulation in mice: role of opioid receptors. <i>Basic and Clinical Pharmacology and Toxicology</i> , 1998 , 82, 3-10		18	
3	Study of rearrests for drunken driving in Norway. Forensic Science International, 1998, 92, 21-8	2.6	17	
2	Recidivism among drunken and drugged drivers in Norway. <i>Alcohol and Alcoholism</i> , 1996 , 31, 609-11	3.5	10	
1	Female drivers suspected for drunken or drugged driving. <i>Forensic Science International</i> , 1995 , 75, 139-4	18 .6	28	