Filippo Valbusa

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1210194/publications.pdf Version: 2024-02-01



FILIDDO VALBUSA

#	Article	IF	CITATIONS
1	Treatment With Multiple Blood Pressure Medications, Achieved Blood Pressure, and Mortality in Older Nursing Home Residents. JAMA Internal Medicine, 2015, 175, 989.	5.1	225
2	Mortality and Cardiovascular Events Are Best Predicted by Low Central/Peripheral Pulse Pressure Amplification But Not by High Blood Pressure Levels in Elderly Nursing Home Subjects. Journal of the American College of Cardiology, 2012, 60, 1503-1511.	2.8	156
3	Non-Alcoholic Fatty Liver Disease Is Associated with an Increased Incidence of Atrial Fibrillation in Patients with Type 2 Diabetes. PLoS ONE, 2013, 8, e57183.	2.5	153
4	Chronic Plaque Psoriasis Is Associated with Increased Arterial Stiffness. Dermatology, 2009, 218, 110-113.	2.1	109
5	Comparative study of methodologies for pulse wave velocity estimation. Journal of Human Hypertension, 2008, 22, 669-677.	2.2	108
6	Nonalcoholic Fatty Liver Disease Is Associated With Ventricular Arrhythmias in Patients With Type 2 Diabetes Referred for Clinically Indicated 24-Hour Holter Monitoring. Diabetes Care, 2016, 39, 1416-1423.	8.6	95
7	Heart valve calcification in patients with type 2 diabetes and nonalcoholic fatty liver disease. Metabolism: Clinical and Experimental, 2015, 64, 879-887.	3.4	82
8	Nonalcoholic Fatty Liver Disease Is Independently Associated with Early Left Ventricular Diastolic Dysfunction in Patients with Type 2 Diabetes. PLoS ONE, 2015, 10, e0135329.	2.5	81
9	Orthostatic hypotension in very old individuals living in nursing homes. Journal of Hypertension, 2012, 30, 53-60.	0.5	78
10	Association of nonalcoholic fatty liver disease with QTc interval in patients with type 2 diabetes. Nutrition, Metabolism and Cardiovascular Diseases, 2014, 24, 663-669.	2.6	77
11	Relation of Elevated Serum Uric Acid Levels to Incidence of Atrial Fibrillation in Patients With Type 2 Diabetes Mellitus. American Journal of Cardiology, 2013, 112, 499-504.	1.6	58
12	Evidence for a Prognostic Role of Orthostatic Hypertension on Survival in a Very Old Institutionalized Population. Hypertension, 2016, 67, 191-196.	2.7	55
13	Simultaneous Measurement of Beat-to-Beat Carotid Diameter and Pressure Changes to Assess Arterial Mechanical Properties. Hypertension, 2008, 52, 896-902.	2.7	54
14	Nonalcoholic Fatty Liver Disease Is Associated with Aortic Valve Sclerosis in Patients with Type 2 Diabetes Mellitus. PLoS ONE, 2014, 9, e88371.	2.5	49
15	Blood pressure and pulse wave velocity values in the institutionalized elderly aged 80 and over: baseline of the PARTAGE study. Journal of Hypertension, 2010, 28, 41-50.	0.5	46
16	Non-alcoholic fatty liver disease and increased risk of all-cause mortality in elderly patients admitted for acute heart failure. International Journal of Cardiology, 2018, 265, 162-168.	1.7	41
17	Nonalcoholic fatty liver disease and increased risk of 1-year all-cause and cardiac hospital readmissions in elderly patients admitted for acute heart failure. PLoS ONE, 2017, 12, e0173398.	2.5	38
18	Increased Aortic Pulse Wave Velocity as Measured by Echocardiography Is Strongly Associated with Poor Prognosis in Patients with Heart Failure. Journal of the American Society of Echocardiography, 2013, 26, 714-720.	2.8	31

FILIPPO VALBUSA

#	Article	IF	CITATIONS
19	Early impairment in left ventricular longitudinal systolic function is associated with an increased risk of incident atrial fibrillation in patients with type 2 diabetes. Journal of Diabetes and Its Complications, 2017, 31, 413-418.	2.3	24
20	Do Arterial Hemodynamic Parameters Predict Cognitive Decline Over a Period of 2ÂYears in Individuals Older Than 80ÂYears Living in Nursing Homes? The PARTAGE Study. Journal of the American Medical Directors Association, 2015, 16, 598-602.	2.5	23
21	Increased Pulse Pressure Independently Predicts Incident Atrial Fibrillation in Patients With Type 2 Diabetes. Diabetes Care, 2012, 35, 2337-2339.	8.6	20
22	Nonalcoholic Fatty Liver Disease Is Associated With Higher 1-year All-Cause Rehospitalization Rates in Patients Admitted for Acute Heart Failure. Medicine (United States), 2016, 95, e2760.	1.0	17
23	Non-Invasive Assessment of Arterial Stiffness: Pulse Wave Velocity, Pulse Wave Analysis and Carotid Cross-Sectional Distensibility: Comparison between Methods. Journal of Clinical Medicine, 2022, 11, 2225.	2.4	13
24	A case of bronchiolitis obliterans organising pneumoniae (BOOP) after nine months post-operative irradiation for breast cancer. Age and Ageing, 2008, 37, 235-235.	1.6	5
25	Unusual echocardiographic image of a left circumflex coronary artery with anomalous origin from the right coronary sinus. European Heart Journal, 2016, 37, 792-792.	2.2	2
26	Increased aortic stiffness in adults with chronic indeterminate Chagas disease. PLoS ONE, 2019, 14, e0220689.	2.5	2
27	Inter-atrial shunt inversion by the sitting position in a patient with a patent foramen ovale and acute pulmonary embolism. European Heart Journal, 2014, 35, 1032-1032.	2.2	0
28	Interest of Combined Blood Pressure Measurements in Very Old Frail Subjects: The PARTAGE Study. American Journal of Hypertension, 2018, 31, 950-956.	2.0	0