

# E F Magann

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/12124501/e-f-magann-publications-by-citations.pdf>

**Version:** 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

92  
papers

3,250  
citations

32  
h-index

54  
g-index

92  
ext. papers

3,668  
ext. citations

2.1  
avg, IF

4.39  
L-index

#	Paper	IF	Citations
92	The spectrum of severe preeclampsia: comparative analysis by HELLP (hemolysis, elevated liver enzyme levels, and low platelet count) syndrome classification. <i>American Journal of Obstetrics and Gynecology</i> , <b>1999</b> , 180, 1373-84	6.4	224
91	Pre-pregnancy body mass index and pregnancy outcomes. <i>International Journal of Gynecology and Obstetrics</i> , <b>2006</b> , 95, 242-7	4	186
90	Perinatal outcome and amniotic fluid index in the antepartum and intrapartum periods: A meta-analysis. <i>American Journal of Obstetrics and Gynecology</i> , <b>1999</b> , 181, 1473-8	6.4	178
89	Antepartum corticosteroids: disease stabilization in patients with the syndrome of hemolysis, elevated liver enzymes, and low platelets (HELLP). <i>American Journal of Obstetrics and Gynecology</i> , <b>1994</b> , 171, 1148-53	6.4	174
88	The amniotic fluid index, single deepest pocket, and two-diameter pocket in normal human pregnancy. <i>American Journal of Obstetrics and Gynecology</i> , <b>2000</b> , 182, 1581-8	6.4	134
87	Maternal mortality associated with HELLP (hemolysis, elevated liver enzymes, and low platelets) syndrome. <i>American Journal of Obstetrics and Gynecology</i> , <b>1999</b> , 181, 924-8	6.4	128
86	Postpartum corticosteroids: accelerated recovery from the syndrome of hemolysis, elevated liver enzymes, and low platelets (HELLP). <i>American Journal of Obstetrics and Gynecology</i> , <b>1994</b> , 171, 1154-8	6.4	121
85	Twelve steps to optimal management of HELLP syndrome. <i>Clinical Obstetrics and Gynecology</i> , <b>1999</b> , 42, 532-50	1.7	114
84	Measurement of amniotic fluid volume: accuracy of ultrasonography techniques. <i>American Journal of Obstetrics and Gynecology</i> , <b>1992</b> , 167, 1533-7	6.4	109
83	Mode of delivery for the morbidly obese with prior cesarean delivery: vaginal versus repeat cesarean section. <i>American Journal of Obstetrics and Gynecology</i> , <b>2001</b> , 185, 349-54	6.4	105
82	Early risk assessment of severe preeclampsia: admission battery of symptoms and laboratory tests to predict likelihood of subsequent significant maternal morbidity. <i>American Journal of Obstetrics and Gynecology</i> , <b>1999</b> , 180, 1407-14	6.4	87
81	The effect of placental management at cesarean delivery on operative blood loss. <i>American Journal of Obstetrics and Gynecology</i> , <b>1992</b> , 167, 1363-7	6.4	65
80	Antenatal testing among 1001 patients at high risk: the role of ultrasonographic estimate of amniotic fluid volume. <i>American Journal of Obstetrics and Gynecology</i> , <b>1999</b> , 180, 1330-6	6.4	62
79	A prospective, randomized trial comparing the efficacy of dexamethasone and betamethasone for the treatment of antepartum HELLP (hemolysis, elevated liver enzymes, and low platelet count) syndrome. <i>American Journal of Obstetrics and Gynecology</i> , <b>2001</b> , 184, 1332-7; discussion 1337-9	6.4	61
78	Employment, exertion, and pregnancy outcome: assessment by kilocalories expended each day. <i>American Journal of Obstetrics and Gynecology</i> , <b>1996</b> , 175, 182-7	6.4	54
77	Amniotic fluid index and single deepest pocket: weak indicators of abnormal amniotic volumes. <i>Obstetrics and Gynecology</i> , <b>2000</b> , 96, 737-40	4.9	51
76	Does an amniotic fluid index of . <i>American Journal of Obstetrics and Gynecology</i> , <b>1999</b> , 180, 1354-9	6.4	49

75	Ultrasonographic assessment of amniotic fluid does not reflect actual amniotic fluid volume. <i>American Journal of Obstetrics and Gynecology</i> , <b>1997</b> , 177, 291-6; discussion 296-7	6.4	46
74	Intrapartum detection of a macrosomic fetus: clinical versus 8 sonographic models. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , <b>1995</b> , 35, 266-70	1.7	46
73	The accuracy of ultrasound evaluation of amniotic fluid volume in singleton pregnancies: the effect of operator experience and ultrasound interpretative technique. <i>Journal of Clinical Ultrasound</i> , <b>1997</b> , 25, 249-53	1	45
72	Obstetric challenges of massive obesity complicating pregnancy. <i>Journal of Perinatology</i> , <b>1994</b> , 14, 10-4	3.1	45
71	Amniotic fluid volume in normal singleton pregnancies. <i>Obstetrics and Gynecology</i> , <b>1997</b> , 90, 524-8	4.9	43
70	Comparative efficacy of two sonographic measurements for the detection of aberrations in the amniotic fluid volume and the effect of amniotic fluid volume on pregnancy outcome. <i>Obstetrics and Gynecology</i> , <b>1994</b> , 83, 959-62	4.9	42
69	The interrelationship of eclampsia, HELLP syndrome, and prematurity: cofactors for significant maternal and perinatal risk. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , <b>1993</b> , 100, 1095-100	3.7	42
68	Infectious morbidity, operative blood loss, and length of the operative procedure after cesarean delivery by method of placental removal and site of uterine repair. <i>Journal of the American College of Surgeons</i> , <b>1995</b> , 181, 517-20	4.4	41
67	Second trimester placental location as a predictor of an adverse pregnancy outcome. <i>Journal of Perinatology</i> , <b>2007</b> , 27, 9-14	3.1	40
66	Intrapartum oligohydramnios does not predict adverse peripartum outcome among high-risk parturients. <i>American Journal of Obstetrics and Gynecology</i> , <b>1997</b> , 176, 1130-6; discussion 1136-8	6.4	38
65	Corticosteroids for the enhancement of fetal lung maturity: impact on the gravida with preeclampsia and the HELLP syndrome. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , <b>1993</b> , 33, 127-31	1.7	38
64	Management of pregnancies beyond forty-one weeks gestation with an unfavorable cervix. <i>American Journal of Obstetrics and Gynecology</i> , <b>1998</b> , 178, 1279-87	6.4	37
63	Immediate postpartum curettage: accelerated recovery from severe preeclampsia. <i>Obstetrics and Gynecology</i> , <b>1993</b> , 81, 502-6	4.9	36
62	A randomized study to assess the efficacy of the amniotic fluid index as a fetal admission test. <i>Obstetrics and Gynecology</i> , <b>1995</b> , 86, 9-13	4.9	32
61	Blood loss at time of cesarean section by method of placental removal and exteriorization versus in situ repair of the uterine incision. <i>Surgery, Gynecology &amp; Obstetrics</i> , <b>1993</b> , 177, 389-92		32
60	Clinical relevance of sonographically estimated amniotic fluid volume: polyhydramnios. <i>Journal of Ultrasound in Medicine</i> , <b>2013</b> , 32, 851-63	2.9	30
59	Cervical ripening before medical induction of labor: a comparison of prostaglandin E2, estradiol, and oxytocin. <i>American Journal of Obstetrics and Gynecology</i> , <b>1995</b> , 172, 1702-6; discussion 1704-8	6.4	29
58	Preoperative skin preparation and intraoperative pelvic irrigation: impact on post-cesarean endometritis and wound infection. <i>Obstetrics and Gynecology</i> , <b>1993</b> , 81, 922-5	4.9	27

57	Dexamethasone compared with betamethasone for glucocorticoid treatment of postpartum HELLP syndrome. <i>International Journal of Gynecology and Obstetrics</i> , <b>2003</b> , 80, 291-7	4	25
56	The effects of physical abuse on pregnancy outcomes in a low-risk obstetric population. <i>American Journal of Obstetrics and Gynecology</i> , <b>2000</b> , 182, 1067-9	6.4	25
55	ADHD and developmental speech/language disorders in late preterm, early term and term infants. <i>Journal of Perinatology</i> , <b>2015</b> , 35, 660-4	3.1	24
54	Corticosteroids for enhanced fetal lung maturation in patients with HELLP syndrome: impact on neonates. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , <b>1993</b> , 33, 131-5	1.7	23
53	Amniotic fluid volume estimation and the biophysical profile: a confusion of criteria. <i>Obstetrics and Gynecology</i> , <b>2000</b> , 96, 640-2	4.9	22
52	Acute tocolysis for fetal distress: terbutaline versus magnesium sulphate. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , <b>1993</b> , 33, 362-4	1.7	22
51	The laboratory evaluation of hypertensive gravidas. <i>Obstetrical and Gynecological Survey</i> , <b>1995</b> , 50, 138-44	4.4	22
50	Effectiveness and safety of a structured swimming program in previously sedentary women during pregnancy. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , <b>2003</b> , 14, 163-9	2	21
49	Accelerated recovery from severe preeclampsia: uterine curettage versus nifedipine. <i>Journal of the Society for Gynecologic Investigation</i> , <b>1994</b> , 1, 210-4		21
48	Neonatal salvage by week of gestation in pregnancies complicated by HELLP syndrome. <i>Journal of the Society for Gynecologic Investigation</i> , <b>1994</b> , 1, 206-9		20
47	Can we decrease postdatism in women with an unfavorable cervix and a negative fetal fibronectin test result at term by serial membrane sweeping?. <i>American Journal of Obstetrics and Gynecology</i> , <b>1998</b> , 179, 890-4	6.4	19
46	Membrane sweeping versus dinoprostone vaginal insert in the management of pregnancies beyond 41 weeks with an unfavorable cervix. <i>Journal of Perinatology</i> , <b>1999</b> , 19, 88-91	3.1	19
45	Intrapartum amniotic fluid index. A poor diagnostic test for adverse perinatal outcome. <i>Journal of reproductive medicine, The</i> , <b>1996</b> , 41, 860-6		19
44	Amniotic fluid volume of third-trimester diamniotic twin pregnancies. <i>Obstetrics and Gynecology</i> , <b>1995</b> , 85, 957-60	4.9	18
43	Standard parameters of preeclampsia: can the clinician depend upon them to reliably identify the patient with the HELLP syndrome?. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , <b>1993</b> , 33, 122-6	1.7	17
42	Cardiopulmonary morbidity as a complication of severe preeclampsia HELLP syndrome. <i>Journal of Perinatology</i> , <b>2000</b> , 20, 78-81	3.1	16
41	Amniotic fluid volume assessment in singleton and twin pregnancies. <i>Obstetrics and Gynecology Clinics of North America</i> , <b>1999</b> , 26, 579-93	3.3	16
40	Dye-dilution techniques using aminohippurate sodium: do they accurately reflect amniotic fluid volume?. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , <b>2002</b> , 11, 167-70	2	15

39	Placental implantation at 18 weeks and migration throughout pregnancy. <i>Southern Medical Journal</i> , <b>1998</b> , 91, 1025-7	0.6	15
38	Childhood Respiratory Morbidity after Late Preterm and Early Term Delivery: a Study of Medicaid Patients in South Carolina. <i>Paediatric and Perinatal Epidemiology</i> , <b>2016</b> , 30, 67-75	2.7	15
37	The accuracy of the summated amniotic fluid index in evaluating amniotic fluid volume in twin pregnancies. <i>American Journal of Obstetrics and Gynecology</i> , <b>1997</b> , 177, 1041-5	6.4	14
36	Critical care of HELLP syndrome with corticosteroids. <i>American Journal of Perinatology</i> , <b>2000</b> , 17, 417-22	3.3	14
35	Corticosteroid-induced arrest of HELLP syndrome progression in a marginally-viable pregnancy. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , <b>1995</b> , 59, 217-9	2.4	14
34	Ultrasonic assessment of the amniotic fluid volume in diamniotic twins. <i>Journal of the Society for Gynecologic Investigation</i> , <b>1995</b> , 2, 609-13		14
33	Pet therapy program for antepartum high-risk pregnancies: a pilot study. <i>Journal of Perinatology</i> , <b>2014</b> , 34, 816-8	3.1	13
32	Ultrasound estimate of amniotic fluid volume: color Doppler overdiagnosis of oligohydramnios. <i>Obstetrics and Gynecology</i> , <b>2001</b> , 98, 71-4	4.9	13
31	Does method of placental removal or site of uterine incision repair alter endometritis after cesarean delivery?. <i>Infectious Diseases in Obstetrics and Gynecology</i> , <b>1993</b> , 1, 65-70	2.4	13
30	Determination of amniotic fluid volume in twin pregnancies: ultrasonographic evaluation versus operator estimation. <i>American Journal of Obstetrics and Gynecology</i> , <b>2000</b> , 182, 1606-9	6.4	12
29	Pregnancy outcome in an active-duty population. <i>Obstetrics and Gynecology</i> , <b>1991</b> , 78, 391-3	4.9	12
28	Climatic factors and preeclampsia-related hypertensive disorders of pregnancy. <i>American Journal of Obstetrics and Gynecology</i> , <b>1995</b> , 172, 204-5	6.4	11
27	Outpatient cervical ripening with prostaglandin E2 and estradiol. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , <b>2002</b> , 11, 113-7	2	10
26	Low amniotic fluid volume is poorly identified in singleton and twin pregnancies using the 2 x 2 cm pocket technique of the biophysical profile. <i>Southern Medical Journal</i> , <b>1999</b> , 92, 802-5	0.6	10
25	Detection of growth-restricted fetuses in preeclampsia: a case-control study. <i>Obstetrics and Gynecology</i> , <b>1999</b> , 93, 687-91	4.9	10
24	Risk factors for secondary arrest of labor among women >41 weeks gestation with an unfavorable cervix undergoing membrane sweeping for cervical ripening. <i>International Journal of Gynecology and Obstetrics</i> , <b>1999</b> , 65, 1-5	4	8
23	Subjective versus objective evaluation of amniotic fluid volume of pregnancies of less than 24 weeks gestation: how can we be accurate?. <i>Journal of Ultrasound in Medicine</i> , <b>2001</b> , 20, 191-5	2.9	7
22	Factors adversely affecting pregnancy outcome in the military. <i>American Journal of Perinatology</i> , <b>1995</b> , 12, 462-6	3.3	7

21	Acute tocolysis for suspected intrapartum fetal distress: maternal effects of terbutaline versus magnesium sulfate. <i>International Journal of Obstetric Anesthesia</i> , <b>1995</b> , 4, 140-4	2.1	7
20	Placenta praevia: does uterine activity cause bleeding?. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , <b>1993</b> , 33, 22-4	1.7	7
19	Cesarean delivery for suspected fetal distress among preterm parturients. <i>Journal of reproductive medicine, The</i> , <b>2000</b> , 45, 395-402		7
18	Domestic violence screening of obstetric triage patients in a military population. <i>Journal of Perinatology</i> , <b>2012</b> , 32, 763-9	3.1	6
17	Detection of fetal growth restriction in patients with chronic hypertension: is it feasible?. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , <b>2003</b> , 14, 324-8	2	6
16	Change in platelet count predicts eventual maternal outcome with syndrome of hemolysis, elevated liver enzymes and low platelet count. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , <b>2001</b> , 10, 28-34	2	6
15	Effect of amniotic fluid volume on neonatal outcome in diamniotic twin pregnancies. <i>Southern Medical Journal</i> , <b>1998</b> , 91, 942-5	0.6	6
14	Intrapartum amniotic fluid index and two-diameter pocket are poor predictors of adverse neonatal outcome. <i>Journal of Perinatology</i> , <b>1997</b> , 17, 221-4	3.1	6
13	Invalid equation to describe the amniotic fluid index in normal pregnancy. <i>American Journal of Obstetrics and Gynecology</i> , <b>1994</b> , 170, 1209-10	6.4	5
12	Maternal morbidity and mortality associated with intrauterine fetal demise: five-year experience in a tertiary referral hospital. <i>Southern Medical Journal</i> , <b>2001</b> , 94, 493-5	0.6	5
11	Ultrasound estimation of amniotic fluid volume using the largest vertical pocket containing umbilical cord: measure to or through the cord?. <i>Ultrasound in Obstetrics and Gynecology</i> , <b>2002</b> , 20, 464-7 <sup>5.8</sup>		4
10	Does uterine wiping influence the rate of post-Cesarean endometritis?. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , <b>2001</b> , 10, 318-322	2	4
9	Accuracy of ultrasonography in evaluating amniotic fluid volume at less than 24 weeks of gestation. <i>Journal of Ultrasound in Medicine</i> , <b>1995</b> , 14, 895-7	2.9	4
8	Do multiple measurements employing different ultrasonic techniques improve the accuracy of amniotic fluid volume assessment?. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , <b>1998</b> , 38, 172-5	1.7	2
7	Absence of seasonal variation on the frequency of HELLP syndrome. <i>Southern Medical Journal</i> , <b>1998</b> , 91, 731-2	0.6	2
6	Military pregnancies and adverse perinatal outcome. <i>International Journal of Gynecology and Obstetrics</i> , <b>1996</b> , 52, 19-24	4	2
5	Factors relevant to mode of preterm delivery with syndrome of HELLP (hemolysis, elevated liver enzymes, and low platelets). <i>American Journal of Obstetrics and Gynecology</i> , <b>1994</b> , 170, 1828-32; discussion 1832-4	6.4	2
4	New-onset hypertension in the pregnant patient. <i>Obstetrics and Gynecology Clinics of North America</i> , <b>1995</b> , 22, 157-71	3.3	1

- 3 Does uterine wiping influence the rate of post-Cesarean endometritis?. *The Journal of Maternal-fetal Medicine*, **2001**, 10, 318-22 1
- 2 Lack of utility of amniotic fluid index in predicting perinatal morbidity and mortality. *American Journal of Obstetrics and Gynecology*, **2001**, 184, 778-9 6.4
- 1 Outpatient cervical ripening with prostaglandin E2 and estradiol. *Journal of Maternal-Fetal and Neonatal Medicine*, **2002**, 11, 113-117 2