E F Magann

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/12124501/e-f-magann-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

92	3,250 citations	32	54
papers		h-index	g-index
92 ext. papers	3,668 ext. citations	2. 1 avg, IF	4.39 L-index

#	Paper	IF	Citations
92	Childhood Respiratory Morbidity after Late Preterm and Early Term Delivery: a Study of Medicaid Patients in South Carolina. <i>Paediatric and Perinatal Epidemiology</i> , 2016 , 30, 67-75	2.7	15
91	ADHD and developmental speech/language disorders in late preterm, early term and term infants. <i>Journal of Perinatology</i> , 2015 , 35, 660-4	3.1	24
90	Pet therapy program for antepartum high-risk pregnancies: a pilot study. <i>Journal of Perinatology</i> , 2014 , 34, 816-8	3.1	13
89	Clinical relevance of sonographically estimated amniotic fluid volume: polyhydramnios. <i>Journal of Ultrasound in Medicine</i> , 2013 , 32, 851-63	2.9	30
88	Domestic violence screening of obstetric triage patients in a military population. <i>Journal of Perinatology</i> , 2012 , 32, 763-9	3.1	6
87	Second trimester placental location as a predictor of an adverse pregnancy outcome. <i>Journal of Perinatology</i> , 2007 , 27, 9-14	3.1	40
86	Pre-pregnancy body mass index and pregnancy outcomes. <i>International Journal of Gynecology and Obstetrics</i> , 2006 , 95, 242-7	4	186
85	Effectiveness and safety of a structured swimming program in previously sedentary women during pregnancy. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2003 , 14, 163-9	2	21
84	Detection of fetal growth restriction in patients with chronic hypertension: is it feasible?. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2003 , 14, 324-8	2	6
83	Dexamethasone compared with betamethasone for glucocorticoid treatment of postpartum HELLP syndrome. <i>International Journal of Gynecology and Obstetrics</i> , 2003 , 80, 291-7	4	25
82	Outpatient cervical ripening with prostaglandin E2 and estradiol. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2002 , 11, 113-7	2	10
81	Dye-dilution techniques using aminohippurate sodium: do they accurately reflect amniotic fluid volume?. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2002 , 11, 167-70	2	15
80	Ultrasound estimation of amniotic fluid volume using the largest vertical pocket containing umbilical cord: measure to or through the cord?. <i>Ultrasound in Obstetrics and Gynecology</i> , 2002 , 20, 464-	·7 ^{5.8}	4
79	Outpatient cervical ripening with prostaglandin E2 and estradiol. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2002 , 11, 113-117	2	
78	A prospective, randomized trial comparing the efficacy of dexamethasone and betamethasone for the treatment of antepartum HELLP (hemolysis, elevated liver enzymes, and low platelet count) syndrome. <i>American Journal of Obstetrics and Gynecology</i> , 2001 , 184, 1332-7; discussion 1337-9	6.4	61
77	Mode of delivery for the morbidly obese with prior cesarean delivery: vaginal versus repeat cesarean section. <i>American Journal of Obstetrics and Gynecology</i> , 2001 , 185, 349-54	6.4	105
76	Change in platelet count predicts eventual maternal outcome with syndrome of hemolysis, elevated liver enzymes and low platelet count. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2001 , 10, 28-34	2	6

75	Lack of utility of amniotic fluid index in predicting perinatal morbidity and mortality. <i>American Journal of Obstetrics and Gynecology</i> , 2001 , 184, 778-9	6.4	
74	Ultrasound estimate of amniotic fluid volume: color Doppler overdiagnosis of oligohydramnios. <i>Obstetrics and Gynecology</i> , 2001 , 98, 71-4	4.9	13
73	Subjective versus objective evaluation of amniotic fluid volume of pregnancies of less than 24 weeksUgestation: how can we be accurate?. <i>Journal of Ultrasound in Medicine</i> , 2001 , 20, 191-5	2.9	7
72	Does uterine wiping influence the rate of post-Cesarean endometritis?. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2001 , 10, 318-322	2	4
71	Maternal morbidity and mortality associated with intrauterine fetal demise: five-year experience in a tertiary referral hospital. <i>Southern Medical Journal</i> , 2001 , 94, 493-5	0.6	5
70	Does uterine wiping influence the rate of post-Cesarean endometritis?. <i>The Journal of Maternal-fetal Medicine</i> , 2001 , 10, 318-22		1
69	Cardiopulmonary morbidity as a complication of severe preeclampsia HELLP syndrome. <i>Journal of Perinatology</i> , 2000 , 20, 78-81	3.1	16
68	The effects of physical abuse on pregnancy outcomes in a low-risk obstetric population. <i>American Journal of Obstetrics and Gynecology</i> , 2000 , 182, 1067-9	6.4	25
67	The amniotic fluid index, single deepest pocket, and two-diameter pocket in normal human pregnancy. <i>American Journal of Obstetrics and Gynecology</i> , 2000 , 182, 1581-8	6.4	134
66	Determination of amniotic fluid volume in twin pregnancies: ultrasonographic evaluation versus operator estimation. <i>American Journal of Obstetrics and Gynecology</i> , 2000 , 182, 1606-9	6.4	12
65	Critical care of HELLP syndrome with corticosteroids. <i>American Journal of Perinatology</i> , 2000 , 17, 417-22	23.3	14
64	Amniotic fluid index and single deepest pocket: weak indicators of abnormal amniotic volumes. <i>Obstetrics and Gynecology</i> , 2000 , 96, 737-40	4.9	51
63	Amniotic fluid volume estimation and the biophysical profile: a confusion of criteria. <i>Obstetrics and Gynecology</i> , 2000 , 96, 640-2	4.9	22
62	Cesarean delivery for suspected fetal distress among preterm parturients. <i>Journal of reproductive medicine, The</i> , 2000 , 45, 395-402		7
61	Low amniotic fluid volume is poorly identified in singleton and twin pregnancies using the 2 x 2 cm pocket technique of the biophysical profile. <i>Southern Medical Journal</i> , 1999 , 92, 802-5	0.6	10
60	Membrane sweeping versus dinoprostone vaginal insert in the management of pregnancies beyond 41 weeks with an unfavorable cervix. <i>Journal of Perinatology</i> , 1999 , 19, 88-91	3.1	19
59	Antenatal testing among 1001 patients at high risk: the role of ultrasonographic estimate of amniotic fluid volume. <i>American Journal of Obstetrics and Gynecology</i> , 1999 , 180, 1330-6	6.4	62
58	Does an amniotic fluid index of . <i>American Journal of Obstetrics and Gynecology</i> , 1999 , 180, 1354-9	6.4	49

57	The spectrum of severe preeclampsia: comparative analysis by HELLP (hemolysis, elevated liver enzyme levels, and low platelet count) syndrome classification. <i>American Journal of Obstetrics and Gynecology</i> , 1999 , 180, 1373-84	6.4	224
56	Early risk assessment of severe preeclampsia: admission battery of symptoms and laboratory tests to predict likelihood of subsequent significant maternal morbidity. <i>American Journal of Obstetrics and Gynecology</i> , 1999 , 180, 1407-14	6.4	87
55	Maternal mortality associated with HELLP (hemolysis, elevated liver enzymes, and low platelets) syndrome. <i>American Journal of Obstetrics and Gynecology</i> , 1999 , 181, 924-8	6.4	128
54	Perinatal outcome and amniotic fluid index in the antepartum and intrapartum periods: A meta-analysis. <i>American Journal of Obstetrics and Gynecology</i> , 1999 , 181, 1473-8	6.4	178
53	Amniotic fluid volume assessment in singleton and twin pregnancies. <i>Obstetrics and Gynecology Clinics of North America</i> , 1999 , 26, 579-93	3.3	16
52	Detection of growth-restricted fetuses in preeclampsia: a case-control study. <i>Obstetrics and Gynecology</i> , 1999 , 93, 687-91	4.9	10
51	Risk factors for secondary arrest of labor among women >41 weeksUgestation with an unfavorable cervix undergoing membrane sweeping for cervical ripening. <i>International Journal of Gynecology and Obstetrics</i> , 1999 , 65, 1-5	4	8
50	Twelve steps to optimal management of HELLP syndrome. <i>Clinical Obstetrics and Gynecology</i> , 1999 , 42, 532-50	1.7	114
49	Do multiple measurements employing different ultrasonic techniques improve the accuracy of amniotic fluid volume assessment?. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , 1998 , 38, 172-5	1.7	2
48	Can we decrease postdatism in women with an unfavorable cervix and a negative fetal fibronectin test result at term by serial membrane sweeping?. <i>American Journal of Obstetrics and Gynecology</i> , 1998 , 179, 890-4	6.4	19
47	Management of pregnancies beyond forty-one weeksUgestation with an unfavorable cervix. <i>American Journal of Obstetrics and Gynecology</i> , 1998 , 178, 1279-87	6.4	37
46	Placental implantation at 18 weeks and migration throughout pregnancy. <i>Southern Medical Journal</i> , 1998 , 91, 1025-7	0.6	15
45	Effect of amniotic fluid volume on neonatal outcome in diamniotic twin pregnancies. <i>Southern Medical Journal</i> , 1998 , 91, 942-5	0.6	6
44	Absence of seasonal variation on the frequency of HELLP syndrome. <i>Southern Medical Journal</i> , 1998 , 91, 731-2	0.6	2
43	Amniotic fluid volume in normal singleton pregnancies. <i>Obstetrics and Gynecology</i> , 1997 , 90, 524-8	4.9	43
42	The accuracy of the summated amniotic fluid index in evaluating amniotic fluid volume in twin pregnancies. <i>American Journal of Obstetrics and Gynecology</i> , 1997 , 177, 1041-5	6.4	14
41	Ultrasonographic assessment of amniotic fluid does not reflect actual amniotic fluid volume. <i>American Journal of Obstetrics and Gynecology</i> , 1997 , 177, 291-6; discussion 296-7	6.4	46
40	Intrapartum oligohydramnios does not predict adverse peripartum outcome among high-risk parturients. <i>American Journal of Obstetrics and Gynecology</i> , 1997 , 176, 1130-6; discussion 1136-8	6.4	38

(1995-1997)

39	of operator experience and ultrasound interpretative technique. <i>Journal of Clinical Ultrasound</i> , 1997 , 25, 249-53	1	45	
38	Intrapartum amniotic fluid index and two-diameter pocket are poor predictors of adverse neonatal outcome. <i>Journal of Perinatology</i> , 1997 , 17, 221-4	3.1	6	
37	Military pregnancies and adverse perinatal outcome. <i>International Journal of Gynecology and Obstetrics</i> , 1996 , 52, 19-24	4	2	
36	Employment, exertion, and pregnancy outcome: assessment by kilocalories expended each day. <i>American Journal of Obstetrics and Gynecology</i> , 1996 , 175, 182-7	6.4	54	
35	Intrapartum amniotic fluid index. A poor diagnostic test for adverse perinatal outcome. <i>Journal of reproductive medicine, The</i> , 1996 , 41, 860-6		19	
34	Accuracy of ultrasonography in evaluating amniotic fluid volume at less than 24 weeksUgestation. Journal of Ultrasound in Medicine, 1995 , 14, 895-7	2.9	4	
33	Factors adversely affecting pregnancy outcome in the military. <i>American Journal of Perinatology</i> , 1995 , 12, 462-6	3.3	7	
32	Acute tocolysis for suspected intrapartum fetal distress: maternal effects of terbutaline versus magnesium sulfate. <i>International Journal of Obstetric Anesthesia</i> , 1995 , 4, 140-4	2.1	7	
31	Amniotic fluid volume of third-trimester diamniotic twin pregnancies. <i>Obstetrics and Gynecology</i> , 1995 , 85, 957-60	4.9	18	
30	A randomized study to assess the efficacy of the amniotic fluid index as a fetal admission test. <i>Obstetrics and Gynecology</i> , 1995 , 86, 9-13	4.9	32	
29	Climatic factors and preeclampsia-related hypertensive disorders of pregnancy. <i>American Journal of Obstetrics and Gynecology</i> , 1995 , 172, 204-5	6.4	11	
28	Cervical ripening before medical induction of labor: a comparison of prostaglandin E2, estradiol, and oxytocin. <i>American Journal of Obstetrics and Gynecology</i> , 1995 , 172, 1702-6; discussion 1704-8	6.4	29	
27	Corticosteroid-induced arrest of HELLP syndrome progression in a marginally-viable pregnancy. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 1995 , 59, 217-9	2.4	14	
26	Intrapartum detection of a macrosomic fetus: clinical versus 8 sonographic models. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , 1995 , 35, 266-70	1.7	46	
25	Infectious morbidity, operative blood loss, and length of the operative procedure after cesarean delivery by method of placental removal and site of uterine repair. <i>Journal of the American College of Surgeons</i> , 1995 , 181, 517-20	4.4	41	
24	New-onset hypertension in the pregnant patient. <i>Obstetrics and Gynecology Clinics of North America</i> , 1995 , 22, 157-71	3.3	1	
23	Ultrasonic assessment of the amniotic fluid volume in diamniotic twins. <i>Journal of the Society for Gynecologic Investigation</i> , 1995 , 2, 609-13		14	
22	The laboratory evaluation of hypertensive gravidas. <i>Obstetrical and Gynecological Survey</i> , 1995 , 50, 138-	4 54	22	

21	Accelerated recovery from severe preeclampsia: uterine curettage versus nifedipine. <i>Journal of the Society for Gynecologic Investigation</i> , 1994 , 1, 210-4		21
20	Neonatal salvage by week& gestation in pregnancies complicated by HELLP syndrome. <i>Journal of the Society for Gynecologic Investigation</i> , 1994 , 1, 206-9		20
19	Antepartum corticosteroids: disease stabilization in patients with the syndrome of hemolysis, elevated liver enzymes, and low platelets (HELLP). <i>American Journal of Obstetrics and Gynecology</i> , 1994 , 171, 1148-53	6.4	174
18	Invalid equation to describe the amniotic fluid index in normal pregnancy. <i>American Journal of Obstetrics and Gynecology</i> , 1994 , 170, 1209-10	6.4	5
17	Postpartum corticosteroids: accelerated recovery from the syndrome of hemolysis, elevated liver enzymes, and low platelets (HELLP). <i>American Journal of Obstetrics and Gynecology</i> , 1994 , 171, 1154-8	6.4	121
16	Comparative efficacy of two sonographic measurements for the detection of aberrations in the amniotic fluid volume and the effect of amniotic fluid volume on pregnancy outcome. <i>Obstetrics and Gynecology</i> , 1994 , 83, 959-62	4.9	42
15	Obstetric challenges of massive obesity complicating pregnancy. <i>Journal of Perinatology</i> , 1994 , 14, 10-4	3.1	45
14	Factors relevant to mode of preterm delivery with syndrome of HELLP (hemolysis, elevated liver enzymes, and low platelets). <i>American Journal of Obstetrics and Gynecology</i> , 1994 , 170, 1828-32; discussion 1832-4	6.4	2
13	Acute tocolysis for fetal distress: terbutaline versus magnesium sulphate. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , 1993 , 33, 362-4	1.7	22
12	Corticosteroids for enhanced fetal lung maturation in patients with HELLP syndrome: impact on neonates. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , 1993 , 33, 131-5	1.7	23
11	Standard parameters of preeclampsia: can the clinician depend upon them to reliably identify the patient with the HELLP syndrome?. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , 1993 , 33, 122-6	1.7	17
10	Corticosteroids for the enhancement of fetal lung maturity: impact on the gravida with preeclampsia and the HELLP syndrome. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , 1993 , 33, 127-31	1.7	38
9	Placenta praevia: does uterine activity cause bleeding?. <i>Australian and New Zealand Journal of Obstetrics and Gynaecology</i> , 1993 , 33, 22-4	1.7	7
8	Does method of placental removal or site of uterine incision repair alter endometritis after cesarean delivery?. <i>Infectious Diseases in Obstetrics and Gynecology</i> , 1993 , 1, 65-70	2.4	13
7	The interrelationship of eclampsia, HELLP syndrome, and prematurity: cofactors for significant maternal and perinatal risk. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 1993 , 100, 109	9 3 :7100	42
6	Blood loss at time of cesarean section by method of placental removal and exteriorization versus in situ repair of the uterine incision. <i>Surgery, Gynecology & Obstetrics</i> , 1993 , 177, 389-92		32
5	Immediate postpartum curettage: accelerated recovery from severe preeclampsia. <i>Obstetrics and Gynecology</i> , 1993 , 81, 502-6	4.9	36
4	Preoperative skin preparation and intraoperative pelvic irrigation: impact on post-cesarean endometritis and wound infection. <i>Obstetrics and Gynecology</i> , 1993 , 81, 922-5	4.9	27

LIST OF PUBLICATIONS

3	Measurement of amniotic fluid volume: accuracy of ultrasonography techniques. <i>American Journal of Obstetrics and Gynecology</i> , 1992 , 167, 1533-7	6.4	109
2	The effect of placental management at cesarean delivery on operative blood loss. <i>American Journal of Obstetrics and Gynecology</i> , 1992 , 167, 1363-7	6.4	65
1	Pregnancy outcome in an active-duty population. Obstetrics and Gynecology, 1991, 78, 391-3	4.9	12