
## Weilong Yao

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1213289/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Mcl-1 levels critically impact the sensitivities of human colorectal cancer cells to APG-1252-M1, a novel Bcl-2/Bcl-XL dual inhibitor that induces Bax-dependent apoptosis. Neoplasia, 2022, 29, 100798.	2.3	5
2	Inhibition of MEK5/ERK5 signaling overcomes acquired resistance to the third generation EGFR inhibitor, osimertinib, via enhancing Bim-dependent apoptosis. Cancer Letters, 2021, 519, 141-149.	3.2	8
3	KLF13 suppresses the proliferation and growth of colorectal cancer cells through transcriptionally inhibiting HMGCS1-mediated cholesterol biosynthesis. Cell and Bioscience, 2020, 10, 76.	2.1	31
4	BRD4 Levels Determine the Response of Human Lung Cancer Cells to BET Degraders That Potently Induce Apoptosis through Suppression of Mcl-1. Cancer Research, 2020, 80, 2380-2393.	0.4	28
5	Downregulation of RPS15A by miR-29a-3p attenuates cell proliferation in colorectal carcinoma. Bioscience, Biotechnology and Biochemistry, 2019, 83, 2057-2064.	0.6	15
6	Co-inhibition of BET and proteasome enhances ER stress and Bim-dependent apoptosis with augmented cancer therapeutic efficacy. Cancer Letters, 2018, 435, 44-54.	3.2	23
7	Expression of Death Receptor 4 Is Positively Regulated by MEK/ERK/AP-1 Signaling and Suppressed upon MEK Inhibition. Journal of Biological Chemistry, 2016, 291, 21694-21702.	1.6	22
8	Enhancing therapeutic efficacy of the MEK inhibitor, MEK162, by blocking autophagy or inhibiting PI3K/Akt signaling in human lung cancer cells. Cancer Letters, 2015, 364, 70-78.	3.2	40
9	The BET bromodomain inhibitor, JQ1, facilitates c-FLIP degradation and enhances TRAIL-induced apoptosis independent of BRD4 and c-Myc inhibition. Oncotarget, 2015, 6, 34669-34679.	0.8	35