

Vadim Iablokov

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/12137545/publications.pdf>

Version: 2024-02-01

9
papers

166
citations

1307594

7
h-index

1720034

7
g-index

9
all docs

9
docs citations

9
times ranked

363
citing authors

#	ARTICLE	IF	CITATIONS
1	Proteinase-activated Receptor 2 (PAR2) Decreases Apoptosis in Colonic Epithelial Cells. <i>Journal of Biological Chemistry</i> , 2014, 289, 34366-34377.	3.4	45
2	Naturally Occurring Glycoalkaloids in Potatoes Aggravate Intestinal Inflammation in Two Mouse Models of Inflammatory Bowel Disease. <i>Digestive Diseases and Sciences</i> , 2010, 55, 3078-3085.	2.3	28
3	The serine protease-mediated increase in intestinal epithelial barrier function is dependent on occludin and requires an intact tight junction. <i>American Journal of Physiology - Renal Physiology</i> , 2016, 311, G466-G479.	3.4	26
4	Tumor necrosis factor α decreases aquaporin 3 expression in intestinal epithelial cells through inhibition of constitutive transcription. <i>Physiological Reports</i> , 2017, 5, e13451.	1.7	23
5	Epidermal growth factor receptor transactivation is required for proteinase-activated receptor-2-induced COX-2 expression in intestinal epithelial cells. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 303, G111-G119.	3.4	22
6	Detecting Proteomic Indicators to Distinguish Diabetic Nephropathy from Hypertensive Nephrosclerosis by Integrating Matrix-Assisted Laser Desorption/Ionization Mass Spectrometry Imaging with High-Mass Accuracy Mass Spectrometry. <i>Kidney and Blood Pressure Research</i> , 2020, 45, 233-248.	2.0	12
7	The Canadian MD/PhD training program needs reinstated support. <i>Nature Medicine</i> , 2015, 21, 1111-1111.	30.7	10
8	Protease-activated receptor 2 (PAR2) inactivates the proapoptotic protein, BAD, via ERK1/2 and PI3K activity to decrease apoptosis in colonic epithelial cells. <i>FASEB Journal</i> , 2013, 27, 727.4.	0.5	0
9	The Ability of Serine Proteases to Induce an Increase in Barrier Function is Dependent on the Tight Junction Protein Occludin. <i>FASEB Journal</i> , 2015, 29, 282.2.	0.5	0