

Abdullah AtlÄ±

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/12141971/publications.pdf>

Version: 2024-02-01

20
papers

552
citations

840776

11
h-index

839539

18
g-index

20
all docs

20
docs citations

20
times ranked

912
citing authors

#	ARTICLE	IF	CITATIONS
1	Bipolar bozukluk tip I hastalarÄ±nÄ±n tedavisinde uzun etkili antipsikotiklerin klinik kullanÄ±mÄ±. Cukurova Medical Journal, 2022, 47, 844-851.	0.2	0
2	Case Reports Presentations. Journal of Theoretical Social Psychology, 2018, 28, 114-270.	1.9	34
3	Oxidative metabolism may be associated with negative symptoms in schizophrenia. Journal of Theoretical Social Psychology, 2017, 27, 54-61.	1.9	12
4	The investigation of factors related to suicide attempts in Southeastern Turkey. Neuropsychiatric Disease and Treatment, 2016, 12, 407.	2.2	11
5	Family History in Patients with Bipolar Disorder. Noropsikiyatri Arsivi, 2016, 53, 276-279.	0.7	14
6	Evaluation of Paraoxonase, Arylesterase and Malondialdehyde Levels in Schizophrenia Patients Taking Typical, Atypical and Combined Antipsychotic Treatment. Clinical Psychopharmacology and Neuroscience, 2016, 14, 345-350.	2.0	21
7	Acute Unilateral Myopia Induced by Add-On Aripiprazole: A Case Report. Journal of Microbiology and Biotechnology, 2016, 26, 68-71.	2.1	2
8	Comparison of Clinical Characteristics of Patients on whom Electroconvulsive Therapy was Applied as Inpatient and Outpatient. Journal of Microbiology and Biotechnology, 2016, 26, 32-38.	2.1	0
9	Diagnostic performance of increased prolidase activity in schizophrenia. Neuroscience Letters, 2016, 613, 36-40.	2.1	21
10	Altered lipid peroxidation markers are related to post-traumatic stress disorder (PTSD) and not trauma itself in earthquake survivors. European Archives of Psychiatry and Clinical Neuroscience, 2016, 266, 329-336.	3.2	36
11	Decreased Prolidase Activity in Patients with Posttraumatic Stress Disorder. Psychiatry Investigation, 2016, 13, 420.	1.6	5
12	Neutrophil–lymphocyte ratio in patients with major depressive disorder undergoing no pharmacological therapy. Neuropsychiatric Disease and Treatment, 2015, 11, 2253.	2.2	92
13	Serum levels of carcinoembryonic antigen (CEA) in patients with bipolar disorder. Acta Neuropsychiatrica, 2015, 27, 177-181.	2.1	4
14	Serum total oxidant and antioxidant status in earthquake survivors with post-traumatic stress disorder. Acta Neuropsychiatrica, 2015, 27, 153-158.	2.1	4
15	Relations between mood characteristics, circadian preferences, and functionality in multiple sclerosis. International Journal of Psychiatry in Clinical Practice, 2015, 19, 148-154.	2.4	3
16	Magnetic Resonance Imaging in Children under Anesthesia: The Relationship between the Degree of Information Provided to Parents and Parentsâ€™ Anxiety Scores. BioMed Research International, 2014, 1-5.	1.9	8
17	Latent class analysis of obsessiveâ€™compulsive symptoms in a clinical sample. Comprehensive Psychiatry, 2014, 55, 604-612.	3.1	10
18	Mood and metabolic consequences of sleep deprivation as a potential endophenotypeâ€™ in bipolar disorder. Journal of Affective Disorders, 2013, 150, 284-294.	4.1	31

#	ARTICLE	IF	CITATIONS
19	Chronotype Differences in Suicidal Behavior and Impulsivity Among Suicide Attempters. Chronobiology International, 2011, 28, 170-175.	2.0	90
20	ASSOCIATIONS BETWEEN CHRONOTYPE, SLEEP QUALITY, SUICIDALITY, AND DEPRESSIVE SYMPTOMS IN PATIENTS WITH MAJOR DEPRESSION AND HEALTHY CONTROLS. Chronobiology International, 2010, 27, 1813-1828.	2.0	154