Cindy L K Lam

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1214636/publications.pdf

Version: 2024-02-01

147801 4,570 174 31 citations h-index papers

177 5906 citing authors

144013

57

g-index

177 all docs

177 docs citations

times ranked

#	Article	IF	CITATIONS
1	Is the standard SF-12 Health Survey valid and equivalent for a Chinese population?. Quality of Life Research, 2005, 14, 539-547.	3.1	451
2	Prevalence, clinical spectrum and health care utilization of gastroâ€oesophageal reflux disease in a Chinese population: a populationâ€based study. Alimentary Pharmacology and Therapeutics, 2003, 18, 595-604.	3.7	238
3	The SF-36 summary scales were valid, reliable, and equivalent in a Chinese population. Journal of Clinical Epidemiology, 2005, 58, 815-822.	5.0	219
4	Anxiety but not depression determines health careâ€seeking behaviour in Chinese patients with dyspepsia and irritable bowel syndrome: a populationâ€based study. Alimentary Pharmacology and Therapeutics, 2002, 16, 2081-2088.	3.7	171
5	Is the SFâ€12 version 2 Health Survey a valid and equivalent substitute for the SFâ€36 version 2 Health Survey for the Chinese?. Journal of Evaluation in Clinical Practice, 2013, 19, 200-208.	1.8	150
6	Valuation of the SF-6D Health States Is Feasible, Acceptable, Reliable, and Valid in a Chinese Population. Value in Health, $2008,11,295-303.$	0.3	111
7	Population based study of noncardiac chest pain in southern Chinese: Prevalence, psychosocial factors and health care utilization. World Journal of Gastroenterology, 2004, 10, 707.	3.3	86
8	A validated symptoms questionnaire (Chinese GERDQ) for the diagnosis of gastroâ€oesophageal reflux disease in the Chinese population. Alimentary Pharmacology and Therapeutics, 2003, 17, 1407-1413.	3.7	85
9	Five-Year Effectiveness of the Multidisciplinary Risk Assessment and Management Programme–Diabetes Mellitus (RAMP-DM) on Diabetes-Related Complications and Health Service Uses—A Population-Based and Propensity-Matched Cohort Study. Diabetes Care, 2018, 41, 49-59.	8.6	82
10	The effect of health-related quality of life (HRQOL) on health service utilisation of a Chinese population. Social Science and Medicine, 2002, 55, 1635-1646.	3.8	69
11	Cost effectiveness of mammography screening for Chinese women. Cancer, 2007, 110, 885-895.	4.1	65
12	Effects of Patient Empowerment Programme (PEP) on Clinical Outcomes and Health Service Utilization in Type 2 Diabetes Mellitus in Primary Care: An Observational Matched Cohort Study. PLoS ONE, 2014, 9, e95328.	2.5	65
13	Psychometric validation of the EuroQoL 5-Dimension 5-Level (EQ-5D-5L) in Chinese patients with adolescent idiopathic scoliosis. Scoliosis and Spinal Disorders, 2016, 11, 19.	2.3	64
14	A systematic review of quality of thyroid-specific health-related quality-of-life instruments recommends ThyPRO for patients with benign thyroid diseases. Journal of Clinical Epidemiology, 2016, 78, 63-72.	5.0	64
15	A short message service (SMS) intervention to prevent diabetes in Chinese professional drivers with pre-diabetes: A pilot single-blinded randomized controlled trial. Diabetes Research and Clinical Practice, 2013, 102, 158-166.	2.8	59
16	Impact of Untreated Psychosis on Quality of Life in Patients with First-episode Schizophrenia. Quality of Life Research, 2005, 14, 1803-1811.	3.1	54
17	A pilot study on the validity and reliability of the Patient Enablement Instrument (PEI) in a Chinese population. Family Practice, 2010, 27, 395-403.	1.9	54
18	Systematic screening for diabetic retinopathy (DR) in Hong Kong: prevalence of DR and visual impairment among diabetic population. British Journal of Ophthalmology, 2016, 100, 151-155.	3.9	53

#	Article	IF	Citations
19	The Hong Kong index of dyspepsia: A validated symptom severity questionnaire for patients with dyspepsia. Journal of Gastroenterology and Hepatology (Australia), 2002, 17, 545-551.	2.8	48
20	Evaluation of the internal and external responsiveness of the Pressure Ulcer Scale for Healing (<scp>PUSH</scp>) tool for assessing acute and chronic wounds. Journal of Advanced Nursing, 2016, 72, 1134-1143.	3.3	47
21	Association of variability in hemoglobin A1c with cardiovascular diseases and mortality in Chinese patients with type 2 diabetes mellitus — A retrospective population-based cohort study. Journal of Diabetes and Its Complications, 2016, 30, 1240-1247.	2.3	46
22	Association of Visit-to-Visit Variability of Systolic Blood Pressure With Cardiovascular Disease and Mortality in Primary Care Chinese Patients With Type 2 Diabetes—A Retrospective Population-Based Cohort Study. Diabetes Care, 2017, 40, 270-279.	8.6	42
23	Five-Year Cost-effectiveness of the Multidisciplinary Risk Assessment and Management Programme–Diabetes Mellitus (RAMP-DM). Diabetes Care, 2018, 41, 250-257.	8.6	42
24	Help-seeking intentions and subsequent 12-month mental health service use in Chinese primary care patients with depressive symptoms. BMJ Open, 2015, 5, e006730-e006730.	1.9	41
25	Long-term effects of the multidisciplinary risk assessment and management program for patients with diabetes mellitus (RAMP-DM): a population-based cohort study. Cardiovascular Diabetology, 2015, 14, 105.	6.8	38
26	Effect of Multifactorial Treatment Targets and Relative Importance of Hemoglobin A1c, Blood Pressure, and Lowâ€Density Lipoproteinâ€Cholesterol on Cardiovascular Diseases in Chinese Primary Care Patients With Type 2 Diabetes Mellitus: A Populationâ€Based Retrospective Cohort Study. Journal of the American Heart Association, 2017, 6, .	3.7	37
27	Positive association between gastroâ€oesophageal reflux disease and irritable bowel syndrome in a Chinese population. Alimentary Pharmacology and Therapeutics, 2007, 25, 1099-1104.	3.7	35
28	Systematic review recommends the European Organization for Research and Treatment of Cancer colorectal cancer–specific module for measuring quality of life in colorectal cancer patients. Journal of Clinical Epidemiology, 2015, 68, 266-278.	5.0	34
29	Quality-adjusted life years: population-specific measurement of the quality component. Hong Kong Medical Journal, 2011, 17 Suppl 6, 17-21.	0.1	34
30	Bridging the Gaps Between Patients and Primary Care in China: A Nationwide Representative Survey. Annals of Family Medicine, 2017, 15, 237-245.	1.9	33
31	Prediction of new onset of end stage renal disease in Chinese patients with type 2 diabetes mellitus – a population-based retrospective cohort study. BMC Nephrology, 2017, 18, 257.	1.8	33
32	Onset and disappearance of reflux symptoms in a Chinese population: a 1-year follow-up study. Alimentary Pharmacology and Therapeutics, 2004, 20, 803-812.	3.7	32
33	Direct medical costs of care for Chinese patients with colorectal neoplasia: a health care service provider perspective. Journal of Evaluation in Clinical Practice, 2012, 18, 1203-1210.	1.8	31
34	Measuring body constitution: Validation of the Body Constitution Questionnaire (BCQ) in Hong Kong. Complementary Therapies in Medicine, 2014, 22, 670-682.	2.7	31
35	A Chinese version of the Sleep Apnea Quality of Life Index was evaluated for reliability, validity, and responsiveness. Journal of Clinical Epidemiology, 2004, 57, 470-478.	5.0	30
36	SF-6D population norms for the Hong Kong Chinese general population. Quality of Life Research, 2018, 27, 2349-2359.	3.1	29

#	Article	IF	Citations
37	Systematic review on the cost-effectiveness of self-management education programme for type 2 diabetes mellitus. Diabetes Research and Clinical Practice, 2017, 127, 21-34.	2.8	28
38	The Health-Related Quality of Life of Chinese Patients on Hemodialysis and Peritoneal Dialysis. Patient, 2017, 10, 799-808.	2.7	28
39	Direct and indirect costs of end-stage renal disease patients in the first and second years after initiation of nocturnal home haemodialysis, hospital haemodialysis and peritoneal dialysis. Nephrology Dialysis Transplantation, 2019, 34, 1565-1576.	0.7	28
40	An instrument to assess mental patients? capacity to appraise and report subjective quality of life. Quality of Life Research, 2005, 14, 687-694.	3.1	27
41	Traditional growing rod versus magnetically controlled growing rod for treatment of early onset scoliosis: Cost analysis from implantation till skeletal maturity. Journal of Orthopaedic Surgery, 2017, 25, 230949901770502.	1.0	27
42	Association of estimated glomerular filtration rate and urine albumin-to-creatinine ratio with incidence of cardiovascular diseases and mortality in chinese patients with type 2 diabetes mellitus – a population-based retrospective cohort study. BMC Nephrology, 2017, 18, 47.	1.8	27
43	Mapping the SRS-22r questionnaire onto the EQ-5D-5L utility score in patients with adolescent idiopathic scoliosis. PLoS ONE, 2017, 12, e0175847.	2.5	27
44	Association of Blood Pressure and Risk of Cardiovascular and Chronic Kidney Disease in Hong Kong Hypertensive Patients. Hypertension, 2019, 74, 331-340.	2.7	27
45	The mediating role of sleep quality in the association between nocturia and health-related quality of life. Health and Quality of Life Outcomes, 2019, 17, 181.	2.4	27
46	Greater variability in lipid measurements associated with cardiovascular disease and mortality: A 10â€year diabetes cohort study. Diabetes, Obesity and Metabolism, 2020, 22, 1777-1788.	4.4	27
47	Validation of the Disease-Specific Components of the Kidney Disease Quality of Life-36 (KDQOL-36) in Chinese Patients Undergoing Maintenance Dialysis. PLoS ONE, 2016, 11, e0155188.	2.5	27
48	Development of a cardiovascular diseases risk prediction model and tools for <scp>C</scp> hinese patients with type 2 diabetes mellitus: <scp>A</scp> populationâ€based retrospective cohort study. Diabetes, Obesity and Metabolism, 2018, 20, 309-318.	4.4	26
49	Cost-effectiveness simulation and analysis of colorectal cancer screening in Hong Kong Chinese population: comparison amongst colonoscopy, guaiac and immunologic fecal occult blood testing. BMC Cancer, 2015, 15, 705.	2.6	25
50	Cost-Effectiveness of a Short Message Service Intervention to Prevent Type 2 Diabetes from Impaired Glucose Tolerance. Journal of Diabetes Research, 2016, 2016, 1-8.	2.3	25
51	Mental Health Mediating the Relationship Between Symptom Severity and Healthâ€Related Quality of Life in Patients with Lower Urinary Tract Symptoms. LUTS: Lower Urinary Tract Symptoms, 2016, 8, 141-149.	1.3	25
52	Poverty and health-related quality of life of people living in Hong Kong: comparison of individuals from low-income families and the general population. Journal of Public Health, 2017, 39, fdw046.	1.8	25
53	Effect of Achieved Systolic Blood Pressure on Cardiovascular Outcomes in Patients With Type 2 Diabetes: A Population-Based Retrospective Cohort Study. Diabetes Care, 2018, 41, 1134-1141.	8.6	25
54	PBL curriculum improves medical students' participation in small-group tutorials. Medical Teacher, 2007, 29, e198-e203.	1.8	24

#	Article	IF	Citations
55	Effectiveness of the multidisciplinary Risk Assessment and Management Program for Patients with Diabetes Mellitus (RAMP-DM) for diabetic microvascular complications: A population-based cohort study. Diabetes and Metabolism, 2016, 42, 424-432.	2.9	24
56	The Impact of Cardiovascular Disease and Chronic Kidney Disease on Life Expectancy and Direct Medical Cost in a 10-Year Diabetes Cohort Study. Diabetes Care, 2020, 43, 1750-1758.	8.6	24
57	Patterns of health-related quality of life and associated factors in Chinese patients undergoing haemodialysis. Health and Quality of Life Outcomes, 2015, 13, 108.	2.4	23
58	Effect of a structured diabetes education programme in primary care on hospitalizations and emergency department visits among people with Type 2 diabetes mellitus: results from the Patient Empowerment Programme. Diabetic Medicine, 2016, 33, 1427-1436.	2.3	23
59	Health-related quality of life and health preference of Chinese patients with diabetes mellitus managed in primary care and secondary care setting: decrements associated with individual complication and number of complications. Health and Quality of Life Outcomes, 2017, 15, 125.	2.4	23
60	Awareness of diabetic retinopathy and its association with attendance for systematic screening at the public primary care setting: a cross-sectional study in Hong Kong. BMJ Open, 2018, 8, e019989.	1.9	23
61	Patient Empowerment Programme (PEP) and Risk of Microvascular Diseases Among Patients With Type 2 Diabetes in Primary Care: A Population-Based Propensity-Matched Cohort Study. Diabetes Care, 2015, 38, e116-e117.	8.6	22
62	Increased number of structured diabetes education attendance was not associated with the improvement in patient-reported health-related quality of life: results from Patient Empowerment Programme (PEP). Health and Quality of Life Outcomes, 2015, 13, 126.	2.4	22
63	Annual direct medical costs associated with diabetesâ€related complications in the event year and in subsequent years in Hong Kong. Diabetic Medicine, 2017, 34, 1276-1283.	2.3	22
64	Generative adversarial networks for imputing missing data for big data clinical research. BMC Medical Research Methodology, 2021, 21, 78.	3.1	22
65	Macrovascular and microvascular disease in obese patients with type 2 diabetes attending structured diabetes education program: a population-based propensity-matched cohort analysis of Patient Empowerment Programme (PEP). Endocrine, 2016, 53, 412-422.	2.3	21
66	How Does the Family Influence Adolescent Eating Habits in Terms of Knowledge, Attitudes and Practices? A Global Systematic Review of Qualitative Studies. Nutrients, 2021, 13, 3717.	4.1	21
67	Five-year change of clinical and complications profile of diabetic patients under primary care: a population-based longitudinal study on 127,977 diabetic patients. Diabetology and Metabolic Syndrome, 2015, 7, 79.	2.7	20
68	Development of a Health Empowerment Programme to improve the health of working poor families: protocol for a prospective cohort study in Hong Kong. BMJ Open, 2016, 6, e010015.	1.9	20
69	Development and pilot evaluation of a mobile app on parent-child exercises to improve physical activity and psychosocial outcomes of Hong Kong Chinese children. BMC Public Health, 2020, 20, 1544.	2.9	20
70	Outcomes of Graves' Disease Patients Following Antithyroid Drugs, Radioactive Iodine, or Thyroidectomy as the First-line Treatment. Annals of Surgery, 2021, 273, 1197-1206.	4.2	20
71	Impact of sleep duration, physical activity, and screen time on health-related quality of life in children and adolescents. Health and Quality of Life Outcomes, 2021, 19, 145.	2.4	20
72	Main predictors in health-related quality of life in Chinese patients with type 2 diabetes mellitus. Quality of Life Research, 2016, 25, 2957-2965.	3.1	19

#	Article	IF	CITATIONS
73	The 12-Month Incidence and Predictors of PHQ-9-Screened Depressive Symptoms in Chinese Primary Care Patients. Annals of Family Medicine, 2016, 14, 47-53.	1.9	19
74	Prediction of five-year all-cause mortality in Chinese patients with type 2 diabetes mellitus $\hat{a} \in A$ population-based retrospective cohort study. Journal of Diabetes and Its Complications, 2017, 31, 939-944.	2.3	19
75	Double-blind, randomized controlled study to assess the effects of lansoprazole 30â€∫mg and lansoprazole 15â€∫mg on 24-h oesophageal and intragastric pH in Chinese subjects with gastro-oesophageal reflux disease. Alimentary Pharmacology and Therapeutics, 2004, 19, 455-462.	3.7	18
76	Psychometric Validation of the Traditional Chinese Version of the Early Onset Scoliosis-24 Item Questionnaire (EOSQ-24). Spine, 2016, 41, E1460-E1469.	2.0	18
77	Cost-effectiveness analysis of vaccinations and decision makings on vaccination programmes in Hong Kong: A systematic review. Vaccine, 2017, 35, 3153-3161.	3.8	18
78	The 21st Century: The Age of Family Medicine Research?. Annals of Family Medicine, 2004, 2, S50-S54.	1.9	17
79	Brief problemâ€solving treatment in primary care (pstâ€pc) was not more effective than placebo for elderly patients screened positive of psychological problems. International Journal of Geriatric Psychiatry, 2010, 25, 968-980.	2.7	17
80	Evaluation of the Effectiveness of Nurse-Led Continence Care Treatments for Chinese Primary Care Patients with Lower Urinary Tract Symptoms. PLoS ONE, 2015, 10, e0129875.	2.5	17
81	Incidence and predictors for cardiovascular disease in Chinese patients with type 2 diabetes mellitus – a population-based retrospective cohort study. Journal of Diabetes and Its Complications, 2016, 30, 444-450.	2.3	17
82	Primary care workforce and continuous medical education in China: lessons to learn from a nationwide cross-sectional survey. BMJ Open, 2017, 7, e015145.	1.9	17
83	Comparative Risks of Nonsteroidal Anti-Inflammatory Drugs on CKD. Clinical Journal of the American Society of Nephrology: CJASN, 2021, 16, 898-907.	4.5	17
84	Responsiveness was similar between direct and mapped SF-6D inÂcolorectal cancer patients who declined. Journal of Clinical Epidemiology, 2014, 67, 219-227.	5.0	16
85	Simple nonâ€laboratory―and laboratoryâ€based risk assessment algorithms and nomogram for detecting undiagnosed diabetes mellitus. Journal of Diabetes, 2016, 8, 414-421.	1.8	16
86	Mental health of Chinese primary care patients with lower urinary tract symptoms. Psychology, Health and Medicine, 2016, 21, 113-127.	2.4	16
87	Morbidity pattern of traditional Chinese medicine primary care in the Hong Kong population. Scientific Reports, 2017, 7, 7513.	3.3	15
88	Spillover Effects of Maternal Chronic Disease on Children's Quality of Life and Behaviors Among Low-Income Families. Patient, 2018, 11, 625-635.	2.7	15
89	Healthâ€related quality of life and health utility of Chinese patients undergoing nocturnal home haemodialysis in comparison with other modes of dialysis. Nephrology, 2019, 24, 630-637.	1.6	15
90	Ageâ€specific associations of <scp>glycated haemoglobin</scp> variability with cardiovascular disease and mortality in patients with type 2 diabetes mellitus: A 10―year cohort study. Diabetes, Obesity and Metabolism, 2020, 22, 1316-1327.	4.4	15

#	Article	IF	CITATIONS
91	Successful Pulmonary Vein Isolation Using Transvenous Catheter Cryoablation Improves Quality-of-Life in Patients with Atrial Fibrillation. PACE - Pacing and Clinical Electrophysiology, 2005, 28, 421-424.	1.2	14
92	Unrecognised psychological problems impair quality of life and increase consultation rates in Chinese elderly patients. International Journal of Geriatric Psychiatry, 2009, 24, 979-989.	2.7	14
93	Can HbA1c replace OGTT for the diagnosis of diabetes mellitus among Chinese patients with impaired fasting glucose?. Family Practice, 2015, 32, cmv077.	1.9	14
94	Maternal mental quality of life mediates the associations between intimate partner abuse against mothers and their children's behaviours and quality of life in low-income Chinese families. Quality of Life Research, 2017, 26, 3297-3306.	3.1	14
95	Direct medical costs of diabetes mellitus in the year of mortality and year preceding the year of mortality. Diabetes, Obesity and Metabolism, 2018, 20, 1470-1478.	4.4	14
96	Fiveâ€year effectiveness of bariatric surgery on disease remission, weight loss, and changes of metabolic parameters in obese patients with type 2 diabetes: A populationâ€based propensity scoreâ€matched cohort study. Diabetes/Metabolism Research and Reviews, 2020, 36, e3236.	4.0	14
97	Utilisation patterns of primary health care services in Hong Kong: does having a family doctor make any difference?. Hong Kong Medical Journal, 2011, 17, 28-32.	0.1	14
98	Bariatric surgery is expensive but improves co-morbidity: 5-year assessment of patients with obesity and type 2 diabetes. British Journal of Surgery, 2021, 108, 554-565.	0.3	14
99	Health-related quality of life mediates associations between multi-morbidity and depressive symptoms in Chinese primary care patients. Family Practice, 2016, 33, 61-68.	1.9	13
100	Longâ€term costâ€effectiveness of a Patient Empowerment Programme for type 2 diabetes mellitus in primary care. Diabetes, Obesity and Metabolism, 2019, 21, 73-83.	4.4	13
101	Association of visit-to-visit variability of systolic blood pressure with cardiovascular disease, chronic kidney disease and mortality in patients with hypertension. Journal of Hypertension, 2020, 38, 943-953.	0.5	13
102	Lifetime cost-effectiveness analysis of first-line dialysis modalities for patients with end-stage renal disease under peritoneal dialysis first policy. BMC Nephrology, 2020, 21, 42.	1.8	13
103	Do We Need a Patient-Centered Target for Systolic Blood Pressure in Hypertensive Patients With Type 2 Diabetes Mellitus?. Hypertension, 2017, 70, 1273-1282.	2.7	12
104	Statin use reduces cardiovascular events and all-cause mortality amongst Chinese patients with type 2 diabetes mellitus: a 5-year cohort study. BMC Cardiovascular Disorders, 2017, 17, 166.	1.7	12
105	A prospective cohort study to investigate parental stress and child health in low-income Chinese families: protocol paper. BMJ Open, 2018, 8, e018792.	1.9	12
106	Effects of bariatric surgery on kidney diseases, cardiovascular diseases, mortality and severe hypoglycaemia among patients with Type 2 diabetes mellitus. Nephrology Dialysis Transplantation, 2021, 36, 1440-1451.	0.7	12
107	The Spill-Over Impact of the Novel Coronavirus-19 Pandemic on Medical Care and Disease Outcomes in Non-communicable Diseases: A Narrative Review. Public Health Reviews, 2022, 43, 1604121.	3.2	12
108	The responsiveness of the International Prostate Symptom Score, Incontinence Impact Questionnaireâ€7 and Depression, Anxiety and Stress Scaleâ€21 in patients with lower urinary tract symptoms. Journal of Advanced Nursing, 2015, 71, 1857-1870.	3.3	11

#	Article	IF	Citations
109	Fiveâ€year costâ€effectiveness of the Patient Empowerment Programme (<scp>PEP</scp>) for type 2 diabetes mellitus in primary care. Diabetes, Obesity and Metabolism, 2017, 19, 1312-1316.	4.4	11
110	Nonrestorative sleep scale: reliable and valid for the Chinese population. Quality of Life Research, 2019, 28, 1685-1692.	3.1	11
111	Prediction models and nomograms for 10â€year risk of endâ€stage renal disease in Chinese type 2 diabetes mellitus patients in primary care. Diabetes, Obesity and Metabolism, 2021, 23, 897-909.	4.4	11
112	Having a Family Doctor is Associated with Some Better Patient-Reported Outcomes of Primary Care Consultations. Frontiers in Medicine, 2014, 1, 29.	2.6	10
113	A Mobile App for Identifying Individuals With Undiagnosed Diabetes and Prediabetes and for Promoting Behavior Change: 2-Year Prospective Study. JMIR MHealth and UHealth, 2018, 6, e10662.	3.7	10
114	Awareness of osteoporosis among physicians in China. Osteoporosis International, 2004, 15, 329-334.	3.1	9
115	Evaluation of patient-reported outcomes data in structured diabetes education intervention: 2-year follow-up data of patient empowerment programme. Endocrine, 2016, 54, 422-432.	2.3	9
116	A curvilinear association of body mass index with cardiovascular diseases in Chinese patients with type 2 diabetes mellitus $\hat{a} \in A$ population-based retrospective cohort study. Journal of Diabetes and Its Complications, 2016, 30, 1261-1268.	2.3	9
117	The impact of a co-payment on the cost-effectiveness of screening for diabetic retinopathy. Journal of Public Health, 2016, 38, 782-792.	1.8	9
118	EQ-5D-5L and SF-6D Utility Measures in Symptomatic benign Thyroid Nodules: Acceptability and Psychometric Evaluation. Patient, 2017, 10, 447-454.	2.7	9
119	Effectiveness of a multidisciplinary risk assessment and management programme—diabetes mellitus (RAMP-DM) on patient-reported outcomes. Endocrine, 2017, 55, 416-426.	2.3	9
120	Cultural adaptation and psychometric properties of the Chinese Burden of Treatment Questionnaire (C-TBQ) in primary care patients with multi-morbidity. Family Practice, 2019, 36, 657-665.	1.9	9
121	Intensification with dipeptidyl peptidase-4 inhibitor, insulin, or thiazolidinediones and risks of all-cause mortality, cardiovascular diseases, and severe hypoglycemia in patients on metformin-sulfonylurea dual therapy: A retrospective cohort study. PLoS Medicine, 2019, 16, e1002999.	8.4	9
122	The test–retest reliability of the <scp>I</scp> ncontinence <scp>Q</scp> uestionnaireâ€ <scp>U</scp> rinary <scp>I</scp> ncontinence <scp>S</scp> hort <scp>F</scp> orm (<scp>ICIQ</scp> â€ <scp>UI SF</scp>) for assessing type of urinary incontinence in males and females. Journal of Clinical Nursing, 2015, 24, 3742-3744.	3.0	8
123	Evaluation of the quality of care of a haemodialysis public-private partnership programme for patients with end-stage renal disease. BMC Nephrology, 2016, 17, 79.	1.8	8
124	Evaluation of the internal and external responsiveness of Short Form-12 Health Survey version 2 (SF-12v2) in patients with type 2 diabetes mellitus. Quality of Life Research, 2018, 27, 2459-2469.	3.1	8
125	Evaluation of the responsiveness of Short Form-12 Health Survey version 2 (SF-12v2) in Chinese patients with hypertension in primary care. Quality of Life Research, 2019, 28, 2851-2857.	3.1	8
126	Greater variability in lipid measurements associated with kidney diseases in patients with type 2 diabetes mellitus in a 10-year diabetes cohort study. Scientific Reports, 2021, 11, 8047.	3.3	8

#	Article	IF	CITATIONS
127	Evaluation of the quality of care of a multi-disciplinary Risk Factor Assessment and Management Programme for Hypertension (RAMP-HT). BMC Family Practice, 2015, 16, 71.	2.9	7
128	The mediating factors in the relationship between lower urinary tract symptoms and health-related quality of life. BMC Research Notes, 2017, 10, 611.	1.4	7
129	Ten-year risk prediction models of complications and mortality of Chinese patients with diabetes mellitus in primary care in Hong Kong: a study protocol. BMJ Open, 2018, 8, e023070.	1.9	7
130	SGLT2i as fourth-line therapy and risk of mortality, end-stage renal diseases and cardiovascular diseases in patients with type 2 diabetes mellitus. Diabetes and Metabolism, 2020, 47, 101196.	2.9	7
131	Nonrestorative sleep scale: a reliable and valid short form of the traditional Chinese version. Quality of Life Research, 2020, 29, 2585-2592.	3.1	7
132	Adolescent Knowledge, Attitudes and Practices of Healthy Eating: Findings of Qualitative Interviews among Hong Kong Families. Nutrients, 2022, 14, 2857.	4.1	7
133	Road to better health and integration: a Delphi study on health service models for Hong Kong migrants. International Journal for Equity in Health, 2014, 13, 127.	3.5	6
134	The effect of timing of incentive payments on response rates for cohort study telephone interviews in primary care setting with cost-minimization analysis, a randomized controlled trial. BMC Medical Research Methodology, 2015, 15, 79.	3.1	6
135	The role of the family doctor in the era of multi-disciplinary primary care. Family Practice, 2016, 33, 447-448.	1.9	6
136	Evaluation of the outcomes of care of nurseâ€led continence care clinics for Chinese patients with lower urinary tract symptoms, a 2â€year prospective longitudinal study. Journal of Advanced Nursing, 2017, 73, 1158-1171.	3.3	6
137	Temporal trends in quality of primary care for patients with type 2 diabetes mellitus: A populationâ€based retrospective cohort study after implementation of a quality improvement initiative. Diabetes/Metabolism Research and Reviews, 2018, 34, e2952.	4.0	6
138	Direct medical costs in the preceding, event and subsequent years of a first severe hypoglycaemia episode requiring hospitalization: A populationâ€based cohort study. Diabetes, Obesity and Metabolism, 2019, 21, 1330-1339.	4.4	6
139	Do Negative Emotional States Play a Role in the Association Between Intimate Partner Violence and Poor Health-Related Quality of Life in Chinese Women From Low-Income Families?. Violence Against Women, 2020, 26, 2041-2061.	1.7	6
140	Improvement in patient-reported outcomes in Chinese adults after bariatric surgery: 1-year follow-up of a prospective cohort. Surgery for Obesity and Related Diseases, 2020, 16, 1563-1572.	1,2	6
141	Author response to: Bariatric surgery is expensive but improves co-morbidity: 5-year assessment of patients with obesity and type 2 diabetes. British Journal of Surgery, 2021, 108, e281-e281.	0.3	6
142	Long-term outcome of patients treated with antithyroid drugs, radioactive iodine or surgery for persistent or relapsed Graves' disease. British Journal of Surgery, 2022, 109, 381-389.	0.3	6
143	Possible Impact of Incremental Cost-Effectiveness Ratio (ICER) on Decision Making for Cancer Screening in Hong Kong: A Systematic Review. Applied Health Economics and Health Policy, 2016, 14, 647-657.	2.1	5
144	Quality of care and volume for patients with diabetes mellitus in the primary care setting: A population based retrospective cohort study. Diabetes Research and Clinical Practice, 2016, 120, 171-181.	2.8	5

#	Article	IF	CITATIONS
145	Relation between HbA1c and incident cardiovascular disease over a period of 6 years in the Hong Kong population. Diabetes and Metabolism, 2018, 44, 415-423.	2.9	5
146	Five-year effectiveness of short messaging service (SMS) for pre-diabetes. BMC Research Notes, 2018, 11, 709.	1.4	5
147	Cost-effectiveness of a primary care multidisciplinary Risk Assessment and Management Program for patients with diabetes mellitus (RAMP-DM) over lifetime. Endocrine, 2019, 63, 259-269.	2.3	5
148	Cultural adaptation and content validity of a Chinese translation of the †Person-Centered Primary Care Measure': findings from cognitive debriefing. Family Medicine and Community Health, 2020, 8, e000621.	1.6	5
149	Healthcare utilization and direct medical cost in the years during and after cancer diagnosis in patients with typeÂ2 diabetes mellitus. Journal of Diabetes Investigation, 2020, 11, 1661-1672.	2.4	5
150	Association between bariatric surgery and risks of cancer among Chinese patients with type 2 diabetes mellitus: A retrospective cohort study. Journal of Diabetes, 2021, 13, 868-881.	1.8	5
151	A pilot study on the validity and psychometric properties of the electronic EQ-5D-5L in routine clinical practice. Health and Quality of Life Outcomes, 2021, 19, 266.	2.4	5
152	Trends in Asthma Therapy in Hong Kong, 1987–1992. Journal of Asthma, 1993, 30, 475-483.	1.7	4
153	Classification Rule for 5-year Cardiovascular Diseases Risk using decision tree in Primary Care Chinese Patients with Type 2 Diabetes Mellitus. Scientific Reports, 2017, 7, 15238.	3.3	4
154	The validity, reliability, sensitivity and responsiveness of a modified Patient Enablement Instrument (PEI-2) as a tool for serial measurements of health enablement. Family Practice, 2020, 38, 339-345.	1.9	4
155	<p>The APEC Digital Hub-WONCA Collaborative Framework on Integration of Mental Health into Primary Care in the Asia Pacific</p> . Journal of Multidisciplinary Healthcare, 2020, Volume 13, 1693-1704.	2.7	4
156	Confucius School of Medicine: the way to great learning. BMJ: British Medical Journal, 2001, 323, 1458-1459.	2.3	3
157	Effects of health-related quality of life on health service utilisation in patients with colorectal neoplasms. European Journal of Cancer Care, 2018, 27, e12926.	1.5	3
158	Prediction models for the risk of cardiovascular diseases in Chinese patients with type 2 diabetes mellitus: a systematic review. Public Health, 2020, 186, 144-156.	2.9	3
159	Psychometric evaluation of the Chinese version of the Toronto Hospital Alertness Test. Journal of Patient-Reported Outcomes, 2020, 4, 32.	1.9	3
160	DPP4i, thiazolidinediones, or insulin and risks of cancer in patients with type 2 diabetes mellitus on metformin–sulfonylurea dual therapy with inadequate control. BMJ Open Diabetes Research and Care, 2020, 8, e001346.	2.8	3
161	Trend in health-related quality of life and health utility and their decrements due to non-communicable diseases and risk factors: analysis of four population-based surveys between 1998 and 2015. Quality of Life Research, 2020, 29, 2921-2934.	3.1	3
162	Retrospective cohort study to investigate the 10-year trajectories of disease patterns in patients with hypertension and/or diabetes mellitus on subsequent cardiovascular outcomes and health service utilisation: a study protocol. BMJ Open, 2021, 11, e038775.	1.9	3

#	Article	IF	CITATIONS
163	Association Between Team-Based Continuity of Care and Risk of Cardiovascular Diseases Among Patients With Diabetes: A Retrospective Cohort Study. Diabetes Care, 2022, 45, 1162-1169.	8.6	3
164	Attaining biochemical euthyroidism early after total thyroidectomy in Graves' disease may lower long-term morbidity risk. BJS Open, 2022, 6, .	1.7	3
165	Lower urinary tract symptoms and health-related quality of life in Hong Kong primary care: a cross-sectional study. Quality of Life Research, 2020, 29, 1311-1321.	3.1	2
166	Association between health behaviours and cardiometabolic dysregulation: a population-based survey among healthy adults in Hong Kong. BMJ Open, 2021, 11, e043503.	1.9	2
167	Unintentional Injury Burden in Hong Kong: Results from a Representative Population-Based Survey. International Journal of Environmental Research and Public Health, 2021, 18, 8826.	2.6	2
168	The association between trajectories of risk factors and risk of cardiovascular disease or mortality among patients with diabetes or hypertension: A systematic review. PLoS ONE, 2022, 17, e0262885.	2.5	2
169	Tree analysis modeling of the associations between PHQ-9 depressive symptoms and doctor diagnosis of depression in primary care. Psychological Medicine, 2019, 49, 449-457.	4.5	1
170	Chinese herbal medicine in the treatment of acute upper respiratory tract infection: a randomised, double blind, placebo-controlled clinical trial. Hong Kong Medical Journal, 2009, 15 Suppl 6, 30-4.	0.1	1
171	Morbidity Patterns in Primary Care in Hong Kong: Protocol for a Practice-Based Morbidity Survey. JMIR Research Protocols, 2022, 11, e37334.	1.0	1
172	Health-related quality of life in patients with colorectal neoplasm and cost-effectiveness of colorectal cancer screening in Hong Kong. Hong Kong Medical Journal, 2015, 21 Suppl 6, 4-8.	0.1	0
173	Identifying and articulating the student experience in the Intercalated Enrichment Year. BMC Medical Education, 2022, 22, 246.	2.4	0
174	Association of genetic variants related to combined exposure to higher BMI and waist-to-hip ratio on lifelong cardiovascular risk in UK Biobank. Public Health Nutrition, 2023, 26, 416-424.	2.2	0