

Giovanni Triolo

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/12155845/giovanni-triolo-publications-by-year.pdf>

Version: 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

79
papers

3,326
citations

34
h-index

56
g-index

79
ext. papers

3,933
ext. citations

4.9
avg, IF

4.73
L-index

#	Paper	IF	Citations
79	The growing role of precision medicine for the treatment of autoimmune diseases; results of a systematic review of literature and Experts' Consensus. <i>Autoimmunity Reviews</i> , 2021 , 20, 102738	13.6	14
78	A phase 2 randomized, double-blind, placebo-controlled, proof-of-concept study of oral seletalisib in primary Sjögren's syndrome. <i>Rheumatology</i> , 2021 , 60, 1364-1375	3.9	10
77	Macrophage Activation Syndrome in Patients Affected by Adult-onset Still Disease: Analysis of Survival Rates and Predictive Factors in the Gruppo Italiano di Ricerca in Reumatologia Clinica e Sperimentale Cohort. <i>Journal of Rheumatology</i> , 2018 , 45, 864-872	4.1	50
76	Interleukin-25 Axis Is Involved in the Pathogenesis of Human Primary and Experimental Murine Sjögren's Syndrome. <i>Arthritis and Rheumatology</i> , 2018 , 70, 1265-1275	9.5	10
75	Sclerostin and Antisclerostin Antibody Serum Levels Predict the Presence of Axial Spondyloarthritis in Patients with Inflammatory Bowel Disease. <i>Journal of Rheumatology</i> , 2018 , 45, 630-637	4.1	15
74	Response to: 'Artery tertiary lymphoid organs in giant cell arteritis are not exclusively located in the media of temporal arteries' by Graver. <i>Annals of the Rheumatic Diseases</i> , 2018 , 77, e17	2.4	2
73	The Emerging Role of IL-1 Inhibition in Patients Affected by Rheumatoid Arthritis and Diabetes. <i>Reviews on Recent Clinical Trials</i> , 2018 , 13, 210-214	1.2	16
72	Cardiovascular Disease in Primary Sjögren's Syndrome. <i>Reviews on Recent Clinical Trials</i> , 2018 , 13, 164-169	1.2	3
71	Dose adjustments and discontinuation in TNF inhibitors treated patients: when and how. A systematic review of literature. <i>Rheumatology</i> , 2018 , 57, vii23-vii31	3.9	6
70	Blocking CD248 molecules in perivascular stromal cells of patients with systemic sclerosis strongly inhibits their differentiation toward myofibroblasts and proliferation: a new potential target for antifibrotic therapy. <i>Arthritis Research and Therapy</i> , 2018 , 20, 223	5.7	19
69	Interstitial lung disease in systemic sclerosis: current and future treatment. <i>Rheumatology International</i> , 2017 , 37, 853-863	3.6	55
68	Ectopic expression of CXCL13, BAFF, APRIL and LT- α s associated with artery tertiary lymphoid organs in giant cell arteritis. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 235-243	2.4	44
67	Dysbiosis and zonulin upregulation alter gut epithelial and vascular barriers in patients with ankylosing spondylitis. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 1123-1132	2.4	140
66	New insights into the pathogenesis of giant cell arteritis. <i>Autoimmunity Reviews</i> , 2017 , 16, 675-683	13.6	30
65	Pharmacological stress, rest perfusion and delayed enhancement cardiac magnetic resonance identifies very early cardiac involvement in systemic sclerosis patients of recent onset. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 1247-1260	2.3	10
64	Brief Report: Functional Interaction of Endoplasmic Reticulum Aminopeptidase 2 and HLA-B27 Activates the Unfolded Protein Response. <i>Arthritis and Rheumatology</i> , 2017 , 69, 1009-1015	9.5	12
63	Subclinical atherosclerosis and history of cardiovascular events in Italian patients with rheumatoid arthritis: Results from a cross-sectional, multicenter GIRACS (Gruppo Italiano di Ricerca in Reumatologia Clinica e Sperimentale) study. <i>Medicine (United States)</i> , 2017 , 96, e8180	1.8	15

62	CD4 T lymphocyte autophagy is upregulated in the salivary glands of primary Sjögren's syndrome patients and correlates with focus score and disease activity. <i>Arthritis Research and Therapy</i> , 2017 , 19, 178	5.7	27
61	Advances in immunopathogenesis of macrophage activation syndrome during rheumatic inflammatory diseases: toward new therapeutic targets?. <i>Expert Review of Clinical Immunology</i> , 2017 , 13, 1041-1047	5.1	27
60	Prevalence of type 2 diabetes and impaired fasting glucose in patients affected by rheumatoid arthritis: Results from a cross-sectional study. <i>Medicine (United States)</i> , 2017 , 96, e7896	1.8	32
59	International consensus: What else can we do to improve diagnosis and therapeutic strategies in patients affected by autoimmune rheumatic diseases (rheumatoid arthritis, spondyloarthritis, systemic sclerosis, systemic lupus erythematosus, antiphospholipid syndrome and Sjogren's syndrome)? The unmet needs and the clinical grey zone in autoimmune disease management.	13.6	84
58	IL-17 polarization of MAIT cells is derived from the activation of two different pathways. <i>European Journal of Immunology</i> , 2017 , 47, 2002-2003	6.1	17
57	Prognostic factors of macrophage activation syndrome, at the time of diagnosis, in adult patients affected by autoimmune disease: Analysis of 41 cases collected in 2 rheumatologic centers. <i>Autoimmunity Reviews</i> , 2017 , 16, 16-21	13.6	49
56	Adalimumab efficacy in enteropathic spondyloarthritis: A 12-mo observational multidisciplinary study. <i>World Journal of Gastroenterology</i> , 2017 , 23, 7139-7149	5.6	10
55	Motor cortex tRNS improves pain, affective and cognitive impairment in patients with fibromyalgia: preliminary results of a randomised sham-controlled trial. <i>Clinical and Experimental Rheumatology</i> , 2017 , 35 Suppl 105, 100-105	2.2	14
54	The role of the gastrointestinal tract in the pathogenesis of rheumatic diseases. <i>Best Practice and Research in Clinical Rheumatology</i> , 2016 , 30, 889-900	5.3	16
53	Clinical efficacy of α integrin block with natalizumab in ankylosing spondylitis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 2053-2054	2.4	14
52	Interleukin-9 Overexpression and Th9 Polarization Characterize the Inflamed Gut, the Synovial Tissue, and the Peripheral Blood of Patients With Psoriatic Arthritis. <i>Arthritis and Rheumatology</i> , 2016 , 68, 1922-31	9.5	59
51	Perivascular Cells in Diffuse Cutaneous Systemic Sclerosis Overexpress Activated ADAM12 and Are Involved in Myofibroblast Transdifferentiation and Development of Fibrosis. <i>Journal of Rheumatology</i> , 2016 , 43, 1340-9	4.1	33
50	Patient-reported impact of spondyloarthritis on work disability and working life: the ATLANTIS survey. <i>Arthritis Research and Therapy</i> , 2016 , 18, 78	5.7	35
49	IL-1 β at the crossroad between rheumatoid arthritis and type 2 diabetes: may we kill two birds with one stone?. <i>Expert Review of Clinical Immunology</i> , 2016 , 12, 849-55	5.1	39
48	Safety and efficacy of intra-articular anti-tumor necrosis factor α agents compared to corticosteroids in a treat-to-target strategy in patients with inflammatory arthritis and monoarthritis flare. <i>International Journal of Immunopathology and Pharmacology</i> , 2016 , 29, 252-66	3	28
47	Subclinical gut inflammation in ankylosing spondylitis. <i>Current Opinion in Rheumatology</i> , 2016 , 28, 89-96	5.3	46
46	Intestinal dysbiosis and innate immune responses in axial spondyloarthritis. <i>Current Opinion in Rheumatology</i> , 2016 , 28, 352-8	5.3	18
45	Adult-onset Still's disease: evaluation of prognostic tools and validation of the systemic score by analysis of 100 cases from three centers. <i>BMC Medicine</i> , 2016 , 14, 194	11.4	85

44	Endothelial progenitor cells: Are they displaying a function in autoimmune disorders?. <i>Mechanisms of Ageing and Development</i> , 2016 , 159, 44-48	5.6	6
43	Adult-onset Still's disease: an Italian multicentre retrospective observational study of manifestations and treatments in 245 patients. <i>Clinical Rheumatology</i> , 2016 , 35, 1683-9	3.9	52
42	Mesenchymal stromal cells and rheumatic diseases: new tools from pathogenesis to regenerative therapies. <i>Cytotherapy</i> , 2015 , 17, 832-49	4.8	14
41	Potential involvement of IL-9 and Th9 cells in the pathogenesis of rheumatoid arthritis. <i>Rheumatology</i> , 2015 , 54, 2264-72	3.9	63
40	Type 3 innate lymphoid cells producing IL-17 and IL-22 are expanded in the gut, in the peripheral blood, synovial fluid and bone marrow of patients with ankylosing spondylitis. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1739-47	2.4	184
39	Difference in the expression of IL-9 and IL-17 correlates with different histological pattern of vascular wall injury in giant cell arteritis. <i>Rheumatology</i> , 2015 , 54, 1596-604	3.9	45
38	Use of Rituximab in the Management of Sjögren's Syndrome. <i>Current Treatment Options in Rheumatology</i> , 2015 , 1, 277-291	1.3	3
37	Cell Immunity in Inflammatory Vasculitis. <i>Current Immunology Reviews</i> , 2015 , 11, 3-11	1.3	
36	Brief Report: Intestinal Dysbiosis in Ankylosing Spondylitis. <i>Arthritis and Rheumatology</i> , 2015 , 67, 686-691	1.5	252
35	Increased level of H-ferritin and its imbalance with L-ferritin, in bone marrow and liver of patients with adult onset Still's disease, developing macrophage activation syndrome, correlate with the severity of the disease. <i>Autoimmunity Reviews</i> , 2015 , 14, 429-37	13.6	38
34	Macrophage phenotype in the subclinical gut inflammation of patients with ankylosing spondylitis. <i>Rheumatology</i> , 2014 , 53, 104-13	3.9	33
33	Efficacy and safety of rituximab with and without methotrexate in the treatment of rheumatoid arthritis patients: results from the GISEA register. <i>Joint Bone Spine</i> , 2014 , 81, 508-12	2.9	15
32	Evidence that autophagy, but not the unfolded protein response, regulates the expression of IL-23 in the gut of patients with ankylosing spondylitis and subclinical gut inflammation. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1566-74	2.4	122
31	Is minor salivary gland biopsy more than a diagnostic tool in primary Sjögren's syndrome? Association between clinical, histopathological, and molecular features: a retrospective study. <i>Seminars in Arthritis and Rheumatism</i> , 2014 , 44, 314-24	5.3	54
30	Activated IL-22 pathway occurs in the muscle tissues of patients with polymyositis or dermatomyositis and is correlated with disease activity. <i>Rheumatology</i> , 2014 , 53, 1307-12	3.9	12
29	Rituximab modulates IL-17 expression in the salivary glands of patients with primary Sjögren's syndrome. <i>Rheumatology</i> , 2014 , 53, 1313-20	3.9	34
28	Response to: 'IL-23 expression and activation of autophagy in synovium and PBMCs of HLA-B27 positive patients with ankylosing spondylitis' by Neerinckx et al. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, e69	2.4	1
27	The role of innate and lymphoid IL-22-producing cells in the immunopathology of primary Sjögren's syndrome. <i>Expert Review of Clinical Immunology</i> , 2014 , 10, 533-41	5.1	20

26	IL-33 is overexpressed in the inflamed arteries of patients with giant cell arteritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 258-64	2.4	44
25	IL-34 is overexpressed in the inflamed salivary glands of patients with Sjogren's syndrome and is associated with the local expansion of pro-inflammatory CD14(bright)CD16+ monocytes. <i>Rheumatology</i> , 2013 , 52, 1009-17	3.9	78
24	Efficacy and safety of rituximab treatment in early primary Sjogren's syndrome: a prospective, multi-center, follow-up study. <i>Arthritis Research and Therapy</i> , 2013 , 15, R172	5.7	112
23	Rituximab modulates the expression of IL-22 in the salivary glands of patients with primary Sjogren's syndrome. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 782-3	2.4	29
22	Increased expression of interleukin-32 in the inflamed ileum of ankylosing spondylitis patients. <i>Rheumatology</i> , 2012 , 51, 1966-72	3.9	16
21	Interleukin-22 and interleukin-22-producing NKp44+ natural killer cells in subclinical gut inflammation in ankylosing spondylitis. <i>Arthritis and Rheumatism</i> , 2012 , 64, 1869-78		98
20	Potential involvement of IL-22 and IL-22-producing cells in the inflamed salivary glands of patients with Sjogren's syndrome. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 295-301	2.4	122
19	One year study of efficacy and safety of infliximab in the treatment of patients with ocular and neurological Behçet's disease refractory to standard immunosuppressive drugs. <i>Rheumatology International</i> , 2011 , 31, 33-7	3.6	49
18	Expression of interleukin-32 in the inflamed arteries of patients with giant cell arteritis. <i>Arthritis and Rheumatism</i> , 2011 , 63, 2097-104		23
17	Over-expression of paneth cell-derived anti-microbial peptides in the gut of patients with ankylosing spondylitis and subclinical intestinal inflammation. <i>Rheumatology</i> , 2010 , 49, 2076-83	3.9	39
16	Phenotype and functional changes of Vgamma9/Vdelta2 T lymphocytes in Behçet's disease and the effect of infliximab on Vgamma9/Vdelta2 T cell expansion, activation and cytotoxicity. <i>Arthritis Research and Therapy</i> , 2010 , 12, R109	5.7	25
15	Expansion of intestinal CD4+CD25(high) Treg cells in patients with ankylosing spondylitis: a putative role for interleukin-10 in preventing intestinal Th17 response. <i>Arthritis and Rheumatism</i> , 2010 , 62, 3625-34		43
14	Overexpression of interleukin-23, but not interleukin-17, as an immunologic signature of subclinical intestinal inflammation in ankylosing spondylitis. <i>Arthritis and Rheumatism</i> , 2009 , 60, 955-65		179
13	Long-term anti-tumour necrosis factor therapy reverses the progression of carotid intima-media thickness in female patients with active rheumatoid arthritis. <i>Rheumatology International</i> , 2009 , 30, 193-8 ⁶	3.6	31
12	New approaches in the treatment of Adamantiades-Behçet's disease. <i>Current Opinion in Rheumatology</i> , 2006 , 18, 3-9	5.3	49
11	NF-kappaB protects Behçet's disease T cells against CD95-induced apoptosis up-regulating antiapoptotic proteins. <i>Arthritis and Rheumatism</i> , 2005 , 52, 2179-91		47
10	Cognitive impairment in Behçet's disease patients without overt neurological involvement. <i>Journal of the Neurological Sciences</i> , 2004 , 220, 99-104	3.2	53
9	Randomized study of subcutaneous low molecular weight heparin plus aspirin versus intravenous immunoglobulin in the treatment of recurrent fetal loss associated with antiphospholipid antibodies. <i>Arthritis and Rheumatism</i> , 2003 , 48, 728-31		166

8	Relationship between circulating E-selectin, DD genotype of angiotensin-converting-enzyme, and cardiovascular damage in central obese subjects. <i>Metabolism: Clinical and Experimental</i> , 2003 , 52, 999-1004	12,7	9
7	Vgamma9/Vdelta2 T lymphocytes in Italian patients with Behçet's disease: evidence for expansion, and tumour necrosis factor receptor II and interleukin-12 receptor beta1 expression in active disease. <i>Arthritis Research</i> , 2003 , 5, R262-8		27
6	Do immune complexes formed with autoantibodies have a role in the maintenance of immune homeostasis through interaction with FC receptors. <i>Autoimmunity</i> , 2000 , 32, 153-60	3	1
5	Enhancement of endothelial cell E-selectin expression by sera from patients with active Behçet's disease: moderate correlation with anti-endothelial cell antibodies and serum myeloperoxidase levels. <i>Clinical Immunology</i> , 1999 , 91, 330-7	9	43
4	Inhibition of Fc gamma receptors in the plasma of subjects with Down's syndrome. <i>Clinical Immunology and Immunopathology</i> , 1993 , 69, 92-6		2
3	Immune complex-mediated inhibition of lymphocyte Fc-gamma receptors in the plasma of patients with type 1 (insulin-dependent) diabetes mellitus: association with anti-ssDNA antibodies. <i>Clinical Immunology and Immunopathology</i> , 1990 , 54, 228-36		5
2	Detection of two forms of HBeAg (free-and IgG-bound HBeAg) in patients with HBe antigenemia using staphylococcus bearing protein A. <i>Liver</i> , 1985 , 5, 205-11		3
1	In vitro synthesis of IgG and IgM in patients with HBsAg-positive and HBsAg-negative chronic active hepatitis. <i>Liver</i> , 1983 , 3, 207-12		1