

# Alexandre Boyer

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1215713/publications.pdf>

Version: 2024-02-01

15  
papers

415  
citations

1307594

7  
h-index

1125743

13  
g-index

15  
all docs

15  
docs citations

15  
times ranked

710  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Hippo Pathway Effectors YAP and TAZ Regulate LH Release by Pituitary Gonadotrope Cells in Mice. <i>Endocrinology</i> , 2022, 163, .	2.8	8
2	Constitutive activation of CTNNB1 results in a loss of spermatogonial stem cell activity in mice. <i>PLoS ONE</i> , 2021, 16, e0251911.	2.5	14
3	Adrenal Cortex Development and Maintenance: Knowledge Acquired From Mouse Models. <i>Endocrinology</i> , 2021, 162, .	2.8	5
4	Updating the Function of Activin A in the Fetal Testis: A New Role in Steroidogenesis. <i>Endocrinology</i> , 2020, 161, .	2.8	0
5	Targeted Disruption of Lats1 and Lats2 in Mice Impairs Adrenal Cortex Development and Alters Adrenocortical Cell Fate. <i>Endocrinology</i> , 2020, 161, .	2.8	9
6	Yes-associated protein expression in germ cells is dispensable for spermatogenesis in mice. <i>Genesis</i> , 2019, 57, e23330.	1.6	5
7	Lats1 and Lats2 are required for the maintenance of multipotency in the Müllerian duct mesenchyme. <i>Development (Cambridge)</i> , 2019, 146, .	2.5	8
8	Yes-associated protein and WW-containing transcription regulator 1 regulate the expression of sex-determining genes in Sertoli cells, but their inactivation does not cause sex reversal. <i>Biology of Reproduction</i> , 2017, 97, 162-175.	2.7	16
9	Targeted Disruption of YAP and TAZ Impairs the Maintenance of the Adrenal Cortex. <i>Endocrinology</i> , 2017, 158, 3738-3753.	2.8	18
10	mTOR Regulates Gap Junction Alpha-1 Protein Trafficking in Sertoli Cells and Is Required for the Maintenance of Spermatogenesis in Mice. <i>Biology of Reproduction</i> , 2016, 95, 13-13.	2.7	59
11	CTNNB1 Signaling in Sertoli Cells Downregulates Spermatogonial Stem Cell Activity via WNT4. <i>PLoS ONE</i> , 2012, 7, e29764.	2.5	51
12	WNT4 is required for normal ovarian follicle development and female fertility. <i>FASEB Journal</i> , 2010, 24, 3010-3025.	0.5	138
13	Seminiferous Tubule Degeneration and Infertility in Mice with Sustained Activation of WNT/CTNNB1 Signaling in Sertoli Cells. <i>Biology of Reproduction</i> , 2008, 79, 475-485.	2.7	83
14	Multiple Reproductive Defects in Fzd1-null Mice. <i>Biology of Reproduction</i> , 2008, 78, 290-290.	2.7	1
15	Sustained Activation of Wnt/ $\beta$ -catenin Signaling in Sertoli Cells Causes Seminiferous Tubule Degeneration. <i>Biology of Reproduction</i> , 2008, 78, 297-297.	2.7	0