

# KubÄ°lay KarÄidaÄ

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/12166866/publications.pdf>

Version: 2024-02-01

12  
papers

1,239  
citations

1683934

5  
h-index

1372474

10  
g-index

12  
all docs

12  
docs citations

12  
times ranked

1944  
citing authors

#	ARTICLE	IF	CITATIONS
1	The risk of cardiovascular events in patients with metabolic syndrome: The results of a population based prospective cohort study (PURE Turkey). <i>Anatolian Journal of Cardiology</i> , 2020, 24, 192-200.	0.5	3
2	Prevalence of metabolic syndrome in middle school children and evaluation of components of metabolic syndrome. <i>Sisli Etfal Hastanesi Tip Bulteni</i> , 2018, 53, 403-408.	0.1	1
3	A Comparison of hs-CRP Levels in New Diabetes Groups Diagnosed Based on FPG, 2-hPG, or HbA1c Criteria. <i>Journal of Diabetes Research</i> , 2016, 2016, 1-9.	1.0	15
4	Availability, affordability, and consumption of fruits and vegetables in 18 countries across income levels: findings from the Prospective Urban Rural Epidemiology (PURE) study. <i>The Lancet Global Health</i> , 2016, 4, e695-e703.	2.9	287
5	HDL-C levels in newly diagnosed type 2 diabetes in Turkey. <i>International Journal of Diabetes in Developing Countries</i> , 2015, 35, 184-188.	0.3	0
6	Abdominal Obesity Perception in Turkish Population. <i>Journal of Nutrition Education and Behavior</i> , 2015, 47, e11.	0.3	0
7	The Effect of GHR/exon-3 Polymorphism and Serum GH, IGF-1 and IGFBP-3 Levels in Diabetes and Coronary Heart Disease. <i>In Vivo</i> , 2015, 29, 371-8.	0.6	2
8	Twelve-year trends in the prevalence and risk factors of diabetes and prediabetes in Turkish adults. <i>European Journal of Epidemiology</i> , 2013, 28, 169-180.	2.5	487
9	Determination of nerve conduction abnormalities in patients with impaired glucose tolerance. <i>Neurological Sciences</i> , 2009, 30, 281-289.	0.9	14
10	Population-Based Study of Diabetes and Risk Characteristics in Turkey: Results of the Turkish Diabetes Epidemiology Study (TURDEP). <i>Diabetes Care</i> , 2002, 25, 1551-1556.	4.3	425
11	Comparison of Acarbose and Gliclazide as First-line Agents in Patients with Type 2 Diabetes. <i>Current Medical Research and Opinion</i> , 2000, 16, 296-306.	0.9	3
12	Relationship of Leukoaraiosis to Vascular Risk Factors and Lesion Type in Stroke Patients. <i>Annals of Saudi Medicine</i> , 1995, 15, 107-109.	0.5	2