

# Sangeeta Chavan

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/12177896/publications.pdf>

Version: 2024-02-01

17  
papers

4,296  
citations

623574

14  
h-index

1058333

14  
g-index

17  
all docs

17  
docs citations

17  
times ranked

5242  
citing authors

#	ARTICLE	IF	CITATIONS
1	Transcutaneous auricular vagus nerve stimulation reduces pain and fatigue in patients with systemic lupus erythematosus: a randomised, double-blind, sham-controlled pilot trial. <i>Annals of the Rheumatic Diseases</i> , 2021, 80, 203-208.	0.5	82
2	Choline Acetyltransferase Administration Decrease Blood Pressure in a Murine Model of Hypertension. <i>FASEB Journal</i> , 2021, 35, .	0.2	0
3	IL-1 $\beta$ -Induced Thermoregulation and Vagus Nerve Activity is Mediated by Transient Receptor Potential Ankyrin 1. <i>FASEB Journal</i> , 2021, 35, .	0.2	0
4	Renal expression and serum levels of high mobility group box 1 protein in lupus nephritis. <i>Arthritis Research and Therapy</i> , 2012, 14, R36.	1.6	64
5	Successful therapy with anti-HMGB1 monoclonal antibodies in two separate experimental arthritis models. <i>Annals of the Rheumatic Diseases</i> , 2011, 70, A77-A78.	0.5	0
6	Acetylcholine-Synthesizing T Cells Relay Neural Signals in a Vagus Nerve Circuit. <i>Science</i> , 2011, 334, 98-101.	6.0	1,158
7	Galantamine Alleviates Inflammation and Other Obesity-Associated Complications in High-Fat Diet-Fed Mice. <i>Molecular Medicine</i> , 2011, 17, 599-606.	1.9	96
8	High-Mobility Group Box-1 Protein (HMGB1) Is Increased in Antineutrophilic Cytoplasmic Antibody (ANCA)-Associated Vasculitis with Renal Manifestations. <i>Molecular Medicine</i> , 2011, 17, 29-35.	1.9	53
9	Protective targeting of high mobility group box chromosomal protein 1 in a spontaneous arthritis model. <i>Arthritis and Rheumatism</i> , 2010, 62, 2963-2972.	6.7	49
10	A critical cysteine is required for HMGB1 binding to Toll-like receptor 4 and activation of macrophage cytokine release. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 11942-11947.	3.3	705
11	The Selective $\beta_7$ Agonist GTS-21 Attenuates Cytokine Production in Human Whole Blood and Human Monocytes Activated by Ligands for TLR2, TLR3, TLR4, TLR9, and RAGE. <i>Molecular Medicine</i> , 2009, 15, 195-202.	1.9	175
12	Brain acetylcholinesterase activity controls systemic cytokine levels through the cholinergic anti-inflammatory pathway. <i>Brain, Behavior, and Immunity</i> , 2009, 23, 41-45.	2.0	378
13	Splenic nerve is required for cholinergic antiinflammatory pathway control of TNF in endotoxemia. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 11008-11013.	3.3	659
14	Modulation of TNF Release by Choline Requires $\beta_7$ Subunit Nicotinic Acetylcholine Receptor-Mediated Signaling. <i>Molecular Medicine</i> , 2008, 14, 567-574.	1.9	288
15	Transcutaneous vagus nerve stimulation reduces serum high mobility group box 1 levels and improves survival in murine sepsis*. <i>Critical Care Medicine</i> , 2007, 35, 2762-2768.	0.4	216
16	Cholinergic Anti-Inflammatory Pathway Activity and High Mobility Group Box-1 (HMGB1) Serum Levels in Patients with Rheumatoid Arthritis. <i>Molecular Medicine</i> , 2007, 13, 210-215.	1.9	162
17	Transcutaneous vagus nerve stimulation reduces serum high mobility group box 1 levels and improves survival in murine sepsis *. <i>Critical Care Medicine</i> , 2007, 35, 2762-2768.	0.4	211