

# Keke Zheng

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/12179538/publications.pdf>

Version: 2024-02-01

9  
papers

307  
citations

1163117  
8  
h-index

1474206  
9  
g-index

9  
all docs

9  
docs citations

9  
times ranked

303  
citing authors

#	ARTICLE	IF	CITATIONS
1	Taurine alone or in combination with fish protein hydrolysate affects growth performance, taurine transport and metabolism in juvenile turbot ( <i>Scophthalmus maximus</i> L.). <i>Aquaculture Nutrition</i> , 2019, 25, 396-405.	2.7	18
2	Cloning and characterization of fatty acid transport proteins in Japanese seabass <i>Lateolabrax japonicus</i> , and their gene expressions in response to dietary arachidonic acid. <i>Aquaculture Research</i> , 2017, 48, 5718-5728.	1.8	9
3	Dietary arachidonic acid differentially regulates the gonadal steroidogenesis in the marine teleost, tongue sole ( <i>Cynoglossus semilaevis</i> ), depending on fish gender and maturation stage. <i>Aquaculture</i> , 2017, 468, 378-385.	3.5	63
4	Cloning and characterization of fatty acid-binding proteins (fabps) from Japanese seabass ( <i>Lateolabrax japonicus</i> ). <i>Biochemistry and Physiology - B Biochemistry and Molecular Biology</i> , 2017, 204, 27-34.	1.6	21
5	Application of different types of protein hydrolysate in high plant protein diets for juvenile turbot ( <i>Scophthalmus maximus</i> ). <i>Aquaculture Research</i> , 2017, 48, 2945-2953.	1.8	14
6	The effect of ultrafiltered fish protein hydrolysate levels on the liver and muscle metabolic profile of juvenile turbot ( <i>Scophthalmus maximus</i> L.) by <sup>1</sup> H NMR-based metabolomics studies. <i>Aquaculture Research</i> , 2017, 48, 3515-3527.	1.8	20
7	Graded levels of fish protein hydrolysate in high plant diets for turbot ( <i>Scophthalmus maximus</i> ): effects on growth performance and lipid accumulation. <i>Aquaculture</i> , 2016, 454, 140-147.	3.5	95
8	Feeding Rates Affect Expression of Heat Shock Protein 70 in Green Sturgeon Fry. <i>North American Journal of Aquaculture</i> , 2015, 77, 206-210.	1.4	8
9	Effect of size-fractionated fish protein hydrolysate on growth and feed utilization of turbot ( <i>Scophthalmus maximus</i> L.). <i>Aquaculture Research</i> , 2013, 44, 895-902.	1.8	59