

Maria K Svensson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1218795/publications.pdf>

Version: 2024-02-01

51
papers

1,643
citations

361413
20
h-index

302126
39
g-index

54
all docs

54
docs citations

54
times ranked

3392
citing authors

#	ARTICLE	IF	CITATIONS
1	Cost-effectiveness of proprotein convertase subtilisin/kexin type 9 inhibition with evolocumab in patients with a history of myocardial infarction in Sweden. <i>European Heart Journal Quality of Care & Clinical Outcomes</i> , 2022, 8, 31-38.	4.0	13
2	Estimation of kidney function in patients with primary neuromuscular diseases: is serum cystatin C a better marker of kidney function than creatinine?. <i>Journal of Nephrology</i> , 2022, 35, 493-503.	2.0	9
3	Role of Estrogen and Its Receptors in Adipose Tissue Glucose Metabolism in Pre- and Postmenopausal Women. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, 107, e1879-e1889.	3.6	11
4	CDKN2C expression in adipose tissue is reduced in type II diabetes and central obesity: impact on adipocyte differentiation and lipid storage?. <i>Translational Research</i> , 2022, 242, 105-121.	5.0	10
5	FC028: Characterization of Adipocyte Size and Glucose Uptake in Adipose Tissue with Focus on Different Perirenal Fat Depots. <i>Nephrology Dialysis Transplantation</i> , 2022, 37, .	0.7	0
6	MO190: Experiences of Being a Living Kidney-Donor-Perceived and Measured Stress Levels before and after Kidney Donation. <i>Nephrology Dialysis Transplantation</i> , 2022, 37, .	0.7	1
7	MO169: Three Doses of SARS-CoV-2 BNT162B2 mRNA Vaccine Results in an Enhanced Immunologic B- And T-Cell Response in Haemodialysis Patients. <i>Nephrology Dialysis Transplantation</i> , 2022, 37, .	0.7	0
8	Burden of Treatment-Induced Peripheral Neuropathy in Patients with Multiple Myeloma in Sweden. <i>Acta Haematologica</i> , 2021, 144, 519-527.	1.4	0
9	Cardiovascular Event Rates After Myocardial Infarction or Ischaemic Stroke in Patients with Additional Risk Factors: A Retrospective Population-Based Cohort Study. <i>Advances in Therapy</i> , 2021, 38, 4695-4708.	2.9	4
10	Humoral and cellular response to SARS-CoV-2 BNT162b2 mRNA vaccine in hemodialysis patients. <i>BMC Immunology</i> , 2021, 22, 70.	2.2	15
11	Statin dose titration patterns and subsequent major cardiovascular events in very high-risk patients: estimates from Swedish population-based registry data. <i>European Heart Journal Quality of Care & Clinical Outcomes</i> , 2020, 6, 323-331.	4.0	10
12	3-Hydroxyisobutyrate, A Strong Marker of Insulin Resistance in Type 2 Diabetes and Obesity That Modulates White and Brown Adipocyte Metabolism. <i>Diabetes</i> , 2020, 69, 1903-1916.	0.6	42
13	Patients with moderate chronic kidney disease without heart disease have reduced coronary flow velocity reserve. <i>ESC Heart Failure</i> , 2020, 7, 2797-2806.	3.1	8
14	Real-World Effectiveness of Anti-Resorptive Treatment in Patients With Incident Fragility Fracturesâ€”The STORM Cohortâ€”A Swedish Retrospective Observational Study. <i>Journal of Bone and Mineral Research</i> , 2020, 37, 649-659.	2.8	2
15	Impact of Abdominal Aortic Calcification on Central Haemodynamics and Decline of Glomerular Filtration Rate in Patients with Chronic Kidney Disease Stages 3 and 4. <i>Kidney and Blood Pressure Research</i> , 2019, 44, 950-960.	2.0	6
16	Age- and sex-specific reference values for non-HDL cholesterol and remnant cholesterol derived from the Nordic Reference Interval Project (NORIP). <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2019, 79, 39-42.	1.2	9
17	Cardiovascular event rates in a high atherosclerotic cardiovascular disease risk population: estimates from Swedish population-based register data. <i>European Heart Journal Quality of Care & Clinical Outcomes</i> , 2019, 5, 225-232.	4.0	30
18	Abdominal Aortic Calcifications Predict Survival in Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2018, 38, 366-373.	2.3	14

#	ARTICLE	IF	CITATIONS
19	Treatment target re-classification of subjects comparing estimation of low-density lipoprotein cholesterol by the Friedewald equation and direct measurement of LDL-cholesterol. Upsala Journal of Medical Sciences, 2018, 123, 94-99.	0.9	6
20	Statement of Retraction. The SNARE Protein SNAP23 and the SNARE-Interacting Protein Munc18c in Human Skeletal Muscle Are Implicated in Insulin Resistance/Type 2 Diabetes. Diabetes 2010;59:1870-1878. DOI: 10.2337/db09-1503. PMID: 20460426. Diabetes, 2017, 66, 1426-1426.	0.6	2
21	Decreased systolic blood pressure is associated with increased risk of all-cause mortality in patients with type 2 diabetes and renal impairment: A nationwide longitudinal observational study of 27,732 patients based on the Swedish National Diabetes Register. Diabetes and Vascular Disease Research, 2017, 14, 226-235.	2.0	2
22	Reduced renal function in patients with Myotonic Dystrophy type 1 and the association to CTG expansion and other potential risk factors for chronic kidney disease. Neuromuscular Disorders, 2017, 27, 1038-1042.	0.6	5
23	Overweight, hyperglycemia and tobacco use are modifiable risk factors for onset of retinopathy 9 and 17 years after the diagnosis of diabetes - A retrospective observational nation-wide cohort study. Diabetes Research and Clinical Practice, 2017, 133, 21-29.	2.8	9
24	Soluble plasma proteins ST2 and CD163 as early biomarkers of nephropathy in Swedish patients with diabetes, 15-34 years of age: a prospective cohort study. Diabetology and Metabolic Syndrome, 2017, 9, 41.	2.7	14
25	Role of cannabinoid receptor 1 in human adipose tissue for lipolysis regulation and insulin resistance. Endocrine, 2017, 55, 839-852.	2.3	39
26	Alterations in heart rate variability during everyday life are linked to insulin resistance. A role of dominating sympathetic over parasympathetic nerve activity?. Cardiovascular Diabetology, 2016, 15, 91.	6.8	25
27	Markers of fibrinolysis may predict development of lower extremity arterial disease in patients with diabetes: A longitudinal prospective cohort study with 10 years of follow-up. Diabetes and Vascular Disease Research, 2016, 13, 183-191.	2.0	2
28	Blood pressure level and risk of major cardiovascular events and all-cause of mortality in patients with type 2 diabetes and renal impairment: an observational study from the Swedish National Diabetes Register. Diabetologia, 2015, 58, 1203-1211.	6.3	18
29	Evaluation of reference genes for gene expression studies in human brown adipose tissue. Adipocyte, 2015, 4, 280-285.	2.8	17
30	Cyclosporine A and Tacrolimus Reduce the Amount of GLUT4 at the Cell Surface in Human Adipocytes: Increased Endocytosis as a Potential Mechanism for the Diabetogenic Effects of Immunosuppressive Agents. Journal of Clinical Endocrinology and Metabolism, 2014, 99, E1885-E1894.	3.6	52
31	Altered DNA Methylation and Differential Expression of Genes Influencing Metabolism and Inflammation in Adipose Tissue From Subjects With Type 2 Diabetes. Diabetes, 2014, 63, 2962-2976.	0.6	326
32	Characterization of brown adipose tissue in the human perirenal depot. Obesity, 2014, 22, 1830-1837.	3.0	45
33	FKBP5 expression in human adipose tissue increases following dexamethasone exposure and is associated with insulin resistance. Metabolism: Clinical and Experimental, 2014, 63, 1198-1208.	3.4	81
34	End-stage renal disease and low level exposure to lead, cadmium and mercury; a population-based, prospective nested case-referent study in Sweden. Environmental Health, 2013, 12, 9.	4.0	75
35	Ongoing treatment with renin-angiotensin-aldosterone-blocking agents does not predict normoalbuminuric renal impairment in a general type 2 diabetes population. Journal of Diabetes and Its Complications, 2013, 27, 229-234.	2.3	19
36	The immunosuppressive agents rapamycin, cyclosporin A and tacrolimus increase lipolysis, inhibit lipid storage and alter expression of genes involved in lipid metabolism in human adipose tissue. Molecular and Cellular Endocrinology, 2013, 365, 260-269.	3.2	63

#	ARTICLE	IF	CITATIONS
37	Albuminuria and renal function as predictors of cardiovascular events and mortality in a general population of patients with type 2 diabetes: A nationwide observational study from the Swedish National Diabetes Register. <i>Diabetes and Vascular Disease Research</i> , 2013, 10, 520-529.	2.0	74
38	Pharmacokineticâ€”Pharmacodynamic Assessment of the Interrelationships Between Tesaglitazar Exposure and Renal Function in Patients With Type 2 Diabetes Mellitus. <i>Journal of Clinical Pharmacology</i> , 2012, 52, 1317-1327.	2.0	14
39	mTOR inhibition with rapamycin causes impaired insulin signalling and glucose uptake in human subcutaneous and omental adipocytes. <i>Molecular and Cellular Endocrinology</i> , 2012, 355, 96-105.	3.2	84
40	Effects of hyperinsulinemia on lipoprotein lipase, angiopoietin-like protein 4, and glycosylphosphatidylinositol-anchored high-density lipoprotein binding protein 1 in subjects with and without type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2012, 61, 652-660.	3.4	34
41	Effects of adrenaline on whole-body glucose metabolism and insulin-mediated regulation of glycogen synthase and PKB phosphorylation in human skeletal muscle. <i>Metabolism: Clinical and Experimental</i> , 2011, 60, 215-226.	3.4	26
42	Risk factors for the development of albuminuria and renal impairment in type 2 diabetes--the Swedish National Diabetes Register (NDR). <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 1236-1243.	0.7	110
43	The SNARE Protein SNAP23 and the SNARE-Interacting Protein Munc18c in Human Skeletal Muscle Are Implicated in Insulin Resistance/Type 2 Diabetes. <i>Diabetes</i> , 2010, 59, 1870-1878.	0.6	38
44	A Model for Glucose, Insulin, and Betaâ€”Cell Dynamics in Subjects With Insulin Resistance and Patients With Type 2 Diabetes. <i>Journal of Clinical Pharmacology</i> , 2010, 50, 861-872.	2.0	36
45	Reversible elevations of serum creatinine levels but no effect on glomerular filtration during treatment with the direct thrombin inhibitor AZD0837. <i>European Journal of Clinical Pharmacology</i> , 2010, 66, 903-910.	1.9	11
46	Low agreement between radio binding assays in analyzing glutamic acid decarboxylase (GAD65Ab) autoantibodies in patients classified with type 2 diabetes. <i>Autoimmunity</i> , 2009, 42, 507-514.	2.6	9
47	Cardiovascular disease in patients with renal disease: the role of statins. <i>Current Medical Research and Opinion</i> , 2009, 25, 271-285.	1.9	34
48	Tissue plasminogen activator (tPA) activity is a novel and early marker of asymptomatic LEAD in type 2 diabetes. <i>Thrombosis Research</i> , 2009, 123, 701-706.	1.7	15
49	Acute hyperinsulinemia raises plasma interleukin-6 in both nondiabetic and type 2 diabetes mellitus subjects, and this effect is inversely associated with body mass index. <i>Metabolism: Clinical and Experimental</i> , 2009, 58, 860-866.	3.4	55
50	Hydrochlorothiazide, but not Candesartan, Aggravates Insulin Resistance and Causes Visceral and Hepatic Fat Accumulation. <i>Hypertension</i> , 2008, 52, 1030-1037.	2.7	115
51	Effects of lipid-lowering treatment intensity and adherence on cardiovascular outcomes in patients with a recent myocardial infarction: a Swedish register-based study. <i>Uppsala Journal of Medical Sciences</i> , 0, 127, .	0.9	3