

Maximus Berger

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/122112/publications.pdf>

Version: 2024-02-01

29
papers

929
citations

687363

13
h-index

552781

26
g-index

30
all docs

30
docs citations

30
times ranked

1584
citing authors

#	ARTICLE	IF	CITATIONS
1	“More than skin deep” stress neurobiology and mental health consequences of racial discrimination. <i>Stress</i> , 2015, 18, 1-10.	1.8	385
2	Cortisol awakening response in patients with psychosis: Systematic review and meta-analysis. <i>Neuroscience and Biobehavioral Reviews</i> , 2016, 68, 157-166.	6.1	86
3	Preventive interventions for individuals at ultra high risk for psychosis: An updated and extended meta-analysis. <i>Clinical Psychology Review</i> , 2021, 86, 102005.	11.4	52
4	The NEURAPRO Biomarker Analysis: Long-Chain Omega-3 Fatty Acids Improve 6-Month and 12-Month Outcomes in Youths at Ultra-High Risk for Psychosis. <i>Biological Psychiatry</i> , 2020, 87, 243-252.	1.3	48
5	Allostatic load is associated with psychotic symptoms and decreases with antipsychotic treatment in patients with schizophrenia and first-episode psychosis. <i>Psychoneuroendocrinology</i> , 2018, 90, 35-42.	2.7	47
6	Cortisol Awakening Response and Acute Stress Reactivity in First Nations People. <i>Scientific Reports</i> , 2017, 7, 41760.	3.3	35
7	Comparison of erythrocyte omega-3 index, fatty acids and molecular phospholipid species in people at ultra-high risk of developing psychosis and healthy people. <i>Schizophrenia Research</i> , 2020, 226, 44-51.	2.0	27
8	Assessment of Insulin Resistance Among Drug-Naive Patients With First-Episode Schizophrenia in the Context of Hormonal Stress Axis Activation. <i>JAMA Psychiatry</i> , 2017, 74, 968.	11.0	26
9	Effects of omega-3 PUFA on immune markers in adolescent individuals at ultra-high risk for psychosis “ Results of the randomized controlled Vienna omega-3 study. <i>Schizophrenia Research</i> , 2017, 188, 110-117.	2.0	23
10	Relationship Between Polyunsaturated Fatty Acids and Psychopathology in the NEURAPRO Clinical Trial. <i>Frontiers in Psychiatry</i> , 2019, 10, 393.	2.6	22
11	Hair cortisol, allostatic load, and depressive symptoms in Australian Aboriginal and Torres Strait Islander people. <i>Stress</i> , 2019, 22, 312-320.	1.8	22
12	Mental health consequences of stress and trauma: allostatic load markers for practice and policy with a focus on Indigenous health. <i>Australasian Psychiatry</i> , 2015, 23, 644-649.	0.7	16
13	Trajectories of symptom severity and functioning over a three-year period in a psychosis high-risk sample: A secondary analysis of the Neurapro trial. <i>Behaviour Research and Therapy</i> , 2020, 124, 103527.	3.1	16
14	Treatment of social anxiety disorder and attenuated psychotic symptoms with cannabidiol. <i>BMJ Case Reports</i> , 2020, 13, e235307.	0.5	16
15	Allostatic load mediates the impact of stress and trauma on physical and mental health in Indigenous Australians. <i>Australasian Psychiatry</i> , 2016, 24, 72-75.	0.7	15
16	Novel biotherapies are needed in youth mental health. <i>Australasian Psychiatry</i> , 2017, 25, 117-120.	0.7	13
17	Relationship between allostatic load and clinical outcomes in youth at ultra-high risk for psychosis in the NEURAPRO study. <i>Schizophrenia Research</i> , 2020, 226, 38-43.	2.0	13
18	State marker properties of niacin skin sensitivity in ultra-high risk groups for psychosis - An optical reflection spectroscopy study. <i>Schizophrenia Research</i> , 2018, 192, 377-384.	2.0	12

#	ARTICLE	IF	CITATIONS
19	Cannabidiol for at risk for psychosis youth: A randomized controlled trial. <i>Microbial Biotechnology</i> , 2022, 16, 419-432.	1.7	9
20	Trauma and posttraumatic stress disorder in transcultural patients with chronic pain. <i>Neuropsychiatrie</i> , 2014, 28, 185-191.	2.5	8
21	Harmonised collection of data in youth mental health: Towards large datasets. <i>Australian and New Zealand Journal of Psychiatry</i> , 2020, 54, 46-56.	2.3	8
22	Cross-sectional association of seafood consumption, polyunsaturated fatty acids and depressive symptoms in two Torres Strait communities. <i>Nutritional Neuroscience</i> , 2020, 23, 353-362.	3.1	8
23	Supplementation with the omega-3 long chain polyunsaturated fatty acids: Changes in the concentrations of omega-3 index, fatty acids and molecular phospholipids of people at ultra high risk of developing psychosis. <i>Schizophrenia Research</i> , 2020, 226, 52-60.	2.0	8
24	Predictors of longer-term outcome in the Vienna omega-3 high-risk study. <i>Schizophrenia Research</i> , 2018, 193, 168-172.	2.0	6
25	Screening for depression in young Indigenous people: building on a unique community initiative. <i>Australian Journal of Primary Health</i> , 2018, 24, 343.	0.9	6
26	The association of plasma inflammatory markers with omega-3 fatty acids and their mediating role in psychotic symptoms and functioning: An analysis of the NEURAPRO clinical trial. <i>Brain, Behavior, and Immunity</i> , 2022, 99, 147-156.	4.1	2
27	8. Allostatic Load is Associated With Positive Symptoms in Schizophrenia and First-Episode Psychosis and Decreases With Antipsychotic Therapy. <i>Schizophrenia Bulletin</i> , 2017, 43, S9-S10.	4.3	0
28	Novel Biological Treatment Strategies. , 2019, , 221-240.		0
29	Testing the Effects of Dietary Seafood Consumption on Depressive Symptoms. <i>Methods in Molecular Biology</i> , 2020, 2138, 233-242.	0.9	0