

Masato Honda

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1222842/publications.pdf>

Version: 2024-02-01

12
papers

426
citations

1307594

7
h-index

1199594

12
g-index

12
all docs

12
docs citations

12
times ranked

514
citing authors

#	ARTICLE	IF	CITATIONS
1	Hydroxylated benzo[c]phenanthrene metabolites cause osteoblast apoptosis and skeletal abnormalities in fish. <i>Ecotoxicology and Environmental Safety</i> , 2022, 234, 113401.	6.0	8
2	Correlation between Polycyclic Aromatic Hydrocarbons in Wharf Roach (<i>Ligia spp.</i>) and Environmental Components of the Intertidal and Supralittoral Zone along the Japanese Coast. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 630.	2.6	4
3	Contamination of Heavy Metals, Polychlorinated Dibenzo-p-Dioxins/Furans and Dioxin-Like Polychlorinated Biphenyls in Wharf Roach <i>Ligia spp.</i> In Japanese Intertidal and Supratidal Zones. <i>Applied Sciences (Switzerland)</i> , 2021, 11, 1856.	2.5	3
4	Toxicities of Polycyclic Aromatic Hydrocarbons for Aquatic Animals. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 1363.	2.6	293
5	The Wharf Roach, <i>Ligia sp.</i> , A Novel Indicator of Polycyclic Aromatic Hydrocarbon Contamination in Coastal Areas. <i>International Journal of Environmental Research</i> , 2018, 12, 1-11.	2.3	15
6	High concentrations of perfluorooctane sulfonate in mucus of tiger puffer fish <i>Takifugu rubripes</i> : a laboratory exposure study. <i>Environmental Science and Pollution Research</i> , 2018, 25, 1551-1558.	5.3	4
7	Pollution of radiocesium and radiosilver in wharf roach (<i>Ligia sp.</i>) by the Fukushima Dai-ichi Nuclear Power Plant accident. <i>Journal of Radioanalytical and Nuclear Chemistry</i> , 2017, 311, 121-126.	1.5	11
8	Contaminants in Liquid Organic Fertilizers Used for Agriculture in Japan. <i>Bulletin of Environmental Contamination and Toxicology</i> , 2017, 99, 131-137.	2.7	7
9	Short-term and persistent impacts on behaviors related to locomotion, anxiety, and startle responses of Japanese medaka (<i>Oryzias latipes</i>) induced by acute, sublethal exposure to chlorpyrifos. <i>Aquatic Toxicology</i> , 2017, 192, 148-154.	4.0	40
10	Alteration of development and gene expression induced by in ovo -nanoinjection of 3-hydroxybenzo[c]phenanthrene into Japanese medaka (<i>Oryzias latipes</i>) embryos. <i>Aquatic Toxicology</i> , 2017, 182, 194-204.	4.0	12
11	Identification of perfluorooctane sulfonate binding protein in the plasma of tiger pufferfish <i>Takifugu rubripes</i> . <i>Ecotoxicology and Environmental Safety</i> , 2014, 104, 409-413.	6.0	18
12	Accumulation of organotins in wharf roach (<i>Ligia exotica</i> Roux) and its ability to serve as a biomonitoring species for coastal pollution. <i>Ecotoxicology and Environmental Safety</i> , 2013, 96, 75-79.	6.0	11