

Huiling Tan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1227517/publications.pdf>

Version: 2024-02-01

31
papers

448
citations

933447

10
h-index

752698

20
g-index

33
all docs

33
docs citations

33
times ranked

773
citing authors

#	ARTICLE	IF	CITATIONS
1	Association of polymorphisms in pre-miRNA with inflammatory biomarkers in rheumatoid arthritis in the Chinese Han population. <i>Human Immunology</i> , 2012, 73, 101-106.	2.4	77
2	Enhanced IL-6/phosphorylated STAT3 signaling is related to the imbalance of circulating T follicular helper/T follicular regulatory cells in patients with rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2018, 20, 200.	3.5	77
3	Association study of single nucleotide polymorphisms in pre-miRNA and rheumatoid arthritis in a Han Chinese population. <i>Molecular Biology Reports</i> , 2011, 38, 4913-4919.	2.3	66
4	Elevated Levels of Urinary Markers of Oxidative DNA and RNA Damage in Type 2 Diabetes with Complications. <i>Oxidative Medicine and Cellular Longevity</i> , 2016, 2016, 1-7.	4.0	34
5	Autoimmune disease associated IFIH1 single nucleotide polymorphism related with IL-18 serum levels in Chinese systemic lupus erythematosus patients. <i>Scientific Reports</i> , 2018, 8, 9442.	3.3	21
6	Association between tumor necrosis factor polymorphisms and rheumatoid arthritis as well as systemic lupus erythematosus: a meta-analysis. <i>Brazilian Journal of Medical and Biological Research</i> , 2019, 52, e7927.	1.5	16
7	IL-23R and IL-17A polymorphisms correlate with susceptibility of ankylosing spondylitis in a Southwest Chinese population. <i>Oncotarget</i> , 2017, 8, 70310-70316.	1.8	15
8	Genetic Polymorphisms of rs3077 and rs9277535 in HLA-DP associated with Systemic lupus erythematosus in a Chinese population. <i>Scientific Reports</i> , 2017, 7, 39757.	3.3	12
9	Genetic polymorphism of rs9277535 in HLA-DP associated with rheumatoid arthritis and anti-CCP production in a Chinese population. <i>Clinical Rheumatology</i> , 2018, 37, 1799-1805.	2.2	11
10	The Expression of LIGHT Was Increased and the Expression of HVEM and BTLA Were Decreased in the T Cells of Patients with Rheumatoid Arthritis. <i>PLoS ONE</i> , 2016, 11, e0155345.	2.5	11
11	Novel method for ANA quantitation using IIF imaging system. <i>Journal of Immunological Methods</i> , 2014, 404, 52-58.	1.4	10
12	Genetic Variation in miR-146a Is Not Associated with Susceptibility to IgA Nephropathy in Adults from a Chinese Han Population. <i>PLoS ONE</i> , 2015, 10, e0139554.	2.5	10
13	IKZF1 polymorphisms are associated with susceptibility, cytokine levels, and clinical features in systemic lupus erythematosus. <i>Medicine (United States)</i> , 2020, 99, e22607.	1.0	9
14	Prevalence of food allergen and aeroallergen sensitization among people in Sichuan, Western China: An 8-year observational study. <i>Journal of Clinical Laboratory Analysis</i> , 2019, 33, e22723.	2.1	8
15	Association of E26 Transformation Specific Sequence 1 Variants with Rheumatoid Arthritis in Chinese Han Population. <i>PLoS ONE</i> , 2015, 10, e0134875.	2.5	8
16	Genetic polymorphisms in HLA-DP and STAT4 are associated with IgA nephropathy in a Southwest Chinese population. <i>Oncotarget</i> , 2018, 9, 7066-7074.	1.8	8
17	Interleukin 18 -607 A/C Gene Polymorphism is Associated With Susceptibility to IgA Nephropathy in a Chinese Han Population. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2017, 25, 725-730.	1.2	7
18	Association of HLA-DP/DQ and STAT4 polymorphisms with ankylosing spondylitis in Southwest China. <i>International Immunopharmacology</i> , 2016, 39, 10-15.	3.8	6

#	ARTICLE	IF	CITATIONS
19	Prevalence of allergen-specific IgE in southern China: a multicenter research. <i>Aging</i> , 2021, 13, 18894-18911.	3.1	6
20	ETS1 polymorphism rs73013527 in relation to serum RANKL levels among patients with RA. <i>Medicine (United States)</i> , 2021, 100, e24562.	1.0	5
21	Anti-Mitotic Spindle Apparatus Autoantibodies: Prevalence and Disease Association in Chinese Population. <i>Journal of Clinical Laboratory Analysis</i> , 2016, 30, 702-708.	2.1	4
22	Full liberation of 2-Aminopurine with nucleases digestion for highly sensitive biosensing. <i>Biosensors and Bioelectronics</i> , 2022, 196, 113721.	10.1	4
23	Analysis of inflammation-related microRNA expression in patients with ankylosing spondylitis. <i>Immunologic Research</i> , 2022, 70, 23-32.	2.9	4
24	Association of genetic polymorphisms in IL-23R and IL-17A with the susceptibility to IgA nephropathy in a Chinese Han population. <i>Genes and Immunity</i> , 2022, 23, 33-41.	4.1	4
25	Association between food allergy and ankylosing spondylitis. <i>Medicine (United States)</i> , 2019, 98, e14421.	1.0	3
26	Detection of IgA Antiphospholipid Antibodies Does not Improve Thrombotic Antiphospholipid Syndrome Classification: A two-Center Study. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2022, 28, 107602962210811.	1.7	3
27	Dense fine speckled immunofluorescence pattern in a Chinese population: Prevalence and clinical association. <i>Journal of Clinical Laboratory Analysis</i> , 2022, 36, e24173.	2.1	3
28	Genetic association of E26 transformation specific sequence 1 polymorphisms with the susceptibility of primary biliary cholangitis in China. <i>Scientific Reports</i> , 2019, 9, 19680.	3.3	2
29	Differences in the Intestinal Flora of Patients with Inflammatory Bowel Disease in Southwest China. <i>Indian Journal of Microbiology</i> , 2022, 62, 384-392.	2.7	2
30	Genetic variations in LIGHT are associated with susceptibility to ankylosing spondylitis in a Chinese Han population. <i>Oncotarget</i> , 2017, 8, 91415-91424.	1.8	1
31	Impact of the polymorphism in vitamin D receptor gene BsmI and the risk of systemic lupus erythematosus. <i>Clinical Rheumatology</i> , 2016, 35, 1121-1121.	2.2	0