

Naoki Takahashi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1231569/publications.pdf>

Version: 2024-02-01

26
papers

517
citations

933447

10
h-index

677142

22
g-index

30
all docs

30
docs citations

30
times ranked

747
citing authors

#	ARTICLE	IF	CITATIONS
1	Fenofibrate reduces cisplatin-induced apoptosis by inhibiting the p53/Puma/Caspase-9 pathway and the MAPK/Caspase-8 pathway rather than by promoting autophagy in murine renal proximal tubular cells. <i>Biochemistry and Biophysics Reports</i> , 2022, 30, 101237.	1.3	1
2	PPAR- δ activation reduces cisplatin-induced apoptosis via inhibiting p53/Bax/caspase-3 pathway without modulating autophagy in murine renal proximal tubular cells. <i>Clinical and Experimental Nephrology</i> , 2021, 25, 598-607.	1.6	5
3	CD8 Positive T Lymphocyte Infiltration of Liver Metastases of Uveal Melanoma: A Case Report. <i>Frontiers in Oncology</i> , 2021, 11, 672660.	2.8	1
4	Glucocorticoid-dependent Tubulointerstitial Nephritis with IgM-positive Plasma Cells Presenting with Intracellular Crystalline Inclusions within the Rough Endoplasmic Reticulum. <i>Internal Medicine</i> , 2021, 60, 3129-3136.	0.7	1
5	Development of Myeloperoxidase Anti-neutrophil Cytoplasmic Antibody-positive Necrotizing Crescentic Glomerulonephritis in an Elderly Patient with Immunological Kidney Disease. <i>Internal Medicine</i> , 2021, , .	0.7	2
6	IgA Vasculitis Developed as an Adverse Effect of Tofacitinib Taken for Rheumatoid Arthritis. <i>Internal Medicine</i> , 2020, 59, 817-821.	0.7	11
7	Tubulointerstitial Nephritis Cases With IgM-Positive Plasma Cells. <i>Kidney International Reports</i> , 2020, 5, 1576-1580.	0.8	3
8	Preemptive HLA Antibody Screening Prior to Episodic Transplant Renal Biopsy Enables Early Diagnosis and Therapeutic Response in Asymptomatic Chronically Active Antibody-Related Rejection: A Case Report. <i>Transplantation Proceedings</i> , 2020, 52, 2750-2753.	0.6	1
9	Chronic hypoxia exacerbates diabetic glomerulosclerosis through mesangiolysis and podocyte injury in db/db mice. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, 1678-1688.	0.7	11
10	Analytical and clinical validation of rapid chemiluminescence enzyme immunoassay for urinary thioredoxin, an oxidative stress-dependent early biomarker of acute kidney injury. <i>Clinica Chimica Acta</i> , 2020, 507, 271-279.	1.1	5
11	β -Hydroxybutyrate, a ketone body, reduces the cytotoxic effect of cisplatin via activation of HDAC5 in human renal cortical epithelial cells. <i>Life Sciences</i> , 2019, 222, 125-132.	4.3	21
12	Elevated Levels of Urinary Extracellular Vesicle Fibroblast-Specific Protein 1 in Patients with Active Crescentic Glomerulonephritis. <i>Nephron</i> , 2019, 141, 177-187.	1.8	15
13	Short-chain fatty acids, GPR41 and GPR43 ligands, inhibit TNF- α -induced MCP-1 expression by modulating p38 and JNK signaling pathways in human renal cortical epithelial cells. <i>Biochemical and Biophysical Research Communications</i> , 2017, 486, 499-505.	2.1	127
14	A multicenter phase II prospective clinical trial of glucocorticoid for patients with untreated IgG4-related disease. <i>Modern Rheumatology</i> , 2017, 27, 849-854.	1.8	62
15	A case of Gitelman syndrome involving bilateral progressive renal cysts that necessitated the initiation of peritoneal dialysis. <i>Nihon Toseki Igakkai Zasshi</i> , 2017, 50, 581-586.	0.1	0
16	Renal resistive index correlates with peritubular capillary loss and arteriosclerosis in biopsy tissues from patients with chronic kidney disease. <i>Clinical and Experimental Nephrology</i> , 2015, 19, 1114-1119.	1.6	26
17	Clinical and biochemical investigation of male patients exhibiting membranous cytoplasmic bodies in biopsied kidney tissues; a pitfall in diagnosis of Fabry disease. <i>Journal of Nephrology</i> , 2015, 4, 91-6.	0.2	4
18	Telmisartan activates endogenous peroxisome proliferator-activated receptor- δ and may have anti-fibrotic effects in human mesangial cells. <i>Hypertension Research</i> , 2014, 37, 422-431.	2.7	29

#	ARTICLE	IF	CITATIONS
19	Telmisartan, a possible PPAR- γ agonist, reduces TNF- α -stimulated VEGF-C production by inhibiting the p38MAPK/HSP27 pathway in human proximal renal tubular cells. <i>Biochemical and Biophysical Research Communications</i> , 2014, 454, 320-327.	2.1	12
20	Renal redox dysregulation in AKI: application for oxidative stress marker of AKI. <i>American Journal of Physiology - Renal Physiology</i> , 2014, 307, F1342-F1351.	2.7	30
21	Severe intraglomerular detachment of podocytes in a Gitelman syndrome patient. <i>Clinical and Experimental Nephrology</i> , 2012, 16, 495-500.	1.6	10
22	Rapid remission of minimal change disease with angiotensin II antagonist treatment in a type 1 diabetic patient with no diabetic nephropathy. <i>CKJ: Clinical Kidney Journal</i> , 2008, 1, 375-376.	2.9	0
23	Quantification of Coronary Artery Calcium with Multi- ϕ Detector row CT: Assessing Interscan Variability with Different Tube Currents- ϕ Pilot Study. <i>Radiology</i> , 2003, 228, 101-106.	7.3	35
24	Complete nucleotide sequence of immunoglobulin γ 2b chain gene cloned from newborn mouse DNA. <i>Nature</i> , 1980, 283, 786-789.	27.8	104
25	Comparison of immunoglobulin .GAMMA.1 and .GAMMA.2b chain genes cloned from newborn mouse. Evolutional divergence of intervening sequences and structural basis for recognition of splicing sites.. <i>Proceedings of the Japan Academy Series B: Physical and Biological Sciences</i> , 1979, 55, 519-524.	3.8	0
26	Fanconi syndrome in an elderly patient with membranous nephropathy during treatment with the immunosuppressant mizoribine. <i>CEN Case Reports</i> , 0, , .	0.9	1