

Joanna Wykrzykowska

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1233958/joanna-wykrzykowska-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

57
papers

1,910
citations

20
h-index

43
g-index

59
ext. papers

2,367
ext. citations

5.7
avg, IF

3.82
L-index

| # | Paper | IF | Citations |
|----|--|------|-----------|
| 57 | Identification of vulnerable plaques and patients by intracoronary near-infrared spectroscopy and ultrasound (PROSPECT II): a prospective natural history study. <i>Lancet, The</i> , 2021 , 397, 985-995 | 4.0 | 45 |
| 56 | Online Quantitative Aortographic Assessment of Aortic Regurgitation After TAVR: Results of the OVAL Study. <i>JACC: Cardiovascular Interventions</i> , 2021 , 14, 531-538 | 5 | 4 |
| 55 | Novel Supreme Drug-Eluting Stents With Early Synchronized Antiproliferative Drug Delivery to Inhibit Smooth Muscle Cell Proliferation After Drug-Eluting Stents Implantation in Coronary Artery Disease: Results of the PIONEER III Randomized Clinical Trial. <i>Circulation</i> , 2021 , 143, 2143-2154 | 16.7 | 3 |
| 54 | Risk factor management and OCT characteristics of plaque vulnerability: the Holy Grail of plaque and patient vulnerability. <i>International Journal of Cardiovascular Imaging</i> , 2021 , 1 | 2.5 | |
| 53 | Safety and Efficacy of 1-Month Dual Antiplatelet Therapy (Ticagrelor + Aspirin) Followed by 23-Month Ticagrelor Monotherapy in Patients Undergoing Staged Percutaneous Coronary Intervention (A Sub-Study from GLOBAL LEADERS). <i>American Journal of Cardiology</i> , 2021 , 138, 1-10 | 3 | 2 |
| 52 | Comparative Assessment of Predictive Performance of PRECISE-DAPT, CRUSADE, and ACUITY Scores in Risk Stratifying 30-Day Bleeding Events. <i>Thrombosis and Haemostasis</i> , 2020 , 120, 1087-1095 | 7 | 8 |
| 51 | Early mobilisation after transfemoral transcatheter aortic valve implantation: results of the MobiTAVI trial. <i>Netherlands Heart Journal</i> , 2020 , 28, 240-248 | 2.2 | 5 |
| 50 | DAPT Score and the Impact of Ticagrelor Monotherapy During the Second Year After PCI. <i>JACC: Cardiovascular Interventions</i> , 2020 , 13, 634-646 | 5 | 6 |
| 49 | The Absorb bioresorbable vascular scaffold in real-world practice: long-term follow-up of the AMC Single Centre Real World PCI Registry. <i>Netherlands Heart Journal</i> , 2020 , 28, 153-160 | 2.2 | 3 |
| 48 | Association of Sex With Outcomes in Patients Undergoing Percutaneous Coronary Intervention: A Subgroup Analysis of the GLOBAL LEADERS Randomized Clinical Trial. <i>JAMA Cardiology</i> , 2020 , 5, 21-29 | 16.2 | 23 |
| 47 | Association of Pulse Pressure With Clinical Outcomes in Patients Under Different Antiplatelet Strategies After Percutaneous Coronary Intervention: Analysis of GLOBAL LEADERS. <i>Canadian Journal of Cardiology</i> , 2020 , 36, 747-755 | 3.8 | 1 |
| 46 | Will coronary artery bypass grafting remain a standard of care for elderly patients with multivessel disease in the contemporary era?. <i>Netherlands Heart Journal</i> , 2020 , 28, 457-459 | 2.2 | |
| 45 | The 2010s in clinical drug-eluting stent and bioresorbable scaffold research: a Dutch perspective. <i>Netherlands Heart Journal</i> , 2020 , 28, 78-87 | 2.2 | 1 |
| 44 | Cardiology fellows-in-training are exposed to relatively high levels of radiation in the cath lab compared with staff interventional cardiologists-insights from the RECAP trial. <i>Netherlands Heart Journal</i> , 2019 , 27, 330-333 | 2.2 | 1 |
| 43 | Identification of patients and plaques vulnerable to future coronary events with near-infrared spectroscopy intravascular ultrasound imaging: a prospective, cohort study. <i>Lancet, The</i> , 2019 , 394, 1629-1637 | 4.0 | 131 |
| 42 | Five-year clinical follow-up of the STENTYS self-apposing stent in complex coronary anatomy: a single-centre experience with report of specific angiographic indications. <i>Netherlands Heart Journal</i> , 2018 , 26, 263-271 | 2.2 | 6 |
| 41 | First-in-Man Trial of SiO Inert-Coated Bare Metal Stent System in Native Coronary Stenosis - The AXETIS FIM Trial. <i>Circulation Journal</i> , 2018 , 82, 477-485 | 2.9 | 2 |

| | | | |
|----|---|------|-----|
| 40 | Angiographic late lumen loss revisited: impact on long-term target lesion revascularization. <i>European Heart Journal</i> , 2018 , 39, 3381-3389 | 9.5 | 17 |
| 39 | Midterm clinical outcomes with everolimus-eluting bioresorbable scaffolds versus everolimus-eluting metallic stents for percutaneous coronary interventions: a meta-analysis of randomised trials. <i>EuroIntervention</i> , 2018 , 13, 1565-1573 | 3.1 | 29 |
| 38 | Appropriate use criteria for optical coherence tomography guidance in percutaneous coronary interventions : Recommendations of the working group of interventional cardiology of the Netherlands Society of Cardiology. <i>Netherlands Heart Journal</i> , 2018 , 26, 473-483 | 2.2 | 5 |
| 37 | Trends in patient characteristics and clinical outcome over 8 years of transcatheter aortic valve implantation. <i>Netherlands Heart Journal</i> , 2018 , 26, 445-453 | 2.2 | 7 |
| 36 | Mid-term and long-term safety and efficacy of bioresorbable vascular scaffolds versus metallic everolimus-eluting stents in coronary artery disease: A weighted meta-analysis of seven randomised controlled trials including 5577 patients. <i>Netherlands Heart Journal</i> , 2017 , 25, 429-438 | 2.2 | 7 |
| 35 | Recommendations for the use of bioresorbable vascular scaffolds in percutaneous coronary interventions : 2017 revision. <i>Netherlands Heart Journal</i> , 2017 , 25, 419-428 | 2.2 | 8 |
| 34 | Change in lumen eccentricity and asymmetry after treatment with Absorb bioresorbable vascular scaffolds in the ABSORB cohort B trial: a five-year serial optical coherence tomography imaging study. <i>EuroIntervention</i> , 2017 , 12, e2244-e2252 | 3.1 | 12 |
| 33 | Is quantitative coronary angiography reliable in assessing the late lumen loss of the everolimus-eluting bioresorbable polylactide scaffold in comparison with the cobalt-chromium metallic stent?. <i>EuroIntervention</i> , 2017 , 13, e585-e594 | 3.1 | 5 |
| 32 | First-in-man six-month results of a surface-modified coronary stent system in native coronary stenosis. <i>EuroIntervention</i> , 2017 , 12, 2118-2127 | 3.1 | 1 |
| 31 | Quantification by optical coherence tomography imaging of the ablation volume obtained with the Orbital Atherectomy System in calcified coronary lesions. <i>EuroIntervention</i> , 2016 , 12, 1126-1134 | 3.1 | 17 |
| 30 | Final results of a self-apposing paclitaxel-eluting stent for the Percutaneous treatment of de novo lesions in native bifurcated coronary arteries study. <i>EuroIntervention</i> , 2016 , 12, 356-8 | 3.1 | 13 |
| 29 | Long-term effectiveness and safety of the sirolimus-eluting BiOSS LIM dedicated bifurcation stent in the treatment of distal left main stenosis: an international registry. <i>EuroIntervention</i> , 2016 , 12, 1246-1254 | 3.1 | 11 |
| 28 | Regular drug-eluting stents versus the dedicated coronary bifurcation sirolimus-eluting BiOSS LIM stent: the randomised, multicentre, open-label, controlled POLBOS II trial. <i>EuroIntervention</i> , 2016 , 12, e1404-e1412 | 3.1 | 23 |
| 27 | Percutaneous Intervention for Concurrent Chronic Total Occlusions in Patients With STEMI: The EXPLORE Trial. <i>Journal of the American College of Cardiology</i> , 2016 , 68, 1622-1632 | 15.1 | 205 |
| 26 | Appropriate use of bioresorbable vascular scaffolds in percutaneous coronary interventions: a recommendation from experienced users : A position statement on the use of bioresorbable vascular scaffolds in the Netherlands. <i>Netherlands Heart Journal</i> , 2015 , 23, 161-5 | 2.2 | 28 |
| 25 | Pressure-controlled intermittent coronary sinus occlusion (PICSO) in acute ST-segment elevation myocardial infarction: results of the Prepare RAMSES safety and feasibility study. <i>EuroIntervention</i> , 2015 , 11, 37-44 | 3.1 | 21 |
| 24 | Six-month clinical outcomes of the Tryton Side Branch Stent for the treatment of bifurcation lesions. <i>Netherlands Heart Journal</i> , 2012 , 20, 439-46 | 2.2 | 14 |
| 23 | Vascular tissue reaction to acute malapposition in human coronary arteries: sequential assessment with optical coherence tomography. <i>Circulation: Cardiovascular Interventions</i> , 2012 , 5, 20-9, S1-8 | 6 | 86 |

| | | | |
|----|---|------|-----|
| 22 | NIRS and IVUS for characterization of atherosclerosis in patients undergoing coronary angiography. <i>JACC: Cardiovascular Imaging</i> , 2011 , 4, 647-55 | 8.4 | 65 |
| 21 | Six-month clinical follow-up of the Tryton side branch stent for the treatment of bifurcation lesions: a two center registry analysis. <i>Catheterization and Cardiovascular Interventions</i> , 2011 , 77, 798-806 | 2.7 | 29 |
| 20 | The prognostic utility of the SYNTAX score on 1-year outcomes after revascularization with zotarolimus- and everolimus-eluting stents: a substudy of the RESOLUTE All Comers Trial. <i>JACC: Cardiovascular Interventions</i> , 2011 , 4, 432-41 | 5 | 83 |
| 19 | Delayed coverage in malapposed and side-branch struts with respect to well-apposed struts in drug-eluting stents: in vivo assessment with optical coherence tomography. <i>Circulation</i> , 2011 , 124, 612-23 | 16.7 | 104 |
| 18 | The outcome of bifurcation lesion stenting using a biolimus-eluting stent with a bio-degradable polymer compared to a sirolimus-eluting stent with a durable polymer. <i>EuroIntervention</i> , 2011 , 6, 928-35 | 3.1 | 16 |
| 17 | Intravascular ultrasound radiofrequency analysis after optimal coronary stenting with initial quantitative coronary angiography guidance: an ATHEROREMO sub-study. <i>EuroIntervention</i> , 2011 , 6, 977-84 | 3.1 | 10 |
| 16 | First-in-man evaluation of intravascular optical frequency domain imaging (OFDI) of Terumo: a comparison with intravascular ultrasound and quantitative coronary angiography. <i>EuroIntervention</i> , 2011 , 6, 1037-45 | 3.1 | 81 |
| 15 | Implantation of the biodegradable polymer biolimus-eluting stent in patients with high SYNTAX score is associated with decreased cardiac mortality compared to a permanent polymer sirolimus-eluting stent: two year follow-up results from the "all-comers" LEADERS trial. <i>EuroIntervention</i> , 2011 , 7, 605-13 | 3.1 | 20 |
| 14 | First clinical evaluation of a luminal self-expanding shield in patients with intermediate coronary lesions. <i>EuroIntervention</i> , 2011 , 7, 780-8 | 3.1 | 3 |
| 13 | The three year follow-up of the randomised "all-comers" trial of a biodegradable polymer biolimus-eluting stent versus permanent polymer sirolimus-eluting stent (LEADERS). <i>EuroIntervention</i> , 2011 , 7, 789-95 | 3.1 | 34 |
| 12 | A new tool for the risk stratification of patients with complex coronary artery disease: the Clinical SYNTAX Score. <i>Circulation: Cardiovascular Interventions</i> , 2010 , 3, 317-26 | 6 | 199 |
| 11 | Very late stent thrombosis of a NIR Advance bare metal stent nine years after implantation. <i>EuroIntervention</i> , 2010 , 5, 752-3 | 3.1 | 1 |
| 10 | Optical coherence tomography for the assessment of pericardium covered stents for the treatment of degenerated saphenous vein grafts. <i>EuroIntervention</i> , 2010 , 6, 78-85 | 3.1 | 11 |
| 9 | How should I treat an unusual referral for heart transplantation?. <i>EuroIntervention</i> , 2010 , 5, 861-5 | 3.1 | 2 |
| 8 | Vascular effects of a low-carbohydrate high-protein diet. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009 , 106, 15418-23 | 11.5 | 132 |
| 7 | Exercise-induced expression of VEGF and salvation of myocardium in the early stage of myocardial infarction. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2009 , 296, H389-95 | 5.2 | 40 |
| 6 | Imaging of inflamed and vulnerable plaque in coronary arteries with ¹⁸ F-FDG PET/CT in patients with suppression of myocardial uptake using a low-carbohydrate, high-fat preparation. <i>Journal of Nuclear Medicine</i> , 2009 , 50, 563-8 | 8.9 | 216 |
| 5 | Impact of aging on the angiogenic potential of the myocardium: implications for angiogenic therapies with emphasis on sirtuin agonists. <i>Recent Patents on Cardiovascular Drug Discovery</i> , 2009 , 4, 119-32 | | 2 |

| | | | |
|---|--|------|----|
| 4 | Biolimus-eluting biodegradable polymer versus sirolimus-eluting permanent polymer stent performance in long lesions: results from the LEADERS multicentre trial substudy. <i>EuroIntervention</i> , 2009 , 5, 310-7 | 3.1 | 12 |
| 3 | Mitral annuloplasty causing left circumflex injury and infarction: novel use of intravascular ultrasound to diagnose suture injury. <i>Journal of Invasive Cardiology</i> , 2006 , 18, 505-8 | 0.7 | 14 |
| 2 | Early regeneration of thymic progenitors in rhesus macaques infected with simian immunodeficiency virus. <i>Journal of Experimental Medicine</i> , 1998 , 187, 1767-78 | 16.6 | 79 |
| 1 | Evaluation of anti-human antibodies for immunohistochemistry on archival nonhuman primate tissues. <i>Journal of Medical Primatology</i> , 1996 , 25, 71-7 | 0.7 | 7 |