Trine Fink

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/1234297/trine-fink-publications-by-year.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

64 1,962 30 43 g-index

69 2,143 4.3 4.44 ext. papers ext. citations avg, IF L-index

| # | Paper | IF | Citations |
|----|--|------|-----------|
| 64 | Multiplex Analysis of Adipose-Derived Stem Cell (ASC) Immunophenotype Adaption to In Vitro Expansion. <i>Cells</i> , 2021 , 10, | 7.9 | 2 |
| 63 | Evolution of ASC Immunophenotypical Subsets During Expansion In Vitro. <i>International Journal of Molecular Sciences</i> , 2020 , 21, | 6.3 | 10 |
| 62 | Distribution of Stromal Cell Subsets in Cultures from Distinct Ocular Surface Compartments. Journal of Ophthalmic and Vision Research, 2020 , 15, 493-501 | 1.2 | |
| 61 | Distribution of Stromal Cell Subsets in Cultures from Distinct Ocular Surface Compartments. Journal of Ophthalmic and Vision Research, 2020 , 15, 493-501 | 1.2 | O |
| 60 | Systematic Review of Stem-Cell-Based Therapy of Burn Wounds: Lessons Learned from Animal and Clinical Studies. <i>Cells</i> , 2020 , 9, | 7.9 | 6 |
| 59 | Fabrication and characterization of extracellular matrix scaffolds obtained from adipose-derived stem cells. <i>Methods</i> , 2020 , 171, 68-76 | 4.6 | 10 |
| 58 | Enhanced Cellular Uptake Of Phenamil Through Inclusion Complex With Histidine Functionalized ECyclodextrin As Penetrative Osteoinductive Agent. <i>International Journal of Nanomedicine</i> , 2019 , 14, 8221-8234 | 7-3 | 6 |
| 57 | Focal Adhesion Kinase Activation Is Necessary for Stretch-Induced Alignment and Enhanced Differentiation of Myogenic Precursor Cells. <i>Tissue Engineering - Part A</i> , 2018 , 24, 631-640 | 3.9 | 8 |
| 56 | Hypoxia Enhances Differentiation of Adipose Tissue-Derived Stem Cells toward the Smooth Muscle Phenotype. <i>International Journal of Molecular Sciences</i> , 2018 , 19, | 6.3 | 7 |
| 55 | Pigmentation Is Associated with Stemness Hierarchy of Progenitor Cells Within Cultured Limbal Epithelial Cells. <i>Stem Cells</i> , 2018 , 36, 1411-1420 | 5.8 | 13 |
| 54 | Hypoxia enhances the wound-healing potential of adipose-derived stem cells in a novel human primary keratinocyte-based scratch assay. <i>International Journal of Molecular Medicine</i> , 2017 , 39, 587-594 | 14.4 | 18 |
| 53 | Maintaining RNA Integrity for Transcriptomic Profiling of Ex Vivo Cultured Limbal Epithelial Stem Cells after Fluorescence-Activated Cell Sorting (FACS). <i>Biological Procedures Online</i> , 2017 , 19, 15 | 8.3 | 2 |
| 52 | Implications of Extracellular Matrix Production by Adipose Tissue-Derived Stem Cells for Development of Wound Healing Therapies. <i>International Journal of Molecular Sciences</i> , 2017 , 18, | 6.3 | 40 |
| 51 | Discrete adipose-derived stem cell subpopulations may display differential functionality after in vitro expansion despite convergence to a common phenotype distribution. <i>Stem Cell Research and Therapy</i> , 2016 , 7, 177 | 8.3 | 16 |
| 50 | Mass spectrometry analysis of adipose-derived stem cells reveals a significant effect of hypoxia on pathways regulating extracellular matrix. <i>Stem Cell Research and Therapy</i> , 2016 , 7, 52 | 8.3 | 33 |
| 49 | Environmental Chemicals Modulate Polar Bear (Ursus maritimus) Peroxisome Proliferator-Activated Receptor Gamma (PPARG) and Adipogenesis in Vitro. <i>Environmental Science & Environmental Science</i> | 10.3 | 33 |
| 48 | Comparative Analysis of Media and Supplements on Initiation and Expansion of Adipose-Derived Stem Cells. <i>Stem Cells Translational Medicine</i> , 2016 , 5, 314-24 | 6.9 | 35 |

(2011-2015)

| 47 | Critical steps in the isolation and expansion of adipose-derived stem cells for translational therapy. <i>Expert Reviews in Molecular Medicine</i> , 2015 , 17, e11 | 6.7 | 32 |
|----|---|-------|----|
| 46 | Uniaxial cyclic strain enhances adipose-derived stem cell fusion with skeletal myocytes. <i>Biochemical and Biophysical Research Communications</i> , 2014 , 450, 1083-8 | 3.4 | 12 |
| 45 | Comparison of human adipose-derived stem cells and bone marrow-derived stem cells in a myocardial infarction model. <i>Cell Transplantation</i> , 2014 , 23, 195-206 | 4 | 59 |
| 44 | Moderate hypoxia influences potassium outward currents in adipose-derived stem cells. <i>PLoS ONE</i> , 2014 , 9, e104912 | 3.7 | 6 |
| 43 | Technical brief: Optimized pipeline for isolation of high-quality RNA from corneal cell subpopulations. <i>Molecular Vision</i> , 2014 , 20, 797-803 | 2.3 | 4 |
| 42 | Hypoxia is a key regulator of limbal epithelial stem cell growth and differentiation. <i>Stem Cell Research</i> , 2013 , 10, 349-60 | 1.6 | 17 |
| 41 | Patterned polymeric surfaces to study the influence of nanotopography on the growth and differentiation of mesenchymal stem cells. <i>Methods in Molecular Biology</i> , 2013 , 1058, 77-88 | 1.4 | 3 |
| 40 | Development of cardiac support bioprostheses for ventricular restoration and myocardial regeneration. <i>European Journal of Cardio-thoracic Surgery</i> , 2013 , 43, 1211-9 | 3 | 29 |
| 39 | Defined xenogeneic-free and hypoxic environment provides superior conditions for long-term expansion of human adipose-derived stem cells. <i>Tissue Engineering - Part C: Methods</i> , 2012 , 18, 593-602 | 2.9 | 35 |
| 38 | Patterned poly(lactic acid) films support growth and spontaneous multilineage gene expression of adipose-derived stem cells. <i>Colloids and Surfaces B: Biointerfaces</i> , 2012 , 93, 92-9 | 6 | 32 |
| 37 | Activation of protease-activated receptor 2 induces VEGF independently of HIF-1. <i>PLoS ONE</i> , 2012 , 7, e46087 | 3.7 | 27 |
| 36 | Prolonged hypoxic culture and trypsinization increase the pro-angiogenic potential of human adipose tissue-derived stem cells. <i>Cytotherapy</i> , 2011 , 13, 318-28 | 4.8 | 82 |
| 35 | Isolation and growth of adipose tissue-derived stem cells. <i>Methods in Molecular Biology</i> , 2011 , 698, 37-4 | 191.4 | 31 |
| 34 | Adipogenic differentiation of human mesenchymal stem cells. <i>Methods in Molecular Biology</i> , 2011 , 698, 243-51 | 1.4 | 53 |
| 33 | Isolation and expansion of adipose-derived stem cells for tissue engineering. <i>Frontiers in Bioscience - Elite</i> , 2011 , 3, 256-63 | 1.6 | 30 |
| 32 | Adipose-derived stem cells from the brown bear (Ursus arctos) spontaneously undergo chondrogenic and osteogenic differentiation in vitro. <i>Stem Cell Research</i> , 2011 , 7, 89-95 | 1.6 | 40 |
| 31 | Hypoxia and adipose-derived stem cell-based tissue regeneration and engineering. <i>Expert Opinion on Biological Therapy</i> , 2011 , 11, 775-86 | 5.4 | 34 |
| 30 | Nanoscale topography reduces fibroblast growth, focal adhesion size and migration-related gene expression on platinum surfaces. <i>Colloids and Surfaces B: Biointerfaces</i> , 2011 , 85, 189-97 | 6 | 55 |

| 29 | Growth of Human Embryonic Stem Cells in Long-Term Hypoxia. Springer Protocols, 2011, 13-24 | 0.3 | |
|----|---|-----|-----|
| 28 | The effect of human embryonic stem cells (hESCs) long-term normoxic and hypoxic cultures on the maintenance of pluripotency. <i>In Vitro Cellular and Developmental Biology - Animal</i> , 2010 , 46, 276-83 | 2.6 | 30 |
| 27 | Notch and hedgehog signaling cooperate to maintain self-renewal of human embryonic stem cells exposed to low oxygen concentration. <i>International Journal of Stem Cells</i> , 2010 , 3, 129-37 | 3 | 6 |
| 26 | Effect of oxygen concentration, culture format and donor variability on in vitro chondrogenesis of human adipose tissue-derived stem cells. <i>Regenerative Medicine</i> , 2009 , 4, 539-48 | 2.5 | 32 |
| 25 | Transcriptional signature of human adipose tissue-derived stem cells (hASCs) preconditioned for chondrogenesis in hypoxic conditions. <i>Experimental Cell Research</i> , 2009 , 315, 1937-52 | 4.2 | 37 |
| 24 | Continuous hypoxic culturing maintains activation of Notch and allows long-term propagation of human embryonic stem cells without spontaneous differentiation. <i>Cell Proliferation</i> , 2009 , 42, 63-74 | 7.9 | 91 |
| 23 | Effect of growth media and serum replacements on the proliferation and differentiation of adipose-derived stem cells. <i>Cytotherapy</i> , 2009 , 11, 189-97 | 4.8 | 74 |
| 22 | Comparative analysis of highly defined proteases for the isolation of adipose tissue-derived stem cells. <i>Regenerative Medicine</i> , 2008 , 3, 705-15 | 2.5 | 40 |
| 21 | Instability of standard PCR reference genes in adipose-derived stem cells during propagation, differentiation and hypoxic exposure. <i>BMC Molecular Biology</i> , 2008 , 9, 98 | 4.5 | 104 |
| 20 | Temporal transcriptome of mouse ATDC5 chondroprogenitors differentiating under hypoxic conditions. <i>Experimental Cell Research</i> , 2006 , 312, 1727-44 | 4.2 | 20 |
| 19 | Optimized chondrogenesis of ATCD5 cells through sequential regulation of oxygen conditions. <i>Tissue Engineering</i> , 2006 , 12, 559-67 | | 20 |
| 18 | Hypoxic treatment inhibits insulin-induced chondrogenesis of ATDC5 cells despite upregulation of DEC1. <i>Connective Tissue Research</i> , 2006 , 47, 119-23 | 3.3 | 7 |
| 17 | Derivation and characterization of four new human embryonic stem cell lines: the Danish experience. <i>Reproductive BioMedicine Online</i> , 2006 , 12, 119-26 | 4 | 21 |
| 16 | Inhibition of osteoblast differentiation but not adipocyte differentiation of mesenchymal stem cells by sera obtained from aged females. <i>Bone</i> , 2006 , 39, 181-8 | 4.7 | 112 |
| 15 | Evaluation of transfection protocols for unmodified and modified peptide nucleic acid (PNA) oligomers. <i>Oligonucleotides</i> , 2006 , 16, 43-57 | | 38 |
| 14 | Separation and detection of compounds in Honeysuckle by integration of ion-exchange chromatography fractionation with reversed-phase liquid chromatography-atmospheric pressure chemical ionization mass spectrometer and matrix-assisted laser desorption/ionization | 3.5 | 36 |
| 13 | Optimized Chondrogenesis of ATCD5 Cells Through Sequential Regulation of Oxygen Conditions. <i>Tissue Engineering</i> , 2006 , 060317120837001 | | |
| 12 | Quantitative transcriptional profiling of ATDC5 mouse progenitor cells during chondrogenesis. <i>Differentiation</i> , 2005 , 73, 350-63 | 3.5 | 36 |

LIST OF PUBLICATIONS

| 11 | factor-1 via MAP kinases and hypoxia-inducible factor-1 in HepG2 cells. <i>Thrombosis and Haemostasis</i> , 2005 , 93, 1176-84 | 7 | 33 |
|----|--|------|-----|
| 10 | Induction of adipocyte-like phenotype in human mesenchymal stem cells by hypoxia. <i>Stem Cells</i> , 2004 , 22, 1346-55 | 5.8 | 145 |
| 9 | Expression of butyrate response factor 1 in HTLV-1-transformed cells and its transactivation by tax protein. <i>Archives of Virology</i> , 2003 , 148, 1787-804 | 2.6 | 3 |
| 8 | Natural killer cell-mediated basal and interferon-enhanced cytotoxicity against liver cancer cells is significantly impaired under in vivo oxygen conditions. <i>Scandinavian Journal of Immunology</i> , 2003 , 58, 607-12 | 3.4 | 41 |
| 7 | Transcriptional activation of immediate-early gene ETR101 by human T-cell leukaemia virus type I Tax. <i>Journal of General Virology</i> , 2003 , 84, 3203-3214 | 4.9 | 7 |
| 6 | Identification of a tightly regulated hypoxia-response element in the promoter of human plasminogen activator inhibitor-1. <i>Blood</i> , 2002 , 99, 2077-83 | 2.2 | 123 |
| 5 | Role of placental cytokines in transcriptional modulation of HIV type 1 in the isolated villous trophoblast. <i>AIDS Research and Human Retroviruses</i> , 2002 , 18, 839-47 | 1.6 | 24 |
| 4 | Biological characterization of three novel variants of IFN-alpha 13 produced by human placental trophoblast. <i>Placenta</i> , 2001 , 22, 673-80 | 3.4 | 5 |
| 3 | Quantitative gene expression profiles of human liver-derived cell lines exposed to moderate hypoxia. <i>Cellular Physiology and Biochemistry</i> , 2001 , 11, 105-14 | 3.9 | 23 |
| 2 | Lack of protection against vertical transmission of HIV-1 by interferons produced during pregnancy in a cohort from East African republic of Malawi. <i>Journal of Medical Virology</i> , 2000 , 61, 195-200 | 19.7 | 3 |
| 1 | Genetic analysis reveals ongoing HIV type 1 evolution in infected human placental trophoblast. <i>AIDS Research and Human Retroviruses</i> , 1999 , 15, 1673-83 | 1.6 | 29 |