Claudia Forjaz

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/1237288/claudia-forjaz-publications-by-year.pdf

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

 143
 2,619
 28
 45

 papers
 citations
 h-index
 g-index

 158
 3,202
 2.4
 4.81

 ext. papers
 ext. citations
 avg, IF
 L-index

#	Paper	IF	Citations
143	Consistency of hemodynamic and autonomic mechanisms underlying post-exercise hypotension. Journal of Human Hypertension, 2021 , 35, 1003-1011	2.6	
142	Activation of Mechanoreflex, but not Central Command, Delays Heart Rate Recovery after Exercise in Healthy Men. <i>International Journal of Sports Medicine</i> , 2021 , 42, 602-609	3.6	0
141	Brazilian Guidelines of Hypertension - 2020. <i>Arquivos Brasileiros De Cardiologia</i> , 2021 , 116, 516-658	1.2	53
140	Comparison of morning versus evening aerobic-exercise training on heart rate recovery in treated hypertensive men: a randomized controlled trial. <i>Blood Pressure Monitoring</i> , 2021 , 26, 388-392	1.3	1
139	Walking Training Improves Ambulatory Blood Pressure Variability in Claudication. <i>Arquivos Brasileiros De Cardiologia</i> , 2021 , 116, 898-905	1.2	O
138	Walking Training Improves Systemic and Local Pathophysiological Processes in Intermittent Claudication. <i>European Journal of Vascular and Endovascular Surgery</i> , 2021 , 61, 954-963	2.3	3
137	Potential Mechanisms Behind the Blood Pressure-Lowering Effect of Dynamic Resistance Training. <i>Current Hypertension Reports</i> , 2021 , 23, 35	4.7	О
136	Effects of resistance training on metabolic and cardiovascular responses to a maximal cardiopulmonary exercise test in Parkinson disease. <i>Einstein (Sao Paulo, Brazil)</i> , 2021 , 19, eAO5940	1.2	O
135	Cardiovascular Responses during and after Maximal Walking in Men and Women with Symptomatic Peripheral Artery Disease. <i>Annals of Vascular Surgery</i> , 2021 , 71, 9-18	1.7	
134	Effects of a Real-Life Park-Based Physical Activity Interventional Program on Cardiovascular Risk and Physical Fitness. <i>Preventing Chronic Disease</i> , 2021 , 18, E18	3.7	2
133	Physiological Responses to Maximal and Submaximal Walking in Patients with Symptomatic Peripheral Artery Disease. <i>Arquivos Brasileiros De Cardiologia</i> , 2021 , 117, 309-316	1.2	O
132	Local and Systemic Inflammation and Oxidative Stress After a Single Bout of Maximal Walking in Patients With Symptomatic Peripheral Artery Disease. <i>Journal of Cardiovascular Nursing</i> , 2021 , 36, 498-	5 6 6	2
131	The influence of a supervised group exercise intervention combined with active lifestyle recommendations on breast cancer survivorsShealth, physical functioning, and quality of life indices: study protocol for a randomized and controlled trial <i>Trials</i> , 2021 , 22, 934	2.8	O
130	Physiological responses during active video games in spinal cord injury: a preliminary study. <i>Physiotherapy Theory and Practice</i> , 2020 , 1-8	1.5	
129	Blood Pressure Increase in Hypertensive Individuals During Resistance Training Protocols With Equated Work to Rest Ratio. <i>Frontiers in Physiology</i> , 2020 , 11, 481	4.6	O
128	Effects of ACEi and ARB on post-exercise hypotension induced by exercises conducted at different times of day in hypertensive men. <i>Clinical and Experimental Hypertension</i> , 2020 , 42, 722-727	2.2	6
127	Post-exercise hypotension and its hemodynamic determinants depend on the calculation approach. <i>Journal of Human Hypertension</i> , 2020 , 34, 719-726	2.6	5

(2018-2020)

126	Reliability and validity of a sedentary behavior questionnaire for South American pediatric population: SAYCARE study. <i>BMC Medical Research Methodology</i> , 2020 , 20, 5	4.7	6
125	Hydration Does Not Change Postexercise Hypotension and Its Mechanisms. <i>Journal of Physical Activity and Health</i> , 2020 , 17, 533-539	2.5	1
124	Effects of postexercise cooling on heart rate recovery in normotensive and hypertensive men. <i>Clinical Physiology and Functional Imaging</i> , 2020 , 40, 114-121	2.4	1
123	Poor sleep quality is associated with cardiac autonomic dysfunction in treated hypertensive men. <i>Journal of Clinical Hypertension</i> , 2020 , 22, 1484-1490	2.3	4
122	Reproducibility of post-exercise heart rate recovery indices: A systematic review. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2019 , 221, 102582	2.4	5
121	Recommendations in Post-exercise Hypotension: Concerns, Best Practices and Interpretation. <i>International Journal of Sports Medicine</i> , 2019 , 40, 487-497	3.6	28
120	Effect of Time of Day on Sustained Postexercise Vasodilation Following Small Muscle-Mass Exercise in Humans. <i>Frontiers in Physiology</i> , 2019 , 10, 762	4.6	3
119	Familial aggregation and heritability of markers of metabolic risk, physical activity, and physical fitness in nuclear families from Muzambinho (Minas Gerais, Brazil). <i>Archives of Endocrinology and Metabolism</i> , 2019 , 63, 215-221	2.2	1
118	Morning versus Evening Aerobic Training Effects on Blood Pressure in Treated Hypertension. <i>Medicine and Science in Sports and Exercise</i> , 2019 , 51, 653-662	1.2	19
117	Reproducibility of Hemodynamic, Cardiac Autonomic Modulation, and Blood Flow Assessments in Patients with Intermittent Claudication. <i>Annals of Vascular Surgery</i> , 2019 , 57, 144-151	1.7	
116	Blood Pressure Response During Resistance Training of Different Work-to-Rest Ratio. <i>Journal of Strength and Conditioning Research</i> , 2019 , 33, 399-407	3.2	5
115	Reliability and Validity of a Questionnaire for Physical Activity Assessment in South American Children and Adolescents: The SAYCARE Study. <i>Obesity</i> , 2018 , 26 Suppl 1, S23-S30	8	9
114	Design and Objectives of the South American Youth/Child Cardiovascular and Environmental (SAYCARE) Study. <i>Obesity</i> , 2018 , 26 Suppl 1, S5-S13	8	13
113	Is the Measurement of Blood Pressure by Automatic Monitor in the South American Pediatric Population Accurate? SAYCARE Study. <i>Obesity</i> , 2018 , 26 Suppl 1, S41-S46	8	3
112	Assessment of physical activity intensity and duration in the paediatric population: evidence to support an a priori hypothesis and sample size in the agreement between subjective and objective methods. <i>Obesity Reviews</i> , 2018 , 19, 810-824	10.6	12
111	Balance and fear of falling in subjects with Parkinson's disease is improved after exercises with motor complexity. <i>Gait and Posture</i> , 2018 , 61, 90-97	2.6	29
110	Aerobic Stimulus Induced by Virtual Reality Games in Stroke Survivors. <i>Archives of Physical Medicine and Rehabilitation</i> , 2018 , 99, 927-933	2.8	6
109	Acute effects of walking and combined exercise on oxidative stress and vascular function in peripheral artery disease. <i>Clinical Physiology and Functional Imaging</i> , 2018 , 38, 69-75	2.4	8

108	Reproducibility of heart rate recovery in patients with intermittent claudication. <i>Clinical Physiology and Functional Imaging</i> , 2018 , 38, 603-609	2.4	4
107	RELA^ [] D ENTRE ATIVIDADE F^ BICA, APTID^ D F^ BICA E RISCO CARDIOVASCULAR: ESTUDO EM MUZAMBINHO, MINAS GERAIS. <i>Revista Brasileira De Medicina Do Esporte</i> , 2018 , 24, 73-77	0.5	1
106	Separate aftereffects of morning and evening exercise on ambulatory blood pressure in prehypertensive men. <i>Journal of Sports Medicine and Physical Fitness</i> , 2018 , 58, 157-163	1.4	O
105	Postexercise hypotension as a clinical tool: a "single brick" in the wall. <i>Journal of the American Society of Hypertension</i> , 2018 , 12, e59-e64		39
104	Reproducibility of ambulatory blood pressure after high-intensity interval training sessions in healthy individuals. <i>Blood Pressure Monitoring</i> , 2018 , 23, 301-304	1.3	2
103	Cardiovascular Responses During Resistance Exercise in Patients With Parkinson Disease. <i>PM and R</i> , 2018 , 10, 1145-1152	2.2	11
102	The Oporto mixed-longitudinal growth, health and performance study. Design, methods and baseline results. <i>Annals of Human Biology</i> , 2017 , 44, 11-20	1.7	2
101	Patients with Parkinson disease present high ambulatory blood pressure variability. <i>Clinical Physiology and Functional Imaging</i> , 2017 , 37, 530-535	2.4	14
100	Resistance training with instability is more effective than resistance training in improving spinal inhibitory mechanisms in Parkinson's disease. <i>Journal of Applied Physiology</i> , 2017 , 122, 1-10	3.7	14
99	Captopril does not Potentiate Post-Exercise Hypotension: A Randomized Crossover Study. <i>International Journal of Sports Medicine</i> , 2017 , 38, 270-277	3.6	8
98	A Single Session of Low-Volume High-Intensity Interval Exercise Reduces Ambulatory Blood Pressure in Normotensive Men. <i>Journal of Strength and Conditioning Research</i> , 2017 , 31, 2263-2269	3.2	9
97	Relationship between walking capacity and ambulatory blood pressure in patients with intermittent claudication. <i>Blood Pressure Monitoring</i> , 2017 , 22, 115-121	1.3	13
96	Instability Resistance Training Improves Neuromuscular Outcome in Parkinson's Disease. <i>Medicine and Science in Sports and Exercise</i> , 2017 , 49, 652-660	1.2	14
95	Walking training at the heart rate of pain threshold improves cardiovascular function and autonomic regulation in intermittent claudication: A randomized controlled trial. <i>Journal of Science and Medicine in Sport</i> , 2017 , 20, 886-892	4.4	28
94	Additive effects of heating and exercise on baroreflex control of heart rate in healthy males. <i>Journal of Applied Physiology</i> , 2017 , 123, 1555-1562	3.7	6
93	Reproducibility (Reliability and Agreement) of Post-exercise Hypotension. <i>International Journal of Sports Medicine</i> , 2017 , 38, 1029-1034	3.6	7
92	Effects of Progressive Resistance Training on Cardiovascular Autonomic Regulation in Patients With Parkinson Disease: A Randomized Controlled Trial. <i>Archives of Physical Medicine and Rehabilitation</i> , 2017 , 98, 2134-2141	2.8	21
91	Blood pressure reactivity to mental stress is attenuated following resistance exercise in older hypertensive women. <i>Clinical Interventions in Aging</i> , 2017 , 12, 793-803	4	4

(2015-2017)

90	Passive Heating Attenuates Post-exercise Cardiac Autonomic Recovery in Healthy Young Males. <i>Frontiers in Neuroscience</i> , 2017 , 11, 727	5.1	10	
89	Relationship between physical activity, physical fitness and multiple metabolic risk in youths from Muzambinhos study. <i>European Journal of Sport Science</i> , 2016 , 16, 618-23	3.9	3	
88	Metaboreflex activation delays heart rate recovery after aerobic exercise in never-treated hypertensive men. <i>Journal of Physiology</i> , 2016 , 594, 6211-6223	3.9	20	
87	Post-Exercise Hypotension Is Mediated by a Decrease in Sympathetic Nerve Activity in Stages 2-3 CKD. <i>American Journal of Nephrology</i> , 2016 , 43, 206-12	4.6	10	
86	Blunted Maximal and Submaximal Responses to Cardiopulmonary Exercise Tests in Patients With Parkinson Disease. <i>Archives of Physical Medicine and Rehabilitation</i> , 2016 , 97, 720-5	2.8	22	
85	A methodological approach to short-term tracking of youth physical fitness: the Oporto Growth, Health and Performance Study. <i>Journal of Sports Sciences</i> , 2016 , 34, 1885-92	3.6	3	
84	Reproducibility of Anaerobic and Pain Thresholds in Male Patients With Intermittent Claudication. Journal of Cardiopulmonary Rehabilitation and Prevention, 2016 , 36, 358-67	3.6	3	
83	Resistance Training with Instability for Patients with Parkinson's Disease. <i>Medicine and Science in Sports and Exercise</i> , 2016 , 48, 1678-87	1.2	52	
82	Modeling longitudinal changes in hypertensive and waist phenotype: The oporto growth, health, and performance study. <i>American Journal of Human Biology</i> , 2016 , 28, 387-93	2.7		
81	Modeling the dynamics of BMI changes during adolescence. The Oporto Growth, Health and Performance Study. <i>International Journal of Obesity</i> , 2015 , 39, 1063-9	5.5	9	
80	Influence of endurance and resistance exercise order on the postexercise hemodynamic responses in hypertensive women. <i>Journal of Strength and Conditioning Research</i> , 2015 , 29, 612-8	3.2	10	
79	Post-walking exercise hypotension in patients with intermittent claudication. <i>Medicine and Science in Sports and Exercise</i> , 2015 , 47, 460-7	1.2	19	
78	Time of day affects heart rate recovery and variability after maximal exercise in pre-hypertensive men. <i>Chronobiology International</i> , 2015 , 32, 1385-90	3.6	7	
77	A session of resistance exercise increases vasodilation in intermittent claudication patients. <i>Applied Physiology, Nutrition and Metabolism</i> , 2015 , 40, 59-64	3	6	
76	Barriers to physical activity in patients with intermittent claudication. <i>International Journal of Behavioral Medicine</i> , 2015 , 22, 70-6	2.6	45	
75	Can a first-order exponential decay model fit heart rate recovery after resistance exercise?. <i>Clinical Physiology and Functional Imaging</i> , 2015 , 35, 98-103	2.4	13	
74	Amlodipine reduces blood pressure during dynamic resistance exercise in hypertensive patients. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2015 , 25, 53-60	4.6	7	
73	Are the barriers for physical activity practice equal for all peripheral artery disease patients?. <i>Archives of Physical Medicine and Rehabilitation</i> , 2015 , 96, 248-52	2.8	31	

72	Post-resistance exercise hemodynamic and autonomic responses: Comparison between normotensive and hypertensive men. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2015 , 25, 486-94	4.6	34
71	High-Intensity Progressive Resistance Training Increases Strength With No Change in Cardiovascular Function and Autonomic Neural Regulation in Older Adults. <i>Journal of Aging and Physical Activity</i> , 2015 , 23, 339-45	1.6	16
70	High-Intensity Resistance Exercise Promotes Postexercise Hypotension Greater than Moderate Intensity and Affects Cardiac Autonomic Responses in Women Who Are Hypertensive. <i>Journal of Strength and Conditioning Research</i> , 2015 , 29, 3486-93	3.2	16
69	Post-Exercise Hypotension and Its Mechanisms Differ after Morning and Evening Exercise: A Randomized Crossover Study. <i>PLoS ONE</i> , 2015 , 10, e0132458	3.7	39
68	Effects of oral N-acetylcysteine on walking capacity, leg reactive hyperemia, and inflammatory and angiogenic mediators in patients with intermittent claudication. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2015 , 309, H897-905	5.2	17
67	Modela [^] [] B multin [^] [Jel e delineamento longitudinal-misto na pesquisa em Educa [^] [] B F [^] [Sica e Ci [^] [Bcias do Esporte. <i>Revista Brasileira De Educa</i> [B F] <i>F</i> [Sica E Esporte: RBEFE, 2015 , 29, 35-45]	0.8	
66	Metaboreflex Activation Delays Heart Rate Recovery after Aerobic Exercise. <i>FASEB Journal</i> , 2015 , 29, 1054.4	0.9	1
65	Resistance Training with Instability Increase Levels of Spinal Inhibition and Decrease the Motor Symptoms of Parkinsonians. <i>FASEB Journal</i> , 2015 , 29, 677.15	0.9	
64	Heart rate recovery: autonomic determinants, methods of assessment and association with mortality and cardiovascular diseases. <i>Clinical Physiology and Functional Imaging</i> , 2014 , 34, 327-39	2.4	110
63	Influence of population and exercise protocol characteristics on hemodynamic determinants of post-aerobic exercise hypotension. <i>Brazilian Journal of Medical and Biological Research</i> , 2014 , 47, 626-36	6 ^{2.8}	57
62	Effects of estrogen therapy and aerobic training on sympathetic activity and hemodynamics in healthy postmenopausal women: a double-blind randomized trial. <i>Menopause</i> , 2014 , 21, 369-75	2.5	10
61	Oral estrogen therapy may mitigate the effects of aerobic training on cardiorespiratory fitness in postmenopausal women: a double-blind, randomized clinical pilot study. <i>Menopause</i> , 2014 , 21, 376-82	2.5	
60	A single bout of resistance exercise does not modify cardiovascular responses during daily activities in patients with peripheral artery disease. <i>Blood Pressure Monitoring</i> , 2014 , 19, 64-71	1.3	14
59	Yoga Relaxation (savasana) decreases cardiac sympathovagal balance in hypertensive patients. <i>Medical Express</i> , 2014 , 1,		4
58	Gender influence on post-resistance exercise hypotension and hemodynamics. <i>International Journal of Sports Medicine</i> , 2013 , 34, 939-44	3.6	26
57	Cardiac work remains high after strength exercise in elderly. <i>International Journal of Sports Medicine</i> , 2013 , 34, 391-7	3.6	17
56	Cardiovascular adaptations to resistance training in elderly postmenopausal women. <i>International Journal of Sports Medicine</i> , 2013 , 34, 806-13	3.6	29
55	Relacao entre o nivel de atividade fisica estimado pelo Baltimore Activity Scale for Intermittent Claudication e a pedometria em pacientes com claudicacao intermitente. <i>Jornal Vascular Brasileiro</i> , 2013, 12, 187-192	0.9	3

(2011-2013)

54	A bout of resistance exercise following the 2007 AHA guidelines decreases asleep blood pressure in Mozambican men. <i>Journal of Strength and Conditioning Research</i> , 2013 , 27, 786-92	3.2	4
53	Prescri [^] [] b de caminhada n [^] b supervisionada, risco cardiovascular e aptid [^] b f [^] sica. <i>Revista Brasileira De Educa F Bica E Esporte: RBEFE</i> , 2013 , 27, 377-386	0.8	2
52	Low-intensity resistance exercise does not affect cardiac autonomic modulation in patients with peripheral artery disease. <i>Clinics</i> , 2013 , 68, 632-7	2.3	8
51	Effect of rest interval on cardiovascular responses after resistance exercise. <i>Motriz Revista De Educacao Fisica</i> , 2013 , 19, 252-260	0.9	1
50	Betabloqueio com atenolol n° B reduz pot° Bicia aer° Bia nem muda limiares ventilat° Bios em hipertensos sedent° Bios. <i>Revista Brasileira De Medicina Do Esporte</i> , 2013 , 19, 339-342	0.5	
49	Fisiologia do Exerc [^] dio para alunos de gradua [^] d d: uso de estrat [^] gias de ensino baseadas na metodologia dial [^] tica. <i>Revista Brasileira De Educad Fâica E Esporte: RBEFE</i> , 2013 , 27, 289-296	0.8	
48	Exercise prescription using the heart of claudication pain onset in patients with intermittent claudication. <i>Clinics</i> , 2013 , 68, 974-8	2.3	23
47	Predictors of walking capacity in peripheral arterial disease patients. <i>Clinics</i> , 2013 , 68, 537-41	2.3	16
46	Sympathetic cardiac modulation and vascular worsening in arteritis: a case report. <i>Journal of Vascular Nursing</i> , 2012 , 30, 21-3	1	5
45	Isokinetic strength and endurance in proximal and distal muscles in patients with peripheral artery disease. <i>Annals of Vascular Surgery</i> , 2012 , 26, 1114-9	1.7	27
44	Stages of health behavior change and factors associated with physical activity in patients with intermittent claudication. <i>Einstein (Sao Paulo, Brazil)</i> , 2012 , 10, 422-7	1.2	7
43	Genetic and environmental influences on blood pressure and physical activity: a study of nuclear families from Muzambinho, Brazil. <i>Brazilian Journal of Medical and Biological Research</i> , 2012 , 45, 1269-7	5 ^{2.8}	7
42	Aerobic training abolishes ambulatory blood pressure increase induced by estrogen therapy: a double blind randomized clinical trial. <i>Maturitas</i> , 2011 , 69, 189-94	5	10
41	Post exercise cardiovascular effects of different resistance exercise protocols for trunk and upper limbs. <i>Motriz Revista De Educacao Fisica</i> , 2011 , 17, 667-674	0.9	2
40	A fisiologia em educa [^] [] [5] f [^] [5] lica e esporte. <i>Revista Brasileira De Educa</i> [6] <i>F</i> [6] <i>ica E Esporte: RBEFE</i> , 2011 , 25, 7-13	0.8	1
39	Acute effect of resistance exercise intensity in cardiac autonomic modulation after exercise. <i>Arquivos Brasileiros De Cardiologia</i> , 2011 , 96, 498-503	1.2	40
38	Post-resistance exercise hypotension in patients with intermittent claudication. <i>Clinics</i> , 2011 , 66, 221-6	2.3	17
37	Strength and power training did not modify cardiovascular responses to aerobic exercise in elderly subjects. <i>Brazilian Journal of Medical and Biological Research</i> , 2011 , 44, 864-70	2.8	16

36	Impact of a supervised strength training or walking training over a subsequent unsupervised therapy period on walking capacity in patients with claudication. <i>Journal of Vascular Nursing</i> , 2011 , 29, 81-6	1	13
35	Post-concurrent exercise hemodynamics and cardiac autonomic modulation. <i>European Journal of Applied Physiology</i> , 2011 , 111, 2069-78	3.4	101
34	Rela [^] [] [5] entre aptid [^] [6] f [^] [5] ica e os indicadores de qualidade de vida de indiv [^] [6] uos com claudica [^] [1] [5] intermitente. <i>Revista Brasileira De Medicina Do Esporte</i> , 2011 , 17, 175-178	0.5	7
33	Risco cardiovascular e pr^ Eica de atividade f^ Eica em crian^ Eis e adolescentes de Muzambinho/MG: influ^ Eicia do g^ Eiero e da idade. <i>Revista Brasileira De Medicina Do Esporte</i> , 2011 , 17, 232-236	0.5	6
32	Associa [^] [] [b] de comorbidades e h [^] [b]itos n [^] [b] saud [^] [Q]eis com a capacidade de caminhada em pacientes com claudica [^] [] [b] intermitente. <i>Revista Brasileira De Educa</i> [b] <i>F</i> [s]ica E Esporte: RBEFE, 2011 , 25, 277-284	0.8	0
31	Effects of walking and strength training on resting and exercise cardiovascular responses in patients with intermittent claudication. <i>Vasa - European Journal of Vascular Medicine</i> , 2011 , 40, 390-7	1.9	30
30	Desafios no estudo de fam [^] [las nucleares: etapas iniciais de an [^] [lse. <i>Revista Brasileira De Educa</i>] <i>F</i> @ica E Esporte: RBEFE, 2011 , 25, 717-730	0.8	1
29	Atenolol blunts blood pressure increase during dynamic resistance exercise in hypertensives. <i>British Journal of Clinical Pharmacology</i> , 2010 , 70, 664-73	3.8	12
28	Acute and chronic effects of aerobic and resistance exercise on ambulatory blood pressure. <i>Clinics</i> , 2010 , 65, 317-25	2.3	122
27	Intra-arterial blood pressure response in hypertensive subjects during low- and high-intensity resistance exercise. <i>Clinics</i> , 2010 , 65, 271-7	2.3	53
26	O desafio da informa [^] [] [5] longitudinal: um "passeio guiado" sobre modela [^] [1] [5] hier [^] [5] fiquica, "tracking" e informa [^] [1] [5] omissa com um conjunto de dados do estudo de Muzambinho. <i>Revista Brasileira De Educa</i> [6] <i>F</i> [6] <i>ica E Esporte: RBEFE</i> , 2010 , 24, 413-431	0.8	1
25	Cardiovascular responses during isokinetic muscle assessment in claudicant patients. <i>Arquivos Brasileiros De Cardiologia</i> , 2010 , 95, 571-6	1.2	7
24	Effects of resistance training on blood pressure in the elderly. <i>Arquivos Brasileiros De Cardiologia</i> , 2010 , 95, 135-40	1.2	18
23	Tabelas de classifica [^] [] [5] da aptid [^] [5] f [^] [5] lica para frequentadores de parques p [^] [5] blicos. <i>Revista Brasileira De Medicina Do Esporte</i> , 2010 , 16, 373-377	0.5	2
22	Finger blood pressure during leg resistance exercise. <i>International Journal of Sports Medicine</i> , 2010 , 31, 590-5	3.6	12
21	Strength training increases walking tolerance in intermittent claudication patients: randomized trial. <i>Journal of Vascular Surgery</i> , 2010 , 51, 89-95	3.5	68
20	Translation and validation of the walking impairment questionnaire in Brazilian subjects with intermittent claudication. <i>Arquivos Brasileiros De Cardiologia</i> , 2009 , 92, 136-49	1.2	37
19	Obesity decreases time to claudication and delays post-exercise hemodynamic recovery in elderly peripheral arterial disease patients. <i>Gerontology</i> , 2009 , 55, 21-6	5.5	19

(1993-2009)

18	Clinic and ambulatory blood pressure responses after resistance exercise. <i>Journal of Strength and Conditioning Research</i> , 2009 , 23, 571-8	3.2	39
17	Pain threshold is achieved at intensity above anaerobic threshold in patients with intermittent claudication. <i>Journal of Cardiopulmonary Rehabilitation and Prevention</i> , 2009 , 29, 396-401	3.6	12
16	Crescimento e desenvolvimento motor de escolares de Muzambinho: um estudo com implica [^] []	0	4
15	Low-dose estrogen therapy does not change postexercise hypotension, sympathetic nerve activity reduction, and vasodilation in healthy postmenopausal women. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2008 , 295, H1802-8	5.2	7
14	A randomized, placebo-controlled trial of the effects of physical exercises and estrogen therapy on health-related quality of life in postmenopausal women. <i>Menopause</i> , 2008 , 15, 613-8	2.5	45
13	Neurovascular and hemodynamic responses to hyperinsulinemia in healthy postmenopausal women. <i>Maturitas</i> , 2007 , 58, 50-8	5	3
12	Aftereffects of exercise and relaxation on blood pressure. <i>Clinical Journal of Sport Medicine</i> , 2006 , 16, 341-7	3.2	28
11	Postexercise hypotension induced by low-intensity resistance exercise in hypertensive women receiving captopril. <i>Blood Pressure Monitoring</i> , 2006 , 11, 183-9	1.3	91
10	Post-resistance exercise hypotension, hemodynamics, and heart rate variability: influence of exercise intensity. <i>European Journal of Applied Physiology</i> , 2006 , 98, 105-12	3.4	200
9	Previous exercise attenuates muscle sympathetic activity and increases blood flow during acute euglycemic hyperinsulinemia. <i>Journal of Applied Physiology</i> , 2005 , 98, 866-71	3.7	26
8	Assessment of the cardiovascular risk and physical activity of individuals exercising at a public park in the city of sˆ D̄ paulo. <i>Arquivos Brasileiros De Cardiologia</i> , 2002 , 79, 35-50	1.2	4
7	Muscle metaboreflex control is diminished in normotensive obese women. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2001 , 281, H469-75	5.2	68
6	Muscle sympathetic nerve activity and hemodynamic alterations in middle-aged obese women. Brazilian Journal of Medical and Biological Research, 2001 , 34, 475-8	2.8	36
5	Factors affecting post-exercise hypotension in normotensive and hypertensive humans. <i>Blood Pressure Monitoring</i> , 2000 , 5, 255-62	1.3	55
4	Postexercise responses of muscle sympathetic nerve activity and blood flow to hyperinsulinemia in humans. <i>Journal of Applied Physiology</i> , 1999 , 87, 824-9	3.7	15
3	Post-exercise changes in blood pressure, heart rate and rate pressure product at different exercise intensities in normotensive humans. <i>Brazilian Journal of Medical and Biological Research</i> , 1998 , 31, 1247	7-358	77
2	Oral glucose ingestion increases endurance capacity in normal and diabetic (type I) humans. <i>Journal of Applied Physiology</i> , 1997 , 83, 608-14	3.7	24
1	Exercise tolerance is lower in type I diabetics compared with normal young men. <i>Metabolism:</i> Clinical and Experimental, 1993 , 42, 191-5	12.7	13