

Roland H Wenger

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1241072/roland-h-wenger-publications-by-year.pdf>

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

158 papers	12,940 citations	56 h-index	112 g-index
168 ext. papers	14,070 ext. citations	6.5 avg, IF	6.1 L-index

#	Paper	IF	Citations
158	Fate-mapping of erythropoietin-producing cells in mouse models of hypoxaemia and renal tissue remodelling reveals repeated recruitment and persistent functionality.. <i>Acta Physiologica</i> , 2022 , e13768	5.6	1
157	Sphk1 and Sphk2 Differentially Regulate Erythropoietin Synthesis in Mouse Renal Interstitial Fibroblast-like Cells. <i>International Journal of Molecular Sciences</i> , 2022 , 23, 5882	6.3	1
156	OTUB1 regulates lung development, adult lung tissue homeostasis, and respiratory control. <i>FASEB Journal</i> , 2021 , 35, e22039	0.9	0
155	Heritability and association with distinct genetic loci of erythropoietin levels in the general population. <i>Haematologica</i> , 2021 , 106, 2499-2501	6.6	0
154	Interfering with Tumor Hypoxia for Radiotherapy Optimization. <i>Journal of Experimental and Clinical Cancer Research</i> , 2021 , 40, 197	12.8	14
153	Hypoxia Reduces the Transcription of Fibrotic Markers in the Intestinal Mucosa. <i>Inflammatory Intestinal Diseases</i> , 2021 , 6, 87-100	2.5	
152	S1P Stimulates Erythropoietin Production in Mouse Renal Interstitial Fibroblasts by S1P and S1P Receptor Activation and HIF-2 α Stabilization. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	5
151	Tumor cell endogenous HIF-1 α activity induces aberrant angiogenesis and interacts with TRAF6 pathway required for colorectal cancer development. <i>Neoplasia</i> , 2020 , 22, 745-758	6.4	3
150	Cre-mediated, loxP independent sequential recombination of a tripartite transcriptional stop cassette allows for partial read-through transcription. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2020 , 1863, 194568	6	5
149	Different subpopulations of kidney interstitial cells produce erythropoietin and factors supporting tissue oxygenation in response to hypoxia in vivo. <i>Kidney International</i> , 2020 , 98, 918-931	9.9	17
148	Hypoxia sensing by hepatic stellate cells leads to VEGF-dependent angiogenesis and may contribute to accelerated liver regeneration. <i>Scientific Reports</i> , 2020 , 10, 4392	4.9	13
147	Simultaneous Three-Dimensional Vascular and Tubular Imaging of Whole Mouse Kidneys With X-ray CT. <i>Microscopy and Microanalysis</i> , 2020 , 26, 731-740	0.5	1
146	The Antioxidative Role of Cytoglobin in Podocytes: Implications for a Role in Chronic Kidney Disease. <i>Antioxidants and Redox Signaling</i> , 2020 , 32, 1155-1171	8.4	17
145	HIF hydroxylase inhibitors decrease cellular oxygen consumption depending on their selectivity. <i>FASEB Journal</i> , 2020 , 34, 2344-2358	0.9	12
144	Thin air - thick science. <i>Nature Reviews Molecular Cell Biology</i> , 2020 , 21, 567	48.7	
143	Inhibition of firefly luciferase activity by a HIF prolyl hydroxylase inhibitor. <i>Journal of Photochemistry and Photobiology B: Biology</i> , 2020 , 210, 111980	6.7	5
142	Distal and proximal hypoxia response elements cooperate to regulate organ-specific erythropoietin gene expression. <i>Haematologica</i> , 2020 , 105, 2774-2784	6.6	12

141	The neuronal oxygen-sensing pathway controls postnatal vascularization of the murine brain. <i>FASEB Journal</i> , 2019 , 33, 12812-12824	0.9	4
140	Tumor necrosis factor stimulates fibroblast growth factor 23 levels in chronic kidney disease and non-renal inflammation. <i>Kidney International</i> , 2019 , 96, 890-905	9.9	32
139	Oxygen-dependent bond formation with FIH regulates the activity of the client protein OTUB1. <i>Redox Biology</i> , 2019 , 26, 101265	11.3	6
138	Now a Nobel gas: oxygen. <i>Pflugers Archiv European Journal of Physiology</i> , 2019 , 471, 1343-1358	4.6	16
137	Generation of renal Epo-producing cell lines by conditional gene tagging reveals rapid HIF-2 driven Epo kinetics, cell autonomous feedback regulation, and a telocyte phenotype. <i>Kidney International</i> , 2019 , 95, 375-387	9.9	22
136	Source and microenvironmental regulation of erythropoietin in the kidney. <i>Current Opinion in Nephrology and Hypertension</i> , 2018 , 27, 277-282	3.5	12
135	Erythropoietin stimulates fibroblast growth factor 23 (FGF23) in mice and men. <i>Pflugers Archiv European Journal of Physiology</i> , 2018 , 470, 1569-1582	4.6	50
134	Protocol for a prospective, controlled, observational study to evaluate the influence of hypoxia on healthy volunteers and patients with inflammatory bowel disease: the Altitude IBD Study. <i>BMJ Open</i> , 2017 , 7, e013477	3	6
133	The functional interplay between the HIF pathway and the ubiquitin system - more than a one-way road. <i>Experimental Cell Research</i> , 2017 , 356, 152-159	4.2	14
132	Hypoxia ameliorates intestinal inflammation through NLRP3/mTOR downregulation and autophagy activation. <i>Nature Communications</i> , 2017 , 8, 98	17.4	125
131	Hypoxia of the growing liver accelerates regeneration. <i>Surgery</i> , 2017 , 161, 666-679	3.6	42
130	FIH Regulates Cellular Metabolism through Hydroxylation of the Deubiquitinase OTUB1. <i>PLoS Biology</i> , 2016 , 14, e1002347	9.7	53
129	Estrogen-dependent downregulation of hypoxia-inducible factor (HIF)-2 α in invasive breast cancer cells. <i>Oncotarget</i> , 2016 , 7, 31153-65	3.3	13
128	Erythropoietin production by PDGFR- α cells. <i>Pflugers Archiv European Journal of Physiology</i> , 2016 , 468, 1479-87	4.6	26
127	ITGA6 is directly regulated by hypoxia-inducible factors and enriches for cancer stem cell activity and invasion in metastatic breast cancer models. <i>Molecular Cancer</i> , 2016 , 15, 26	42.1	110
126	Intermittent hypoxia confers pro-metastatic gene expression selectively through NF- κ B in inflammatory breast cancer cells. <i>Free Radical Biology and Medicine</i> , 2016 , 101, 129-142	7.8	33
125	Frequently asked questions in hypoxia research. <i>Hypoxia (Auckland, N Z)</i> , 2015 , 3, 35-43	2.1	122
124	Induction of long noncoding RNA MALAT1 in hypoxic mice. <i>Hypoxia (Auckland, N Z)</i> , 2015 , 3, 45-52	2.1	55

123	Destruction of a distal hypoxia response element abolishes trans-activation of the PAG1 gene mediated by HIF-independent chromatin looping. <i>Nucleic Acids Research</i> , 2015 , 43, 5810-23	20.1	17
122	Hypoxia attenuates the proinflammatory response in colon cancer cells by regulating IB. <i>Oncotarget</i> , 2015 , 6, 20288-301	3.3	16
121	A novel distal upstream hypoxia response element regulating oxygen-dependent erythropoietin gene expression. <i>Haematologica</i> , 2014 , 99, e45-8	6.6	23
120	TET1-mediated hydroxymethylation facilitates hypoxic gene induction in neuroblastoma. <i>Cell Reports</i> , 2014 , 7, 1343-1352	10.6	115
119	Hypoxia-inducible factor-mediated induction of WISP-2 contributes to attenuated progression of breast cancer. <i>Hypoxia (Auckland, N Z)</i> , 2014 , 2, 23-33	2.1	10
118	PAS kinase is a nutrient and energy sensor in hypothalamic areas required for the normal function of AMPK and mTOR/S6K1. <i>Molecular Neurobiology</i> , 2014 , 50, 314-26	6.2	19
117	Neutrophil expression of ICAM1, CXCR1, and VEGFR1 in patients with breast cancer before and after adjuvant chemotherapy. <i>Anticancer Research</i> , 2014 , 34, 4693-9	2.3	5
116	Combined whole-body vibration, resistance exercise, and sustained vascular occlusion increases PGC-1 α and VEGF mRNA abundances. <i>European Journal of Applied Physiology</i> , 2013 , 113, 1081-90	3.4	16
115	Onconeural antigen Cdr2 correlates with HIF prolyl-4-hydroxylase PHD1 and worse prognosis in renal cell carcinoma. <i>Experimental and Molecular Pathology</i> , 2013 , 94, 453-7	4.4	5
114	Dysregulation of hypoxia-inducible factor by presenilin/ β -secretase loss-of-function mutations. <i>Journal of Neuroscience</i> , 2013 , 33, 1915-26	6.6	14
113	Hematopoiesis and the Kidney 2013 , 3087-3124		
112	HIF mediated and DNA damage independent histone H2AX phosphorylation in chronic hypoxia. <i>Biological Chemistry</i> , 2013 , 394, 519-28	4.5	27
111	Hypoxia enhances lipid uptake in macrophages: role of the scavenger receptors Lox1, SRA, and CD36. <i>Atherosclerosis</i> , 2013 , 229, 110-7	3.1	86
110	Breast tumor kinase (Brk/PTK6) is a mediator of hypoxia-associated breast cancer progression. <i>Cancer Research</i> , 2013 , 73, 5810-20	10.1	39
109	Congenital erythrocytosis associated with gain-of-function HIF2A gene mutations and erythropoietin levels in the normal range. <i>Haematologica</i> , 2013 , 98, 1624-32	6.6	21
108	The transcription factor encyclopedia. <i>Genome Biology</i> , 2012 , 13, R24	18.3	86
107	Identification and functional characterization of pVHL-dependent cell surface proteins in renal cell carcinoma. <i>Neoplasia</i> , 2012 , 14, 535-46	6.4	40
106	Non-canonical HIF-2 β function drives autonomous breast cancer cell growth via an AREG-EGFR/ErbB4 autocrine loop. <i>Oncogene</i> , 2012 , 31, 2283-97	9.2	55

105	Synthetic transactivation screening reveals ETV4 as broad coactivator of hypoxia-inducible factor signaling. <i>Nucleic Acids Research</i> , 2012 , 40, 1928-43	20.1	24
104	Distinct deregulation of the hypoxia inducible factor by PHD2 mutants identified in germline DNA of patients with polycythemia. <i>Haematologica</i> , 2012 , 97, 9-14	6.6	40
103	Erythropoietin. <i>Comprehensive Physiology</i> , 2011 , 1, 1759-94	7.7	32
102	The putative RNA helicase HELZ promotes cell proliferation, translation initiation and ribosomal protein S6 phosphorylation. <i>PLoS ONE</i> , 2011 , 6, e22107	3.7	15
101	Vitamin C is dispensable for oxygen sensing in vivo. <i>Blood</i> , 2011 , 117, 5485-93	2.2	56
100	Substrate preference and phosphatidylinositol monophosphate inhibition of the catalytic domain of the Per-Arnt-Sim domain kinase PASKIN. <i>FEBS Journal</i> , 2011 , 278, 1757-68	5.7	10
99	Prolyl-4-hydroxylase PHD2- and hypoxia-inducible factor 2-dependent regulation of amphiregulin contributes to breast tumorigenesis. <i>Oncogene</i> , 2011 , 30, 548-60	9.2	58
98	A graphical simulation software for instruction in cardiovascular mechanics physiology. <i>BioMedical Engineering OnLine</i> , 2011 , 10, 8	4.1	6
97	IRadiation promotes immunological recognition of cancer cells through increased expression of cancer-testis antigens in vitro and in vivo. <i>PLoS ONE</i> , 2011 , 6, e28217	3.7	108
96	Regulated oxygen sensing by protein hydroxylation in renal erythropoietin-producing cells. <i>American Journal of Physiology - Renal Physiology</i> , 2010 , 298, F1287-96	4.3	70
95	Longitudinal and multimodal in vivo imaging of tumor hypoxia and its downstream molecular events. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009 , 106, 14004-9	11.5	54
94	Inhibitor of DNA binding/differentiation 2 induced by hypoxia promotes synovial fibroblast-dependent osteoclastogenesis. <i>Arthritis and Rheumatism</i> , 2009 , 60, 3663-75		13
93	The PAS-domain kinase PASKIN: a new sensor in energy homeostasis. <i>Cellular and Molecular Life Sciences</i> , 2009 , 66, 876-83	10.3	23
92	Taking advantage of tumor cell adaptations to hypoxia for developing new tumor markers and treatment strategies. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2009 , 24 Suppl 1, 1-39	5.6	153
91	Hypoxia-inducible factor prolyl-4-hydroxylase PHD2 protein abundance depends on integral membrane anchoring of FKBP38. <i>Journal of Biological Chemistry</i> , 2009 , 284, 23046-58	5.4	56
90	HIF prolyl-4-hydroxylase interacting proteins: consequences for drug targeting. <i>Current Pharmaceutical Design</i> , 2009 , 15, 3886-94	3.3	18
89	Impaired DNA double-strand break repair contributes to chemoresistance in HIF-1 alpha-deficient mouse embryonic fibroblasts. <i>Carcinogenesis</i> , 2008 , 29, 2306-16	4.6	36
88	Induction of activating transcription factor 3 by anoxia is independent of p53 and the hypoxic HIF signalling pathway. <i>Oncogene</i> , 2007 , 26, 284-9	9.2	63

87	Lack of hypoxia-inducible factor-1 alpha impairs midbrain neural precursor cells involving vascular endothelial growth factor signaling. <i>Journal of Neuroscience</i> , 2007 , 27, 412-21	6.6	92
86	The peptidyl prolyl cis/trans isomerase FKBP38 determines hypoxia-inducible transcription factor prolyl-4-hydroxylase PHD2 protein stability. <i>Molecular and Cellular Biology</i> , 2007 , 27, 3758-68	4.8	90
85	Male germ cell expression of the PAS domain kinase PASKIN and its novel target eukaryotic translation elongation factor eEF1A1. <i>Cellular Physiology and Biochemistry</i> , 2007 , 20, 227-40	3.9	22
84	Glucose-stimulated insulin production in mice deficient for the PAS kinase PASKIN. <i>Diabetes</i> , 2007 , 56, 113-7	0.9	16
83	Oxygen-dependent ATF-4 stability is mediated by the PHD3 oxygen sensor. <i>Blood</i> , 2007 , 110, 3610-7	2.2	161
82	Time course of hypoxia-induced lung injury in rats. <i>Respiratory Physiology and Neurobiology</i> , 2007 , 159, 45-54	2.8	17
81	Determination and modulation of prolyl-4-hydroxylase domain oxygen sensor activity. <i>Methods in Enzymology</i> , 2007 , 435, 43-60	1.7	16
80	Regulated function of the prolyl-4-hydroxylase domain (PHD) oxygen sensor proteins. <i>Antioxidants and Redox Signaling</i> , 2007 , 9, 1329-38	8.4	28
79	Cutting edge: hypoxia-inducible factor 1alpha and its activation-inducible short isoform I.1 negatively regulate functions of CD4+ and CD8+ T lymphocytes. <i>Journal of Immunology</i> , 2006 , 177, 4962-5	5.3	166
78	Increased prolyl 4-hydroxylase domain proteins compensate for decreased oxygen levels. Evidence for an autoregulatory oxygen-sensing system. <i>Journal of Biological Chemistry</i> , 2006 , 281, 23482-91	5.4	213
77	TSGA10 prevents nuclear localization of the hypoxia-inducible factor (HIF)-1alpha. <i>FEBS Letters</i> , 2006 , 580, 3731-8	3.8	27
76	Mitochondria: oxygen sinks rather than sensors?. <i>Medical Hypotheses</i> , 2006 , 66, 380-3	3.8	26
75	The hypoxic testis and post-meiotic expression of PAS domain proteins. <i>Seminars in Cell and Developmental Biology</i> , 2005 , 16, 547-53	7.5	37
74	Copper-dependent activation of hypoxia-inducible factor (HIF)-1: implications for ceruloplasmin regulation. <i>Blood</i> , 2005 , 105, 4613-9	2.2	224
73	Integration of oxygen signaling at the consensus HRE. <i>Science Signaling</i> , 2005 , 2005, re12	8.8	622
72	Induction of the hypoxia-inducible factor system by low levels of heat shock protein 90 inhibitors. <i>Cancer Research</i> , 2005 , 65, 11094-100	10.1	35
71	Norepinephrine-induced acute heart failure in transgenic mice overexpressing erythropoietin. <i>Cardiovascular Research</i> , 2004 , 61, 105-14	9.9	10
70	Simultaneous exposure of rats to dioxin and carbon monoxide reduces the xenobiotic but not the hypoxic response. <i>Biological Chemistry</i> , 2004 , 385, 291-4	4.5	17

69	Cardiac remodeling in erythropoietin-transgenic mice. <i>Cellular Physiology and Biochemistry</i> , 2004 , 14, 277-84	3.9	13
68	Mitochondrial reactive oxygen species control the transcription factor CHOP-10/GADD153 and adipocyte differentiation: a mechanism for hypoxia-dependent effect. <i>Journal of Biological Chemistry</i> , 2004 , 279, 40462-9	5.4	181
67	A dominant-negative isoform of hypoxia-inducible factor-1 alpha specifically expressed in human testis. <i>Biology of Reproduction</i> , 2004 , 71, 331-9	3.9	28
66	Physiologic responses to hypoxia and implications for hypoxia-inducible factors in the pathogenesis of rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2004 , 50, 10-23		83
65	Interaction of the PAS B domain with HSP90 accelerates hypoxia-inducible factor-1alpha stabilization. <i>Cellular Physiology and Biochemistry</i> , 2004 , 14, 351-60	3.9	107
64	The antimycotic ciclopirox olamine induces HIF-1alpha stability, VEGF expression, and angiogenesis. <i>FASEB Journal</i> , 2003 , 17, 761-3	0.9	87
63	Mechanisms of hypoxic gene regulation of angiogenesis factor Cyr61 in melanoma cells. <i>Journal of Biological Chemistry</i> , 2003 , 278, 45651-60	5.4	81
62	Targeted disruption of the mouse PAS domain serine/threonine kinase PASKIN. <i>Molecular and Cellular Biology</i> , 2003 , 23, 6780-9	4.8	32
61	Hypoxic up-regulation of erythroid 5-aminolevulinate synthase. <i>Blood</i> , 2003 , 101, 348-50	2.2	52
60	Hemostasis and coagulation at a hematocrit level of 0.85: functional consequences of erythrocytosis. <i>Blood</i> , 2003 , 101, 4416-22	2.2	89
59	Oxygen-regulated expression of TGF-beta 3, a growth factor involved in trophoblast differentiation. <i>Placenta</i> , 2003 , 24, 941-50	3.4	56
58	Iron chelation, angiogenesis and tumor therapy. <i>International Journal of Cancer</i> , 2003 , 106, 458-9	7.5	11
57	The cellular oxygen tension regulates expression of the endoplasmic oxidoreductase ERO1-Lalpha. <i>FEBS Journal</i> , 2003 , 270, 2228-35		109
56	The hypoxia-inducible factor-1 alpha is a negative factor for tumor therapy. <i>Oncogene</i> , 2003 , 22, 3213-20	5.2	293
55	Chronic mild hypoxia protects heart-derived H9c2 cells against acute hypoxia/reoxygenation by regulating expression of the SUR2A subunit of the ATP-sensitive K+ channel. <i>Journal of Biological Chemistry</i> , 2003 , 278, 31444-55	5.4	74
54	Cellular adaptation to hypoxia: O2-sensing protein hydroxylases, hypoxia-inducible transcription factors, and O2-regulated gene expression. <i>FASEB Journal</i> , 2002 , 16, 1151-62	0.9	954
53	Physiologically low oxygen concentrations in fetal skin regulate hypoxia-inducible factor 1 and transforming growth factor-beta3. <i>FASEB Journal</i> , 2002 , 16, 411-3	0.9	64
52	Heat induction of the unphosphorylated form of hypoxia-inducible factor-1alpha is dependent on heat shock protein-90 activity. <i>Journal of Biological Chemistry</i> , 2002 , 277, 9262-7	5.4	116

51	DNA-binding activity of hypoxia-inducible factors (HIFs). <i>Methods in Molecular Biology</i> , 2002 , 196, 117-291.4	8
50	Isoform-specific expression of hypoxia-inducible factor-1alpha during the late stages of mouse spermiogenesis. <i>Molecular Endocrinology</i> , 2002 , 16, 234-43	43
49	Normoxic induction of the hypoxia-inducible factor 1alpha by insulin and interleukin-1beta involves the phosphatidylinositol 3-kinase pathway. <i>FEBS Letters</i> , 2002 , 512, 157-62	3.8 220
48	Characterization of HIF-1 alpha overexpressing HeLa cells and implications for gene therapy. <i>Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology</i> , 2002 , 133, 475-81	3.2 10
47	Isoform-Specific Expression of Hypoxia-Inducible Factor-1 During the Late Stages of Mouse Spermiogenesis. <i>Molecular Endocrinology</i> , 2002 , 16, 234-243	27
46	Rescue of hypoxia-inducible factor-1alpha-deficient tumor growth by wild-type cells is independent of vascular endothelial growth factor. <i>Cancer Research</i> , 2002 , 62, 2962-70	10.1 37
45	Attenuation of HIF-1 DNA-binding activity limits hypoxia-inducible endothelin-1 expression. <i>Pflugers Archiv European Journal of Physiology</i> , 2001 , 443, 240-9	4.6 58
44	HIF-1 is expressed in normoxic tissue and displays an organ-specific regulation under systemic hypoxia. <i>FASEB Journal</i> , 2001 , 15, 2445-53	0.9 580
43	Hypoxia affects expression of circadian genes PER1 and CLOCK in mouse brain. <i>FASEB Journal</i> , 2001 , 15, 2613-22	0.9 104
42	Dissecting hypoxia-dependent and hypoxia-independent steps in the HIF-1alpha activation cascade: implications for HIF-1alpha gene therapy. <i>FASEB Journal</i> , 2001 , 15, 2715-7	0.9 63
41	Mammalian PASKIN, a PAS-serine/threonine kinase related to bacterial oxygen sensors. <i>Biochemical and Biophysical Research Communications</i> , 2001 , 288, 757-64	3.4 25
40	Regulation of the hypoxia-inducible factor-1 alpha. ARNT is not necessary for hypoxic induction of HIF-1 alpha in the nucleus. <i>Advances in Experimental Medicine and Biology</i> , 2000 , 475, 87-99	3.6 16
39	Genetically Modified Mouse Models in Studies on Cutaneous Wound Healing. <i>Experimental Physiology</i> , 2000 , 85, 687-704	2.4 22
38	Efficient translation of mouse hypoxia-inducible factor-1alpha under normoxic and hypoxic conditions. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 2000 , 1493, 125-34	61
37	Effects of hypobaric hypoxia on vascular endothelial growth factor and the acute phase response in subjects who are susceptible to high-altitude pulmonary oedema. <i>European Journal of Applied Physiology</i> , 2000 , 81, 497-503	3.4 48
36	Hypoxia-regulated gene expression in fetal wound regeneration and adult wound repair. <i>Pediatric Surgery International</i> , 2000 , 16, 232-6	2.1 48
35	Epolones induce erythropoietin expression via hypoxia-inducible factor-1 activation. <i>Blood</i> , 2000 , 96, 1558-1565	2.2 75
34	Nitric oxide prevents cardiovascular disease and determines survival in polyglobulic mice overexpressing erythropoietin. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000 , 97, 11609-13	11.5 216

33	Epilones induce erythropoietin expression via hypoxia-inducible factor-1 α activation. <i>Blood</i> , 2000 , 96, 1558-1565	2.2	1
32	General applicability of chicken egg yolk antibodies: the performance of IgY immunoglobulins raised against the hypoxia-inducible factor 1 α . <i>FASEB Journal</i> , 1999 , 13, 81-8	0.9	102
31	Oxygen tension modulates beta-globin switching in embryoid bodies. <i>FASEB Journal</i> , 1999 , 13, 285-95	0.9	25
30	Molecular Biology of Hypoxia-Inducible Factor-1 1999 , 269-276		1
29	Oxygen-regulated erythropoietin gene expression is dependent on a CpG methylation-free hypoxia-inducible factor-1 DNA-binding site. <i>FEBS Journal</i> , 1998 , 253, 771-7		106
28	Cellular and developmental control of O ₂ homeostasis by hypoxia-inducible factor 1 α . <i>Genes and Development</i> , 1998 , 12, 149-62	12.6	1842
27	Mouse Hypoxia-Inducible Factor-1 α s Encoded by Two Different mRNA Isoforms: Expression From a Tissue-Specific and a Housekeeping-Type Promoter. <i>Blood</i> , 1998 , 91, 3471-3480	2.2	65
26	Mouse Hypoxia-Inducible Factor-1 α s Encoded by Two Different mRNA Isoforms: Expression From a Tissue-Specific and a Housekeeping-Type Promoter. <i>Blood</i> , 1998 , 91, 3471-3480	2.2	4
25	Distinct costimulatory molecules are required for the induction of effector and memory cytotoxic T lymphocytes. <i>Journal of Experimental Medicine</i> , 1997 , 185, 251-62	16.6	117
24	Oxygen-regulated transferrin expression is mediated by hypoxia-inducible factor-1. <i>Journal of Biological Chemistry</i> , 1997 , 272, 20055-62	5.4	289
23	HIF-1, a Mediator of the Molecular Response to Hypoxia. <i>Physiology</i> , 1997 , 12, 214-218	9.8	3
22	Detection of erythropoietin in human liquor: intrinsic erythropoietin production in the brain. <i>Kidney International</i> , 1997 , 51, 416-8	9.9	114
21	Hypoxia-inducible factor-1 α is regulated at the post-mRNA level. <i>Kidney International</i> , 1997 , 51, 560-3	9.9	106
20	The hypoxia-inducible factor-1 DNA recognition site is cAMP-responsive. <i>Kidney International</i> , 1997 , 51, 564-6	9.9	43
19	Oxygen- and dioxin-regulated gene expression in mouse hepatoma cells. <i>Kidney International</i> , 1997 , 51, 567-74	9.9	67
18	The mouse gene for hypoxia-inducible factor-1 α --genomic organization, expression and characterization of an alternative first exon and 5'Tflanking sequence. <i>FEBS Journal</i> , 1997 , 246, 155-65		41
17	Nucleotide sequence, chromosomal assignment and mRNA expression of mouse hypoxia-inducible factor-1 α . <i>Biochemical and Biophysical Research Communications</i> , 1996 , 223, 54-9	3.4	108
16	Oxygen supply and oxygen-dependent gene expression in differentiating embryonic stem cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996 , 93, 2867-72	11.5	120

15	Little difference. <i>Nature</i> , 1996 , 380, 100	50.4	11
14	Erythropoietin gene expression in human, monkey and murine brain. <i>European Journal of Neuroscience</i> , 1996 , 8, 666-76	3.5	477
13	B-cell maturation in chimaeric mice deficient for the heat stable antigen (HSA/mouse CD24). <i>Transgenic Research</i> , 1995 , 4, 173-83	3.3	29
12	Hypoxia, a novel inducer of acute phase gene expression in a human hepatoma cell line. <i>Journal of Biological Chemistry</i> , 1995 , 270, 27865-70	5.4	90
11	Localization of specific erythropoietin binding sites in defined areas of the mouse brain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995 , 92, 3717-20	11.5	423
10	The transcription factors ATF-1 and CREB-1 bind constitutively to the hypoxia-inducible factor-1 (HIF-1) DNA recognition site. <i>Nucleic Acids Research</i> , 1995 , 23, 4542-50	20.1	207
9	Mitochondria contaminate databases. <i>Trends in Genetics</i> , 1995 , 11, 167-8	8.5	7
8	The heat-stable antigen can alter very late antigen 4-mediated adhesion. <i>Journal of Experimental Medicine</i> , 1994 , 179, 1391-5	16.6	52
7	Reannealing of artificial heteroduplexes generated during PCR-mediated isotyping. <i>Trends in Genetics</i> , 1991 , 7, 178	8.5	10
6	The genes for a mouse hematopoietic differentiation marker called the heat-stable antigen. <i>European Journal of Immunology</i> , 1991 , 21, 1039-46	6.1	50
5	The 5'flanking region and chromosomal localization of the gene encoding human platelet membrane glycoprotein Ib alpha. <i>Gene</i> , 1989 , 85, 517-24	3.8	57
4	Isolation and Characterization of Human Blood Platelet mRNA and Construction of a cDNA Library in λ t11. <i>Thrombosis and Haemostasis</i> , 1989 , 61, 448-453	7	35
3	Structure of the human blood platelet membrane glycoprotein Ib alpha gene. <i>Biochemical and Biophysical Research Communications</i> , 1988 , 156, 389-95	3.4	112
2	Refolding of bacteriorhodopsin. Protease V8 fragmentation and chromophore reconstitution from proteolytic V8 fragments. <i>FEBS Journal</i> , 1988 , 177, 125-33		34
1	Androglobin, a chimeric mammalian globin, is required for male fertility. <i>ELife</i> , 11,	8.9	1