Finn M Radtke

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/124739/publications.pdf Version: 2024-02-01



FINN M PADTER

#	Article	IF	CITATIONS
1	Presurgical cognitive impairment is associated with postoperative delirium and postoperative cognitive dysfunction. Minerva Anestesiologica, 2020, 86, 394-403.	0.6	18
2	Câ€reactive protein for risk prediction of postâ€operative delirium and postâ€operative neurocognitive disorder. Acta Anaesthesiologica Scandinavica, 2019, 63, 1282-1289.	0.7	23
3	Electroencephalography-Guided Anesthetic Administration and Postoperative Delirium. JAMA - Journal of the American Medical Association, 2019, 321, 2469.	3.8	4
4	Diabetes, but Not Hypertension and Obesity, Is Associated with Postoperative Cognitive Dysfunction. Dementia and Geriatric Cognitive Disorders, 2018, 46, 193-206.	0.7	24
5	Association of obesity, diabetes and hypertension with cognitive impairment in older age. Clinical Epidemiology, 2018, Volume 10, 853-862.	1.5	29
6	Patients prone for postoperative delirium. Current Opinion in Anaesthesiology, 2016, 29, 384-390.	0.9	42
7	No convincing association between postâ€operative delirium and postâ€operative cognitive dysfunction: a secondary analysis. Acta Anaesthesiologica Scandinavica, 2016, 60, 1404-1414.	0.7	30
8	How can postoperative delirium be predicted in advance? A secondary analysis comparing three methods of early assessment in elderly patients. Minerva Anestesiologica, 2016, 82, 751-9.	0.6	10
9	Postoperative delirium is an independent risk factor for posttraumatic stress disorder in the elderly patient. European Journal of Anaesthesiology, 2015, 32, 147-151.	0.7	53
10	Delirium, Sedation and Analgesia in the Intensive Care Unit: A Multinational, Two-Part Survey among Intensivists. PLoS ONE, 2014, 9, e110935.	1.1	89
11	Incidence and Risk Factors for Cognitive Dysfunction in Patients with Severe Systemic Disease. Journal of International Medical Research, 2012, 40, 612-620.	0.4	13
12	How to implement monitoring tools for sedation, pain and delirium in the intensive care unit: an experimental cohort study. Intensive Care Medicine, 2012, 38, 1974-1981.	3.9	69
13	Delirium in the postanaesthesia period. Current Opinion in Anaesthesiology, 2011, 24, 670-675.	0.9	57
14	Documented Intraoperative Hypotension According to the Three Most Common Definitions Does Not Match the Application of Antihypotensive Medication. Journal of International Medical Research, 2011, 39, 846-856.	0.4	16
15	Different assessment tools for intensive care unit delirium: Which score to use?*. Critical Care Medicine, 2010, 38, 409-418.	0.4	297
16	Duration of fluid fasting and choice of analgesic are modifiable factors for early postoperative delirium. European Journal of Anaesthesiology, 2010, 27, 411-416.	0.7	125
17	A Comparison of Three Scores to Screen for Delirium on the Surgical Ward. World Journal of Surgery, 2010, 34, 487-494.	0.8	74
18	Postoperative cognitive dysfunction: computerized and conventional tests showed only moderate inter-rater reliability. Journal of Anesthesia, 2010, 24, 518-525.	0.7	10

FINN M RADTKE

#	Article	IF	CITATIONS
19	Documentation of Post-operative Nausea and Vomiting in Routine Clinical Practice. Journal of International Medical Research, 2010, 38, 1034-1041.	0.4	44
20	Diagnosing Delirium. JAMA - Journal of the American Medical Association, 2010, 304, 2124.	3.8	5
21	Delayed Treatment of Delirium Increases Mortality Rate in Intensive Care Unit Patients. Journal of International Medical Research, 2010, 38, 1584-1595.	0.4	119
22	Remifentanil Reduces the Incidence of Post-Operative Delirium. Journal of International Medical Research, 2010, 38, 1225-1232.	0.4	30
23	Comparison of three scores to screen for delirium in the recovery room. British Journal of Anaesthesia, 2008, 101, 338-343.	1.5	105