Ping-Jen Chen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1262091/publications.pdf

Version: 2024-02-01

567281 580821 43 720 15 25 citations h-index g-index papers 47 47 47 1222 docs citations times ranked citing authors all docs

#	Article	IF	Citations
1	Relapse insomnia increases greater risk of anxiety and depression: evidence from a population-based 4-year cohort study. Sleep Medicine, 2017, 38, 122-129.	1.6	57
2	Predicting Cause-Specific Mortality of Older Men Living in the Veterans Home by Handgrip Strength and Walking Speed: A 3-Year, Prospective Cohort Study in Taiwan. Journal of the American Medical Directors Association, 2012, 13, 517-521.	2.5	56
3	Hip fracture and risk of acute myocardial infarction: A nationwide study. Journal of Bone and Mineral Research, 2013, 28, 404-411.	2.8	54
4	Methodological Factors in Determining Rates of Dementia in Transient Ischemic Attack and Stroke. Stroke, 2015, 46, 641-646.	2.0	53
5	Methodological Factors in Determining Risk of Dementia After Transient Ischemic Attack and Stroke. Stroke, 2015, 46, 1494-1500.	2.0	48
6	Long-term Mortality Risk After Hyperglycemic Crisis Episodes in Geriatric Patients With Diabetes: A National Population-Based Cohort Study. Diabetes Care, 2015, 38, 746-751.	8.6	36
7	Communication and Behavior of Palliative Care Physicians of Patients With Cancer Near End of Life in Three East Asian Countries. Journal of Pain and Symptom Management, 2021, 61, 315-322.e1.	1.2	36
8	Implementing a novel model for hospice and palliative care in the emergency department. Medicine (United States), 2017, 96, e6943.	1.0	26
9	Advance Care Planning for Older People with Cancer and Its Implications in Asia: Highlighting the Mental Capacity and Relational Autonomy. Geriatrics (Switzerland), 2018, 3, 43.	1.7	26
10	Primary Sjögren's syndrome and risk of ischemic stroke: a nationwide study. Clinical Rheumatology, 2014, 33, 931-937.	2.2	25
11	Lower Urinary Tract Symptoms Are Associated with Increased Risk of Dementia among the Elderly: A Nationwide Study. BioMed Research International, 2015, 2015, 1-7.	1.9	25
12	Comparison of healthcare utilization and lifeâ€sustaining interventions between elderly patients with dementia and those with cancer near the end of life: A nationwide, populationâ€based study in Taiwan. Geriatrics and Gerontology International, 2017, 17, 2545-2551.	1.5	25
13	Association between palliative care and life-sustaining treatments for patients with dementia: A nationwide 5-year cohort study. Palliative Medicine, 2018, 32, 622-630.	3.1	25
14	2019 Taipei Declaration on Advance Care Planning: A Cultural Adaptation of End-of-Life Care Discussion. Journal of Palliative Medicine, 2019, 22, 1175-1177.	1.1	22
15	Cumulative use of therapeutic bladder anticholinergics and the risk of dementia in patients with lower urinary tract symptoms: a nationwide 12-year cohort study. BMC Geriatrics, 2019, 19, 380.	2.7	21
16	Feasibility and acceptability of a culturally adapted advance care planning intervention for people living with advanced cancer and their families: A mixed methods study. Palliative Medicine, 2020, 34, 651-666.	3.1	18
17	Emergency geriatric assessment: A novel comprehensive screen tool for geriatric patients in the emergency department. American Journal of Emergency Medicine, 2018, 36, 143-146.	1.6	15
18	Primary Sjögren's Syndrome and the Risk of Acute Myocardial Infarction: A Nationwide Study. Acta Cardiologica Sinica, 2013, 29, 124-31.	0.2	15

#	Article	IF	CITATIONS
19	Septic arthritis significantly increased the long-term mortality in geriatric patients. BMC Geriatrics, 2017, 17, 178.	2.7	13
20	Prevalence of Geriatric Syndromes and the Need for Hospice Care in Older Patients of the Emergency Department: A Study in an Asian Medical Center. Emergency Medicine International, 2020, 2020, 1-9.	0.8	11
21	To hydrate or not to hydrate? The effect of hydration on survival, symptoms and quality of dying among terminally ill cancer patients. BMC Palliative Care, 2021, 20, 13.	1.8	11
22	Cognitively-Related Basic Activities of Daily Living Impairment Greatly Increases the Risk of Death in Alzheimers Disease. PLoS ONE, 2016, 11, e0160671.	2.5	10
23	Depression is a major risk factor for the development of dementia in people with lower urinary tract symptoms: A nationwide population-based study. PLoS ONE, 2019, 14, e0217984.	2.5	9
24	The Association between Home Healthcare and Burdensome Transitions at the End-of-Life in People with Dementia: A 12-Year Nationwide Population-Based Cohort Study. International Journal of Environmental Research and Public Health, 2020, 17, 9255.	2.6	8
25	End-of-life care in cancer and dementia: a nationwide population-based study of palliative care policy changes. BMJ Supportive and Palliative Care, 2022, 12, e384-e392.	1.6	7
26	HOme-based Longitudinal Investigation of the multidiSciplinary Team Integrated Care (HOLISTIC): protocol of a prospective nationwide cohort study. BMC Geriatrics, 2020, 20, 511.	2.7	7
27	Preferences on the Timing of Initiating Advance Care Planning and Withdrawing Life-Sustaining Treatment between Terminally-Ill Cancer Patients and Their Main Family Caregivers: A Prospective Study. International Journal of Environmental Research and Public Health, 2020, 17, 7954.	2.6	7
28	Burden of Healthcare Utilization among Chronic Obstructive Pulmonary Disease Patients with and without Cancer Receiving Palliative Care: A Population-Based Study in Taiwan. International Journal of Environmental Research and Public Health, 2020, 17, 4980.	2.6	6
29	Prevalence and severity of symptoms and signs in patients with advanced cancer in the last days of life: the East Asian collaborative cross-cultural study to elucidate the dying process (EASED). Supportive Care in Cancer, 2022, 30, 5499-5508.	2.2	6
30	Comparison of the accuracy of clinicians' prediction of survival and Palliative Prognostic Score: an East Asian cross-cultural study. Supportive Care in Cancer, 2022, 30, 2367-2374.	2.2	5
31	Impact of home healthcare on end-of-life outcomes for people with dementia: a systematic review. BMC Geriatrics, 2022, 22, 80.	2.7	5
32	Increased risk of chronic kidney disease in patients with rosacea: A nationwide population-based matched cohort study. PLoS ONE, 2017, 12, e0180446.	2.5	4
33	Assessment of Changes in Symptoms Is Feasible and Prognostic in the Last Weeks of Life: An International Multicenter Cohort Study. Journal of Palliative Medicine, 2022, 25, 388-395.	1.1	4
34	Professionals' experiences and attitudes toward use of Traditional Chinese Medicine in hospice palliative inpatient care units: A multicenter survey in Taiwan. Integrative Medicine Research, 2021, 10, 100642.	1.8	3
35	Physical and psychological symptoms and signs in dying digestive tract cancer patients: the East Asian collaborative cross-cultural Study to Elucidate the Dying process (EASED). Supportive Care in Cancer, 2021, 29, 3603-3612.	2.2	3
36	Medical care needs for patients receiving home healthcare in Taiwan: Do gender and income matter?. PLoS ONE, 2021, 16, e0247622.	2.5	3

#	Article	IF	CITATIONS
37	Factors related to spiritual well-being in the last days of life in three East Asian countries: An international multicenter prospective cohort study. Palliative Medicine, 2021, 35, 1564-1577.	3.1	3
38	Association between Anti-Psychotic Drugs Use and Hip Fractures in Patients with Dementia: A Nationwide Population-Based Study. International Journal of Environmental Research and Public Health, 2021, 18, 8118.	2.6	3
39	Reliability and Validity of the Traditional Chinese Version of the Advance Care Planning Engagement Survey: A Pilot Evaluation in Taiwanese Outpatients. Journal of Palliative Care, 2022, 37, 273-279.	1.0	3
40	Differences between dedicated and not dedicated hospice physicians in symptoms and signs improvement among advanced cancer patients. Medicine (United States), 2021, 100, e26915.	1.0	2
41	Association between the amount of artificial hydration and quality of dying among terminally ill patients with cancer: The East Asian Collaborative Crossâ€Cultural Study to Elucidate the Dying Process. Cancer, 2022, 128, 1699-1708.	4.1	2
42	Exploring the Impact of Different Types of Do-Not-Resuscitate Consent on End-of-Life Treatments among Patients with Advanced Kidney Disease: An Observational Study. International Journal of Environmental Research and Public Health, 2021, 18, 8194.	2.6	1
43	Comparison of Objective Prognostic Score and Palliative Prognostic Score performance in inpatients with advanced cancer in Japan and Korea. Palliative and Supportive Care, 0, , 1-9.	1.0	1