

Yong-June Kim

List of Publications by Citations

Source: <https://exaly.com/author-pdf/1263368/yong-june-kim-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

62

papers

966

citations

15

h-index

29

g-index

64

ext. papers

1,148

ext. citations

3.5

avg, IF

3.66

L-index

#	Paper	IF	Citations
62	Expression signature of E2F1 and its associated genes predict superficial to invasive progression of bladder tumors. <i>Journal of Clinical Oncology</i> , 2010 , 28, 2660-7	2.2	230
61	HOXA9, ISL1 and ALDH1A3 methylation patterns as prognostic markers for nonmuscle invasive bladder cancer: array-based DNA methylation and expression profiling. <i>International Journal of Cancer</i> , 2013 , 133, 1135-42	7.5	72
60	Methylation of the RUNX3 promoter as a potential prognostic marker for bladder tumor. <i>Journal of Urology</i> , 2008 , 180, 1141-5	2.5	67
59	Impact of obesity in patients with urolithiasis and its prognostic usefulness in stone recurrence. <i>Journal of Urology</i> , 2008 , 179, 570-4	2.5	66
58	EFEMP1 as a novel DNA methylation marker for prostate cancer: array-based DNA methylation and expression profiling. <i>Clinical Cancer Research</i> , 2011 , 17, 4523-30	12.9	53
57	Electrochemical detection of methylated DNA on a microfluidic chip with nanoelectrokinetic pre-concentration. <i>Biosensors and Bioelectronics</i> , 2018 , 107, 103-110	11.8	41
56	Gene signatures for the prediction of response to Bacillus Calmette-Guerin immunotherapy in primary pT1 bladder cancers. <i>Clinical Cancer Research</i> , 2010 , 16, 2131-7	12.9	38
55	Multi-institutional analysis of localized renal cell carcinoma that demonstrates the impact of diabetic status on prognosis after nephrectomy. <i>Annals of Surgical Oncology</i> , 2013 , 20, 3662-8	3.1	23
54	The establishment of KORCC (Korean Renal Cell Carcinoma) database. <i>Investigative and Clinical Urology</i> , 2016 , 57, 50-7	1.9	22
53	Body mass index influences prostate-specific antigen in men younger than 60 years of age. <i>International Journal of Urology</i> , 2007 , 14, 1009-12	2.3	20
52	Impact of Young Age at Diagnosis on Survival in Patients with Surgically Treated Renal Cell Carcinoma: a Multicenter Study. <i>Journal of Korean Medical Science</i> , 2016 , 31, 1976-1982	4.7	18
51	Prognostic Significance of Preoperative Neutrophil-to-Lymphocyte Ratio in Nonmetastatic Renal Cell Carcinoma: A Large, Multicenter Cohort Analysis. <i>BioMed Research International</i> , 2016 , 2016, 5634148	3.8	17
50	Prognostic Impact of Nutritional Status Assessed by the Controlling Nutritional Status (CONUT) Score in Patients with Surgically Treated Renal Cell Carcinoma. <i>Nutrition and Cancer</i> , 2018 , 70, 886-894	2.8	17
49	DNA methylation of GSTP1 in human prostate tissues: pyrosequencing analysis. <i>Korean Journal of Urology</i> , 2012 , 53, 200-5		15
48	The association between metabolic syndrome and prostate-specific antigen levels. <i>International Journal of Urology</i> , 2008 , 15, 905-9	2.3	15
47	Association between Perioperative Blood Transfusion and Oncologic Outcomes after Curative Surgery for Renal Cell Carcinoma. <i>Journal of Cancer</i> , 2016 , 7, 965-72	4.5	15
46	Low preoperative serum cholesterol level is associated with aggressive pathologic features and poor cancer-specific survival in patients with surgically treated renal cell carcinoma. <i>International Journal of Clinical Oncology</i> , 2018 , 23, 142-150	4.2	14

45	Hypertriglyceridemia is associated with increased risk for stone recurrence in patients with urolithiasis. <i>Urology</i> , 2014 , 84, 766-71	1.6	13
44	Hypertension influences recurrent stone formation in nonobese stone formers. <i>Urology</i> , 2011 , 77, 1059-63	6.3	13
43	mRNA Expression Is Associated with the Aggressiveness of Prostate Cancer. <i>Journal of Korean Medical Science</i> , 2018 , 33, e303	4.7	13
42	Effect of renal insufficiency on stone recurrence in patients with urolithiasis. <i>Journal of Korean Medical Science</i> , 2014 , 29, 1132-7	4.7	10
41	GSTT1 as a Prognosticator for Recurrence and Progression in Patients with Non-Muscle-Invasive Bladder Cancer. <i>Disease Markers</i> , 2010 , 29, 81-87	3.2	10
40	The age-adjusted Charlson comorbidity index as a predictor of overall survival of surgically treated non-metastatic clear cell renal cell carcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020 , 146, 187-196	4.9	10
39	Role of 1,25-dihydroxy vitamin D3 and parathyroid hormone in urinary calcium excretion in calcium stone formers. <i>Yonsei Medical Journal</i> , 2014 , 55, 1326-32	3	9
38	Impact of the ASA Physical Status Score on Adjuvant Chemotherapy Eligibility and Survival of Upper Tract Urothelial Carcinoma Patients: a Multicenter Study. <i>Journal of Korean Medical Science</i> , 2017 , 32, 335-342	4.7	8
37	Molecular Progression Risk Score for Prediction of Muscle Invasion in Primary T1 High-Grade Bladder Cancer. <i>Clinical Genitourinary Cancer</i> , 2018 , 16, 274-280	3.3	8
36	Clinical Implications and Prognostic Values of Prostate Cancer Susceptibility Candidate Methylation in Primary Nonmuscle Invasive Bladder Cancer. <i>Disease Markers</i> , 2015 , 2015, 402963	3.2	8
35	A Low Geriatric Nutritional Risk Index is Associated with Aggressive Pathologic Characteristics and Poor Survival after Nephrectomy in Clear Renal Cell Carcinoma: A Multicenter Retrospective Study. <i>Nutrition and Cancer</i> , 2020 , 72, 88-97	2.8	8
34	Age-dependent prognostic value of body mass index for non-metastatic clear cell renal cell carcinoma: A large multicenter retrospective analysis. <i>Journal of Surgical Oncology</i> , 2018 , 118, 199-205	2.8	8
33	Chronological Trends in Clinical and Urinary Metabolic Features over 20 Years in Korean Urolithiasis Patients. <i>Journal of Korean Medical Science</i> , 2017 , 32, 1496-1501	4.7	7
32	Lower Levels of Human MOB3B Are Associated with Prostate Cancer Susceptibility and Aggressive Clinicopathological Characteristics. <i>Journal of Korean Medical Science</i> , 2015 , 30, 937-42	4.7	7
31	The predictive value of GSTT1 polymorphisms in predicting the early response to induction BCG therapy in patients with non-muscle invasive bladder cancer. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 458-65	2.8	7
30	Methylation Signature for Prediction of Progression Free Survival in Surgically Treated Clear Cell Renal Cell Carcinoma. <i>Journal of Korean Medical Science</i> , 2019 , 34, e144	4.7	7
29	Sex-Specific Prognostic Significance of Obesity in Nonmetastatic Clear-Cell Renal-Cell Carcinoma in Korea: A Large Multicenter Cohort Analysis. <i>Clinical Genitourinary Cancer</i> , 2017 ,	3.3	6
28	Long-term validation of a molecular progression-associated gene classifier for prediction of muscle invasion in primary non-muscle-invasive bladder cancer. <i>Oncology Letters</i> , 2017 , 14, 2468-2474	2.6	6

27	Pyrosequencing Analysis of APC Methylation Level in Human Prostate Tissues: A Molecular Marker for Prostate Cancer. <i>Korean Journal of Urology</i> , 2013 , 54, 194-8		6
26	Changes in urinary lithogenic features over time in patients with urolithiasis. <i>Urology</i> , 2009 , 74, 51-5	1.6	6
25	Twenty-four-hour urine osmolality as a representative index of adequate hydration and a predictor of recurrence in patients with urolithiasis. <i>International Urology and Nephrology</i> , 2019 , 51, 1129-1135	2.3	5
24	Comparison of Perioperative Outcomes of Extraperitoneal Laparoscopic Radical Prostatectomy (ELRP) versus Open Radical Retropubic Prostatectomy (RRP): Single Surgeon's Initial Experience. <i>Korean Journal of Urology</i> , 2007 , 48, 131		5
23	Impact of preoperative thrombocytosis on prognosis after surgical treatment in pathological T1 and T2 renal cell carcinoma: results of a multi-institutional comprehensive study. <i>Oncotarget</i> , 2017 , 8, 64449-64458	3.3	5
22	Metabolic Characteristics and Risks Associated with Stone Recurrence in Korean Young Adult Stone Patients. <i>Journal of Endourology</i> , 2017 , 31, 806-811	2.7	4
21	Distinct metabolic characteristics and risk of stone recurrence in patients with multiple stones at the first-time presentation. <i>Urology</i> , 2014 , 84, 274-8	1.6	4
20	Utility of Smo as a Prognostic Marker for Human Bladder Tumors. <i>Korean Journal of Urology</i> , 2007 , 48, 997		4
19	ZNF492 and GPR149 methylation patterns as prognostic markers for clear cell renal cell carcinoma: Array-based DNA methylation profiling. <i>Oncology Reports</i> , 2019 , 42, 453-460	3.5	4
18	A high basal metabolic rate is an independent predictor of stone recurrence in obese patients. <i>Investigative and Clinical Urology</i> , 2021 , 62, 195-200	1.9	3
17	Impact of transobturator tape treatment on overactive bladder symptoms, particularly nocturia, in patients with mixed urinary incontinence. <i>Korean Journal of Urology</i> , 2014 , 55, 520-6		2
16	Biomarkers in Bladder Cancer: Present Status and Perspectives. <i>Biomarker Insights</i> , 2007 , 2, 1177271907092002		2
15	The Relationship between RUNX3 Inactivation and Its Pathological Features in Renal Cell Carcinoma. <i>Korean Journal of Urology</i> , 2009 , 50, 432		2
14	A Prospective Multicenter Trial of the Efficacy and Tolerability of Neoadjuvant Sunitinib for Inoperable Metastatic Renal Cell Carcinoma. <i>Journal of Korean Medical Science</i> , 2016 , 31, 1983-1988	4.7	2
13	Gender- and cholesterol-specific predictive value of body mass index in renal cell carcinoma: A multicenter study. <i>Asia-Pacific Journal of Clinical Oncology</i> , 2019 , 15, e36-e42	1.9	2
12	Change in Prostate Specific Antigen Concentration in Men with Prostate Specific Antigen Less than 2.5 ng/ml Taking Low Dose Finasteride or Dutasteride for Male Androgenetic Alopecia. <i>Journal of Urology</i> , 2017 , 198, 1340-1345	2.5	1
11	Can We Use Single Nucleotide Polymorphism and Runt Domain Transcription Factor 3 Methylation as Tumor Markers for Bladder Cancer?. <i>Korean Journal of Urology</i> , 2009 , 50, 311		1
10	Trends in clinical, operative, and pathologic characteristics of surgically treated renal mass in a Korean center: A surgical series from 1988 through 2015. <i>Investigative and Clinical Urology</i> , 2019 , 60, 184-194	1.9	1

9	A novel urinary mRNA signature using the droplet digital polymerase chain reaction platform improves discrimination between prostate cancer and benign prostatic hyperplasia within the prostate-specific antigen gray zone. <i>Investigative and Clinical Urology</i> , 2020 , 61, 411-418	1.9	1
8	Effect of pre-operative internal obturator muscle mass index in MRI on biochemical recurrence of prostate cancer patients after radical prostatectomy: a multi-center study. <i>BMC Urology</i> , 2021 , 21, 85	2.2	1
7	Integrative Transcriptome Profiling Reveals as a Novel Prognostic Marker in Non-Muscle Invasive Bladder Cancer. <i>Cancers</i> , 2021 , 13,	6.6	1
6	Nutritional status assessed by the Controlling Nutritional Status (CONUT) score as a predictor of recurrence of urolithiasis. <i>Investigative and Clinical Urology</i> , 2021 , 62, 553-559	1.9	1
5	Intraoperative allogeneic blood transfusion is associated with adverse oncological outcomes in patients with surgically treated non-metastatic clear cell renal cell carcinoma. <i>International Journal of Clinical Oncology</i> , 2020 , 25, 1551-1561	4.2	0
4	Expression of phosphorylated p21-activated kinase 4 is associated with aggressive histologic characteristics and poor prognosis in patients with surgically treated renal cell carcinoma. <i>Investigative and Clinical Urology</i> , 2021 , 62, 399-407	1.9	0
3	The prognostic value of the pretreatment serum albumin to globulin ratio for predicting adverse pathology in patients undergoing radical prostatectomy for prostate cancer. <i>Investigative and Clinical Urology</i> , 2021 , 62, 545-552	1.9	0
2	Efficacy and Tolerability of Solifenacin Fumarate with Overactive Bladder Patients: A Multicenter Observational Study. <i>Urogenital Tract Infection</i> , 2021 , 16, 8-15	0	
1	Urinary hsv2-miR-H9 to hsa-miR-3659 ratio is an effective marker for discriminating prostate cancer from benign prostate hyperplasia in patients within the prostate-specific antigen grey zone.. <i>Investigative and Clinical Urology</i> , 2022 , 63, 238-244	1.9	