

Wookyung Chung

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1272200/publications.pdf>

Version: 2024-02-01

42
papers

688
citations

687363

13
h-index

610901

24
g-index

43
all docs

43
docs citations

43
times ranked

993
citing authors

#	ARTICLE	IF	CITATIONS
1	KNOW-CKD (KoreaN cohort study for Outcome in patients With Chronic Kidney Disease): design and methods. <i>BMC Nephrology</i> , 2014, 15, 80.	1.8	156
2	Urinary Potassium Excretion and Progression of CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2019, 14, 330-340.	4.5	50
3	Association of Blood Pressure With the Progression of CKD: Findings From KNOW-CKD Study. <i>American Journal of Kidney Diseases</i> , 2021, 78, 236-245.	1.9	39
4	The clinicopathological relevance of pretransplant anti-angiotensin II type 1 receptor antibodies in renal transplantation. <i>Nephrology Dialysis Transplantation</i> , 2017, 32, gfv375.	0.7	33
5	Does Routine Bioimpedance-Guided Fluid Management Provide Additional Benefit to Non-Anuric Peritoneal Dialysis Patients? Results from Compass Clinical Trial. <i>Peritoneal Dialysis International</i> , 2018, 38, 131-138.	2.3	33
6	Baseline General Characteristics of the Korean Chronic Kidney Disease: Report from the KoreaN Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). <i>Journal of Korean Medical Science</i> , 2017, 32, 221.	2.5	31
7	Serum hepcidin may be a novel uremic toxin, which might be related to erythropoietin resistance. <i>Scientific Reports</i> , 2017, 7, 4260.	3.3	27
8	Circulating levels of soluble receptor for advanced glycation end product are inversely associated with vascular calcification in patients on haemodialysis independent of S100A12 (EN-RAGE) levels. <i>Nephrology</i> , 2013, 18, 777-782.	1.6	25
9	Effects of Interleukin-6 T15A Single Nucleotide Polymorphism on Baseline Peritoneal Solute Transport Rate in Incident Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , 2009, 29, 81-88.	2.3	24
10	Sleep Duration and Health-Related Quality of Life in Predialysis CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2018, 13, 858-865.	4.5	24
11	Dietary Protein Intake, Protein Energy Wasting, and the Progression of Chronic Kidney Disease: Analysis from the KNOW-CKD Study. <i>Nutrients</i> , 2019, 11, 121.	4.1	19
12	Associations between Soluble Receptor for Advanced Glycation End Products (sRAGE) and S100A12 (EN-RAGE) with Mortality in Long-term Hemodialysis Patients. <i>Journal of Korean Medical Science</i> , 2017, 32, 54.	2.5	18
13	Hyperuricemia Is an Independent Risk Factor for Mortality Only if Chronic Kidney Disease Is Present. <i>American Journal of Nephrology</i> , 2013, 37, 452-461.	3.1	16
14	Low Resistin Level is Associated with Poor Hospitalization-Free Survival in Hemodialysis Patients. <i>Journal of Korean Medical Science</i> , 2012, 27, 377.	2.5	12
15	Comparison of the Efficacy and Safety Profile of Morning Administration of Controlled-release Simvastatin Versus Evening Administration of Immediate-release Simvastatin in Chronic Kidney Disease Patients With Dyslipidemia. <i>Clinical Therapeutics</i> , 2014, 36, 1182-1190.	2.5	12
16	Incidence of cardiovascular events and mortality in Korean patients with chronic kidney disease. <i>Scientific Reports</i> , 2021, 11, 1131.	3.3	12
17	Body Mass Index, waist circumference, and health-related quality of life in adults with chronic kidney disease. <i>Quality of Life Research</i> , 2019, 28, 1075-1083.	3.1	11
18	Liver cirrhosis leads to poorer survival in patients with end-stage renal disease. <i>Korean Journal of Internal Medicine</i> , 2016, 31, 730-738.	1.7	11

#	ARTICLE	IF	CITATIONS
19	Serum Hepcidin and Iron Indices Affect Anemia Status Differently According to the Kidney Function of Non-Dialysis Chronic Kidney Disease Patients: Korean Cohort Study For Outcome in Patients with Chronic Kidney Disease (KNOW-CKD). <i>Kidney and Blood Pressure Research</i> , 2017, 42, 1183-1192.	2.0	10
20	Discrepant glomerular filtration rate trends from creatinine and cystatin C in patients with chronic kidney disease: results from the KNOW-CKD cohort. <i>BMC Nephrology</i> , 2020, 21, 280.	1.8	10
21	Relationship between brachial-ankle and heart-femoral pulse wave velocities and the rapid decline of kidney function. <i>Scientific Reports</i> , 2018, 8, 821.	3.3	9
22	The Correlation of Serum Osteoprotegerin with Non-Traditional Cardiovascular Risk Factors and Arterial Stiffness in Patients with Pre-Dialysis Chronic Kidney Disease: Results from the KNOW-CKD Study. <i>Journal of Korean Medical Science</i> , 2018, 33, e322.	2.5	9
23	Rapid Weight Change Over Time Is a Risk Factor for Adverse Outcomes in Patients With Predialysis Chronic Kidney Disease: A Prospective Cohort Study. , 2021, 31, 569-578.		9
24	Inflammation Alters Relationship Between High-Density Lipoprotein Cholesterol and Cardiovascular Risk in Patients With Chronic Kidney Disease: Results From KNOW-CKD. <i>Journal of the American Heart Association</i> , 2021, 10, e021731.	3.7	9
25	Albuminuria as a Risk Factor for Anemia in Chronic Kidney Disease: Result from the Korean Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). <i>PLoS ONE</i> , 2015, 10, e0139747.	2.5	8
26	Association of serum adiponectin concentration with aortic arterial stiffness in chronic kidney disease: from the KNOW-CKD study. <i>Clinical and Experimental Nephrology</i> , 2017, 21, 608-616.	1.6	7
27	Smoking Cessation and Coronary Artery Calcification in CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2021, 16, 870-879.	4.5	7
28	Association of blood pressure with cardiovascular outcome and mortality: results from the KNOW-CKD study. <i>Nephrology Dialysis Transplantation</i> , 2022, 37, 1722-1730.	0.7	7
29	Association Between Systolic Blood Pressure Variability and Major Adverse Cardiovascular Events in Korean Patients With Chronic Kidney Disease: Findings From KNOW-CKD. <i>Journal of the American Heart Association</i> , 2022, 11, .	3.7	7
30	Hepcidin, iron indices and bone mineral metabolism in non-dialysis chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, 147-154.	0.7	6
31	The effect of interactions between proteinuria, activity of fibroblast growth factor 23 and serum phosphate on renal progression in patients with chronic kidney disease: a result from the Korean cohort study for Outcome in patients With Chronic Kidney Disease study. <i>Nephrology Dialysis Transplantation</i> . 2020. 35. 438-446.	0.7	6
32	Mayo imaging classification is a good predictor of rapid progress among Korean patients with autosomal dominant polycystic kidney disease: results from the KNOW-CKD study. <i>Kidney Research and Clinical Practice</i> , 2022, 41, 432-441.	2.2	6
33	Association between serum osteoprotegerin level and renal prognosis in nondialysis patients with chronic kidney disease in the Korean Cohort Study for Outcomes in Patients with Chronic Kidney Disease (the KNOW-CKD Study). <i>Kidney Research and Clinical Practice</i> , 2022, 41, 200-208.	2.2	5
34	Low serum adiponectin level is associated with better physical health-related quality of life in chronic kidney disease. <i>Scientific Reports</i> , 2021, 11, 10928.	3.3	4
35	Polypharmacy and the Progression of Chronic Kidney Disease: Korean Cohort Study for Outcome in Patients with Chronic Kidney Disease. <i>Kidney and Blood Pressure Research</i> , 2021, 46, 460-468.	2.0	4
36	Coronary artery calcification in Korean patients with incident dialysis. <i>Hemodialysis International</i> , 2017, 21, 367-374.	0.9	3

#	ARTICLE	IF	CITATIONS
37	Efficacy and Safety of CKD-11101 (Proposed Biosimilar of Darbepoetin-Alfa) Compared with Darbepoetin-Alfa in Patients on Hemodialysis: A Randomized, Double-Blinded, Parallel-Group Phase III Study. <i>BioDrugs</i> , 2020, 34, 99-110.	4.6	2
38	Elevated levels of soluble ST2 but not galectin-3 are associated with increased risk of mortality in hemodialysis patients. <i>Kidney Research and Clinical Practice</i> , 2021, 40, 109-119.	2.2	2
39	Atherosclerotic Renovascular Hypertension : Lessons from Recent Clinical Studies. <i>Electrolyte and Blood Pressure</i> , 2010, 8, 87.	1.8	1
40	Moderateâ€“Vigorous Physical Activity and Clinical Outcomes in Adults with Nondialysis Chronic Kidney Disease. <i>Journal of Clinical Medicine</i> , 2021, 10, 3365.	2.4	1
41	Volume Control by Using the Body Composition Monitor in a Puerperal Patient on Hemodialysis. <i>Electrolyte and Blood Pressure</i> , 2011, 9, 63.	1.8	0
42	Association between the transtubular potassium gradient and progression of chronic kidney disease: results from KNOW-CKD. <i>Journal of Nephrology</i> , 2021, 34, 2063-2072.	2.0	0