

Vivienne A Russell

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1274474/publications.pdf>

Version: 2024-02-01

49
papers

2,585
citations

257357

24
h-index

206029

48
g-index

50
all docs

50
docs citations

50
times ranked

2887
citing authors

#	ARTICLE	IF	CITATIONS
1	Differential effects of social isolation rearing on glutamate- and GABA-stimulated noradrenaline release in the rat prefrontal cortex and hippocampus. <i>European Neuropsychopharmacology</i> , 2020, 36, 111-120.	0.3	10
2	Maternal separation stress reduced prenatal-ethanol-induced increase in exploratory behaviour and extracellular signal-regulated kinase activity. <i>Behavioural Brain Research</i> , 2019, 356, 470-482.	1.2	5
3	Early-Ethanol Exposure Induced Region-Specific Changes in Metabolic Proteins in the Rat Brain: A Proteomics Study. <i>Journal of Molecular Neuroscience</i> , 2018, 65, 277-288.	1.1	8
4	Epigenetics: a link between addiction and social environment. <i>Cellular and Molecular Life Sciences</i> , 2017, 74, 2735-2747.	2.4	50
5	Early ethanol exposure and vincocetine treatment alter learning and memory related proteins in the rat hippocampus and prefrontal cortex. <i>Journal of Neuroscience Research</i> , 2017, 95, 1204-1215.	1.3	24
6	Notes on the Recent History of Neuroscience in Africa. <i>Frontiers in Neuroanatomy</i> , 2017, 11, 96.	0.9	9
7	<i>Searsia chirindensis</i> reverses the potentiating effect of prenatal stress on the development of febrile seizures and decreased plasma interleukin-1 β levels. <i>Neuroscience Research</i> , 2016, 103, 54-58.	1.0	7
8	Neuroenergetics. <i>Current Directions in Psychological Science</i> , 2016, 25, 124-129.	2.8	13
9	Developmental stress elicits preference for methamphetamine in the spontaneously hypertensive rat model of attention-deficit/hyperactivity disorder. <i>Behavioral and Brain Functions</i> , 2016, 12, 18.	1.4	5
10	Genetic predisposition and early life experience interact to determine glutamate transporter (GLT1) and solute carrier family 12 member 5 (KCC2) levels in rat hippocampus. <i>Metabolic Brain Disease</i> , 2016, 31, 169-182.	1.4	10
11	Genetically determined differences in noradrenergic function: The spontaneously hypertensive rat model. <i>Brain Research</i> , 2016, 1641, 291-305.	1.1	9
12	Effect of cocaine on striatal dopamine clearance in a rat model of developmental stress and attention-deficit/hyperactivity disorder. <i>Stress</i> , 2016, 19, 78-82.	0.8	4
13	Proteomic analysis of maternal separation-induced striatal changes in a rat model of ADHD: The spontaneously hypertensive rat. <i>Journal of Neuroscience Methods</i> , 2015, 252, 64-74.	1.3	9
14	Impaired Energy Metabolism and Disturbed Dopamine and Glutamate Signalling in the Striatum and Prefrontal Cortex of the Spontaneously Hypertensive Rat Model of Attention-Deficit Hyperactivity Disorder. <i>Journal of Molecular Neuroscience</i> , 2015, 56, 696-707.	1.1	20
15	Tat-induced histopathological alterations mediate hippocampus-associated behavioural impairments in rats. <i>Behavioral and Brain Functions</i> , 2015, 11, 3.	1.4	22
16	Nicotine-stimulated release of [3 H]norepinephrine is reduced in the hippocampus of an animal model of attention-deficit/hyperactivity disorder, the spontaneously hypertensive rat. <i>Brain Research</i> , 2014, 1572, 1-10.	1.1	9
17	The interaction between stress and exercise, and its impact on brain function. <i>Metabolic Brain Disease</i> , 2014, 29, 255-260.	1.4	18
18	Effect of diet on brain metabolites and behavior in spontaneously hypertensive rats. <i>Behavioural Brain Research</i> , 2014, 270, 240-247.	1.2	15

#	ARTICLE	IF	CITATIONS
19	Maternal separation increases GABAA receptor-mediated modulation of norepinephrine release in the hippocampus of a rat model of ADHD, the spontaneously hypertensive rat. <i>Brain Research</i> , 2013, 1497, 23-31.	1.1	45
20	Evidence for reduced tonic levels of GABA in the hippocampus of an animal model of ADHD, the spontaneously hypertensive rat. <i>Brain Research</i> , 2013, 1541, 52-60.	1.1	26
21	A behavioral neuroenergetics theory of ADHD. <i>Neuroscience and Biobehavioral Reviews</i> , 2013, 37, 625-657.	2.9	73
22	Synergistic tonic and phasic activity of the locus coeruleus norepinephrine (LC-NE) arousal system is required for optimal attentional performance. <i>Metabolic Brain Disease</i> , 2012, 27, 267-274.	1.4	118
23	Maternal separation enhances object location memory and prevents exercise-induced MAPK/ERK signalling in adult Sprague-Dawley rats. <i>Metabolic Brain Disease</i> , 2012, 27, 377-385.	1.4	25
24	Exercise normalizes altered expression of proteins in the ventral hippocampus of rats subjected to maternal separation. <i>Experimental Physiology</i> , 2012, 97, 239-247.	0.9	30
25	The impact of voluntary exercise on relative telomere length in a rat model of developmental stress. <i>BMC Research Notes</i> , 2012, 5, 697.	0.6	17
26	Clozapine decreases exploratory activity and increases anxiety-like behaviour in the Wistar-Kyoto rat but not the spontaneously hypertensive rat model of attention-deficit/hyperactivity disorder. <i>Brain Research</i> , 2012, 1467, 91-103.	1.1	23
27	Maternal separation affects dopamine transporter function in the Spontaneously Hypertensive Rat: An in vivo electrochemical study. <i>Behavioral and Brain Functions</i> , 2011, 7, 49.	1.4	30
28	Effects of early life trauma are dependent on genetic predisposition: a rat study. <i>Behavioral and Brain Functions</i> , 2011, 7, 11.	1.4	46
29	Effect of exercise on dopamine neuron survival in prenatally stressed rats. <i>Metabolic Brain Disease</i> , 2009, 24, 525-539.	1.4	37
30	Effect of exercise on learning and memory in a rat model of developmental stress. <i>Metabolic Brain Disease</i> , 2009, 24, 643-657.	1.4	65
31	Effect of exercise on synaptophysin and calcium/calmodulin-dependent protein kinase levels in prefrontal cortex and hippocampus of a rat model of developmental stress. <i>Metabolic Brain Disease</i> , 2009, 24, 701-709.	1.4	28
32	Increased glutamate-stimulated release of dopamine in substantia nigra of a rat model for attention-deficit/hyperactivity disorder—lack of effect of methylphenidate. <i>Metabolic Brain Disease</i> , 2009, 24, 599-613.	1.4	27
33	The spontaneously hypertensive rat model of ADHD — The importance of selecting the appropriate reference strain. <i>Neuropharmacology</i> , 2009, 57, 619-626.	2.0	176
34	Cross-fostering does not alter the neurochemistry or behavior of spontaneously hypertensive rats. <i>Behavioral and Brain Functions</i> , 2009, 5, 24.	1.4	37
35	Triggering endogenous neuroprotective processes through exercise in models of dopamine deficiency. <i>Parkinsonism and Related Disorders</i> , 2009, 15, S42-S45.	1.1	94
36	Effect of ageing on Ca ²⁺ uptake via NMDA receptors into barrel cortex slices of spontaneously hypertensive rats. <i>Metabolic Brain Disease</i> , 2008, 23, 1-8.	1.4	5

#	ARTICLE	IF	CITATIONS
37	Development of a mild prenatal stress rat model to study long term effects on neural function and survival. <i>Metabolic Brain Disease</i> , 2008, 23, 31-42.	1.4	14
38	Effects of development and dopamine depletion on striatal NMDA receptor-mediated calcium uptake. <i>Metabolic Brain Disease</i> , 2008, 23, 9-30.	1.4	2
39	Response variability in Attention-Deficit/Hyperactivity Disorder: a neuronal and glial energetics hypothesis. <i>Behavioral and Brain Functions</i> , 2006, 2, 30.	1.4	202
40	Stress reduces the neuroprotective effect of exercise in a rat model for Parkinson's disease. <i>Behavioural Brain Research</i> , 2005, 165, 210-220.	1.2	57
41	Animal models of attention-deficit hyperactivity disorder. <i>Behavioral and Brain Functions</i> , 2005, 1, 9.	1.4	189
42	Rodent Models of Attention-Deficit/Hyperactivity Disorder. <i>Biological Psychiatry</i> , 2005, 57, 1239-1247.	0.7	409
43	NMDA-stimulated Ca ²⁺ uptake into barrel cortex slices of spontaneously hypertensive rats. <i>Metabolic Brain Disease</i> , 2001, 16, 133-141.	1.4	21
44	Increased AMPA receptor function in slices containing the prefrontal cortex of spontaneously hypertensive rats. , 2001, 16, 143-149.		42
45	The nucleus accumbens motor-limbic interface of the spontaneously hypertensive rat as studied in vitro by the superfusion slice technique. <i>Neuroscience and Biobehavioral Reviews</i> , 2000, 24, 133-136.	2.9	77
46	Development of a method to evaluate glutamate receptor function in rat barrel cortex slices. <i>Metabolic Brain Disease</i> , 2000, 15, 305-314.	1.4	6
47	Increased noradrenergic activity in prefrontal cortex slices of an animal model for attention-deficit hyperactivity disorder " the spontaneously hypertensive rat. <i>Behavioural Brain Research</i> , 2000, 117, 69-74.	1.2	99
48	Differences between electrically-, ritalin- and d-amphetamine-stimulated release of [3H]dopamine from brain slices suggest impaired vesicular storage of dopamine in an animal model of Attention-Deficit Hyperactivity Disorder. <i>Behavioural Brain Research</i> , 1998, 94, 163-171.	1.2	136
49	Altered dopaminergic function in the prefrontal cortex, nucleus accumbens and caudate-putamen of an animal model of attention-deficit hyperactivity disorder " the spontaneously hypertensive rat. <i>Brain Research</i> , 1995, 676, 343-351.	1.1	182