

# Qi Xu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/127685/publications.pdf>

Version: 2024-02-01

14  
papers

365  
citations

1040056

9  
h-index

1281871

11  
g-index

14  
all docs

14  
docs citations

14  
times ranked

593  
citing authors

#	ARTICLE	IF	CITATIONS
1	Epigallocatechin Gallate Has a Neurorescue Effect in a Mouse Model of Parkinson Disease. <i>Journal of Nutrition</i> , 2017, 147, 1926-1931.	2.9	111
2	Neuroprotective effect of the natural iron chelator, phytic acid in a cell culture model of Parkinson's disease. <i>Toxicology</i> , 2008, 245, 101-108.	4.2	107
3	Total flavonoids of astragalus attenuates experimental autoimmune encephalomyelitis by suppressing the activation and inflammatory responses of microglia via JNK/AKT/NF $\kappa$ B signaling pathway. <i>Phytomedicine</i> , 2021, 80, 153385.	5.3	27
4	Luteolin-7-O-glucoside protects dopaminergic neurons by activating estrogen-receptor-mediated signaling pathway in MPTP-induced mice. <i>Toxicology</i> , 2019, 426, 152256.	4.2	26
5	Phytic Acid Protects against 6-Hydroxydopamine-Induced Dopaminergic Neuron Apoptosis in Normal and Iron Excess Conditions in a Cell Culture Model. <i>Parkinson's Disease</i> , 2011, 2011, 1-6.	1.1	24
6	The Immunoregulatory Effects of Traditional Chinese Medicine on Psoriasis via its Action on Interleukin: Advances and Considerations. <i>The American Journal of Chinese Medicine</i> , 2018, 46, 739-750.	3.8	20
7	Hepcidin Plays a Key Role in 6-OHDA Induced Iron Overload and Apoptotic Cell Death in a Cell Culture Model of Parkinson's Disease. <i>Parkinson's Disease</i> , 2016, 2016, 1-7.	1.1	18
8	A-Type Cinnamon Procyanidin Oligomers Protect Against 1-Methyl-4-Phenyl-1,2,3,6-Tetrahydropyridine-Induced Neurotoxicity in Mice Through Inhibiting the P38 Mitogen-Activated Protein Kinase/P53/BCL-2 Associated X Protein Signaling Pathway. <i>Journal of Nutrition</i> , 2020, 150, 1731-1737.	2.9	12
9	Epigallocatechin Gallate Protects against TNF $\alpha$ - or H <sub>2</sub> O <sub>2</sub> - Induced Apoptosis by Modulating Iron Related Proteins in a Cell Culture Model. <i>International Journal for Vitamin and Nutrition Research</i> , 2018, 88, 158-165.	1.5	10
10	Acute and chronic toxicity of a polyherbal preparation "Jueyin granules. <i>BMC Complementary and Alternative Medicine</i> , 2018, 18, 148.	3.7	9
11	Neuroprotective Effects of B-Type Cinnamon Procyanidin Oligomers on MPP <sup>+</sup> -Induced Apoptosis in a Cell Culture Model of Parkinson's Disease. <i>Molecules</i> , 2021, 26, 6422.	3.8	1
12	Hepcidin plays a key role in 6-OHDA induced iron overload and apoptotic cell death in a cell culture model of Parkinson's disease (1038.2). <i>FASEB Journal</i> , 2014, 28, 1038.2.	0.5	0
13	Calcein's Quenching In Vitro Method for Assessing Dietary Iron Bioavailability. <i>FASEB Journal</i> , 2015, 29, LB336.	0.5	0
14	( $\alpha$ )-Epigallocatechin-3-Gallate Protects against TNF alpha and Hydrogen Peroxide Induced Apoptosis in a Cell Culture Model of Parkinson's Disease. <i>FASEB Journal</i> , 2015, 29, 922.9.	0.5	0