Qi Xu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/127685/publications.pdf

Version: 2024-02-01

1281871 1040056 14 365 9 11 citations h-index g-index papers 593 14 14 14 all docs docs citations times ranked citing authors

#	Article	IF	CITATIONS
1	Total flavonoids of astragalus attenuates experimental autoimmune encephalomyelitis by suppressing the activation and inflammatory responses of microglia via JNK/AKT/NFΰB signaling pathway. Phytomedicine, 2021, 80, 153385.	5.3	27
2	Neuroprotective Effects of B-Type Cinnamon Procyanidin Oligomers on MPP+-Induced Apoptosis in a Cell Culture Model of Parkinson's Disease. Molecules, 2021, 26, 6422.	3.8	1
3	A-Type Cinnamon Procyanidin Oligomers Protect Against 1-Methyl-4-Phenyl-1,2,3,6-Tetrahydropyridine-Induced Neurotoxicity in Mice Through Inhibiting the P38 Mitogen-Activated Protein Kinase/P53/BCL-2 Associated X Protein Signaling Pathway. Journal of Nutrition. 2020. 150. 1731-1737.	2.9	12
4	Luteolin-7-O-glucoside protects dopaminergic neurons by activating estrogen-receptor-mediated signaling pathway in MPTP-induced mice. Toxicology, 2019, 426, 152256.	4.2	26
5	The Immunoregulatory Effects of Traditional Chinese Medicine on Psoriasis via its Action on Interleukin: Advances and Considerations. The American Journal of Chinese Medicine, 2018, 46, 739-750.	3.8	20
6	Acute and chronic toxicity of a polyherbal preparation – Jueyin granules. BMC Complementary and Alternative Medicine, 2018, 18, 148.	3.7	9
7	Epigallocatechin Gallate Protects against TNFα- or H2O2- Induced Apoptosis by Modulating Iron Related Proteins in a Cell Culture Model. International Journal for Vitamin and Nutrition Research, 2018, 88, 158-165.	1.5	10
8	Epigallocatechin Gallate Has a Neurorescue Effect in a Mouse Model of Parkinson Disease. Journal of Nutrition, 2017, 147, 1926-1931.	2.9	111
9	Hepcidin Plays a Key Role in 6-OHDA Induced Iron Overload and Apoptotic Cell Death in a Cell Culture Model of Parkinson's Disease. Parkinson's Disease, 2016, 2016, 1-7.	1.1	18
10	Calcein's Quenching In Vitro Method for Assessing Dietary Iron Bioavailability. FASEB Journal, 2015, 29, LB336.	0.5	0
11	(â^')â€Epigallocatechinâ€3â€Gallate Protects against TNF alpha and Hydrogen Peroxide Induced Apoptosis in a Cell Culture Model of Parkinson's Disease. FASEB Journal, 2015, 29, 922.9.	0.5	O
12	Hepcidin plays a key role in 6â€OHDA induced iron overload and apoptotic cell death in a cell culture model of Parkinson's disease (1038.2). FASEB Journal, 2014, 28, 1038.2.	0.5	0
13	Phytic Acid Protects against 6-Hydroxydopamine-Induced Dopaminergic Neuron Apoptosis in Normal and Iron Excess Conditions in a Cell Culture Model. Parkinson's Disease, 2011, 2011, 1-6.	1.1	24
14	Neuroprotective effect of the natural iron chelator, phytic acid in a cell culture model of Parkinson's disease. Toxicology, 2008, 245, 101-108.	4.2	107