

# Murtaza Emre Durakoglugil

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/128476/publications.pdf>

Version: 2024-02-01

74  
papers

1,302  
citations

516710

16  
h-index

395702

33  
g-index

75  
all docs

75  
docs citations

75  
times ranked

2510  
citing authors

#	ARTICLE	IF	CITATIONS
1	An echocardiographic predictor of contrast-induced nephropathy following percutaneous coronary intervention in acute coronary syndrome: aortic valve sclerosis. <i>Scandinavian Cardiovascular Journal</i> , 2020, 54, 227-231.	1.2	0
2	The relationship between total atrial conduction time and left atrial global strain in patients with psoriasis vulgaris. <i>Archives of Medical Science</i> , 2019, 15, 865-871.	0.9	4
3	The role of serum bilirubin levels in determining venous thromboembolism. <i>Journal of Vascular Surgery: Venous and Lymphatic Disorders</i> , 2019, 7, 635-639.	1.6	8
4	Epicardial Adipose Tissue Predicts Carotid Intima-Media Thickness Independently of Body Mass Index and Waist Circumference. <i>Acta Cardiologica Sinica</i> , 2019, 35, 32-41.	0.2	8
5	What have we learned from Turkish familial hypercholesterolemia registries (A-HIT1 and A-HIT2)? <i>Atherosclerosis</i> , 2018, 277, 341-346.	0.8	15
6	The Relationship between Serum Apelin Levels and the Severity of Calcific Aortic Stenosis. <i>Acta Cardiologica Sinica</i> , 2018, 34, 259-266.	0.2	10
7	The effect of epicardial adipose tissue thickness with irritable bowel syndrome. <i>JPMA the Journal of the Pakistan Medical Association</i> , 2018, 68, 1456-1460.	0.2	1
8	Cardiovascular Efficacy and Safety of Bococizumab in High-Risk Patients. <i>New England Journal of Medicine</i> , 2017, 376, 1527-1539.	27.0	510
9	Prolonged QRS duration on surface electrocardiogram is associated with left ventricular restrictive filling pattern. <i>Interventional Medicine &amp; Applied Science</i> , 2017, 9, 9-14.	0.2	2
10	An increase in epicardial adipose tissue is strongly associated with carotid-intima media thickness and atherosclerotic plaque, but LDL only with the plaque. <i>Anatolian Journal of Cardiology</i> , 2017, 17, 56-63.	0.9	17
11	Hair whitening and obesity are independently related to ascending aorta dilatation in young-middle aged men. <i>Åstanbul Kuzey Klinikleri</i> , 2017, 6, 33-39.	0.3	2
12	The role of urotensin II and atherosclerotic risk factors in patients with slow coronary flow. <i>Interventional Medicine &amp; Applied Science</i> , 2016, 8, 158-163.	0.2	5
13	Total Bilirubin Levels Predict Subclinical Atherosclerosis in Patients With Prediabetes. <i>Angiology</i> , 2016, 67, 909-915.	1.8	20
14	Multiple simultaneous venous and arterial thromboses in a patient with factor V Leiden disorder: Detection by multislice computed tomography. <i>Interventional Medicine &amp; Applied Science</i> , 2015, 7, 38-41.	0.2	1
15	Relation of Angiographic Thrombus Burden with Severity of Coronary Artery Disease in Patients with ST Segment Elevation Myocardial Infarction. <i>Medical Science Monitor</i> , 2015, 21, 3540-3546.	1.1	16
16	Evaluation of Para- and Perirenal Fat Thickness and its Association with Metabolic Disorders in Polycystic Ovary Syndrome. <i>Endocrine Practice</i> , 2015, 21, 878-886.	2.1	8
17	The relationship of plasma catestatin concentrations with metabolic and vascular parameters in untreated hypertensive patients: Influence on high-density lipoprotein cholesterol. <i>Anatolian Journal of Cardiology</i> , 2015, 15, 577-585.	0.9	27
18	Systolic murmur in a young man who had previous ventricular septal defect repair: The double-chambered right ventricle. <i>Indian Heart Journal</i> , 2015, 67, 482-484.	0.5	2

#	ARTICLE	IF	CITATIONS
19	The relationship between epicardial adipose tissue and P wave and QT dispersions. <i>Türk Kardiyoloji Dernegi Arsivi</i> , 2015, 43, 621-9.	0.5	7
20	Heart rate variability can be affected by gender, blood pressure, and insulin resistance--author's reply. <i>Anatolian Journal of Cardiology</i> , 2015, 15, 263.	0.9	0
21	Bilirubin Level is Associated with Left Ventricular Hypertrophy Independent of Blood Pressure in Previously Untreated Hypertensive Patients. <i>Korean Circulation Journal</i> , 2014, 44, 336.	1.9	5
22	The effect of irritable bowel syndrome on carotid intima-media thickness, pulse wave velocity, and heart rate variability. <i>Anatolian Journal of Cardiology</i> , 2014, 14, 525-530.	0.4	5
23	Reply. <i>Muscle and Nerve</i> , 2014, 50, 303-303.	2.2	0
24	Characterization of Echocardiographic Measures of Cardiac Structure and Function in Healthy Octogenarians. <i>Echocardiography</i> , 2014, 31, 1062-1070.	0.9	5
25	The Relation Between Carotid Intima Media Thickness and Serum Osteoprotegerin Levels in Nonalcoholic Fatty Liver Disease. <i>Metabolic Syndrome and Related Disorders</i> , 2014, 12, 283-289.	1.3	12
26	Corrigendum to 'The effect of combined hyperbaric oxygen and iloprost treatment on the prevention of spinal cord ischaemia-reperfusion injury: an experimental study' [ <i>Eur J Cardiothorac Surg</i> 2013;44:e332-e340]. <i>European Journal of Cardio-thoracic Surgery</i> , 2014, 45, 209-209.	1.4	0
27	Arterial Stiffness Measured Via Carotid Femoral Pulse Wave Velocity Is Associated With Disease Severity in COPD. <i>Respiratory Care</i> , 2014, 59, 274-280.	1.6	37
28	Epicardial adipose tissue thickness and plasma homocysteine in patients with metabolic syndrome and normal coronary arteries. <i>Diabetology and Metabolic Syndrome</i> , 2014, 6, 62.	2.7	11
29	The Effect of Different $\beta$ -Blockers on Vascular Graft Nitric Oxide Levels: Comparison of Nebivolol Versus Metoprolol. <i>European Journal of Vascular and Endovascular Surgery</i> , 2014, 47, 204-208.	1.5	8
30	Successful thrombolysis using recombinant tissue plasminogen activator in cases of severe pulmonary embolism with mobile thrombi in the right atrium. <i>Interventional Medicine &amp; Applied Science</i> , 2014, 6, 89-92.	0.2	6
31	Retrieval of embolized tip of port catheter from branch of right pulmonary artery using a macro snare catheter. <i>Interventional Medicine &amp; Applied Science</i> , 2014, 6, 93-95.	0.2	4
32	Relationship between fragmented QRS complexes and left ventricular systolic and diastolic functions. <i>Herz</i> , 2013, 38, 665-670.	1.1	16
33	Cardiac syncope due to pain. <i>Herz</i> , 2013, 38, 683-684.	1.1	0
34	Aortic valve sclerosis is a high predictive marker of systemic endothelial dysfunction in hypertensive patients. <i>Herz</i> , 2013, 38, 915-921.	1.1	11
35	Increased serum resistin levels in patients with coronary slow-flow phenomenon. <i>Herz</i> , 2013, 38, 773-778.	1.1	14
36	Elevated serum YKL-40 level predicts myocardial reperfusion and in-hospital MACE in patients with STEMI. <i>Herz</i> , 2013, 38, 202-209.	1.1	15

#	ARTICLE	IF	CITATIONS
37	Increased Pulse Wave Velocity and Carotid Intima-Media Thickness in Patients with Ulcerative Colitis. <i>Digestive Diseases and Sciences</i> , 2013, 58, 2293-2300.	2.3	26
38	Increased pulse-wave velocity and carotid intima-media thickness in patients with lower urinary tract symptoms. <i>Scandinavian Journal of Urology</i> , 2013, 47, 393-398.	1.0	12
39	Non-dipping pattern in untreated hypertensive patients is related to increased pulse wave velocity independent of raised nocturnal blood pressure. <i>Blood Pressure</i> , 2013, 22, 34-38.	1.5	32
40	Effect of epicardial adipose tissue on diastolic functions and left atrial dimension in untreated hypertensive patients with normal systolic function. <i>Journal of Cardiology</i> , 2013, 61, 359-364.	1.9	31
41	Decreased heart rate variability in irritable bowel syndrome has no effect on carotid intima-media thickness and pulse wave velocity. <i>Journal of the American College of Cardiology</i> , 2013, 62, C208.	2.8	0
42	The effect of combined hyperbaric oxygen and iloprost treatment on the prevention of spinal cord ischaemia-reperfusion injury: an experimental study. <i>European Journal of Cardio-thoracic Surgery</i> , 2013, 44, e332-e340.	1.4	11
43	Epicardial adipose tissue is independently associated with increased left ventricular mass in untreated hypertensive patients: an observational study. <i>Anatolian Journal of Cardiology</i> , 2013, 13, 320-7.	0.4	6
44	YKL-40 levels in patients with coronary artery ectasia. <i>Anatolian Journal of Cardiology</i> , 2013, 14, 97-8.	0.4	3
45	Increased YKL-40 levels in patients with isolated coronary artery ectasia: an observational study. <i>Anatolian Journal of Cardiology</i> , 2013, 13, 465-70.	0.4	13
46	The risk of developing AF after cardiac surgery. <i>Anatolian Journal of Cardiology</i> , 2013, 13, 502.	0.4	0
47	YKL-40 as new cardiac biomarker. <i>Anatolian Journal of Cardiology</i> , 2013, 14, 98-9.	0.4	1
48	The independent relationship of epicardial adipose tissue with carotid intima-media thickness and endothelial functions. <i>Blood Pressure Monitoring</i> , 2013, 18, 85-93.	0.8	26
49	Increased Carotid and Brachial Intima-Media Thickness is Related to Diffuse Coronary Involvement Rather Than Focal Lesions. <i>Angiology</i> , 2013, 64, 356-363.	1.8	5
50	Increased pulse wave velocity and carotid intima-media thickness in patients with carpal tunnel syndrome. <i>Muscle and Nerve</i> , 2013, 47, 872-877.	2.2	6
51	Premature Hair Whitening is an Independent Predictor of Carotid Intima-media Thickness in Young and Middle-aged Men. <i>Internal Medicine</i> , 2013, 52, 29-36.	0.7	18
52	A rare complication: Undeatable balloon of the stent. <i>Interventional Medicine &amp; Applied Science</i> , 2013, 5, 43-45.	0.2	5
53	The degree of premature hair graying as an independent risk marker for coronary artery disease: a predictor of biological age rather than chronological age. <i>Anatolian Journal of Cardiology</i> , 2012, 12, 457-63.	0.4	35
54	Increased circulating soluble CD40 levels in patients with slow coronary flow phenomenon: an observational study. <i>Anatolian Journal of Cardiology</i> , 2012, 13, 39-44.	0.4	9

#	ARTICLE	IF	CITATIONS
55	Fragmented QRS may predict postoperative atrial fibrillation in patients undergoing isolated coronary artery bypass graft surgery. <i>Anatolian Journal of Cardiology</i> , 2012, 12, 576-83.	0.4	14
56	Independent determinants of ascending aortic dilatation in hypertensive patients. <i>Blood Pressure Monitoring</i> , 2012, 17, 223-230.	0.8	12
57	Absence and Resolution of Fragmented QRS Predict Reversible Myocardial Ischemia With Higher Probability of ST Segment Resolution in Patients With ST Segment Elevation Myocardial Infarction. <i>Korean Circulation Journal</i> , 2012, 42, 674.	1.9	14
58	Relationship of fragmented QRS complexes with inadequate coronary collaterals in patients with chronic total occlusion. <i>Journal of Cardiovascular Medicine</i> , 2012, 13, 499-504.	1.5	8
59	Increased Epicardial Adipose Tissue in Patients with Isolated Coronary Artery Ectasia. <i>Internal Medicine</i> , 2012, 51, 833-838.	0.7	12
60	Increased Epicardial Adipose Tissue Thickness Is Correlated with Ascending Aortic Diameter. <i>Tohoku Journal of Experimental Medicine</i> , 2012, 226, 183-190.	1.2	11
61	Increased Serum Bilirubin Level is Related to Good Collateral Development in Patients with Chronic Total Coronary Occlusion. <i>Internal Medicine</i> , 2012, 51, 249-255.	0.7	24
62	A rare combination: Single coronary artery originating from the left sinus of valsalva coexisting with perimembranous ventricular septal defect. <i>Journal of Cardiology Cases</i> , 2012, 6, e39-e41.	0.5	0
63	Elevated serum osteoprotegerin levels predict in-hospital major adverse cardiac events in patients with ST elevation myocardial infarction. <i>Journal of Cardiology</i> , 2012, 60, 355-360.	1.9	11
64	Relationship of fragmented QRS with prognostic markers and in-hospital MACE in patients undergoing CABG. <i>Scandinavian Cardiovascular Journal</i> , 2012, 46, 107-113.	1.2	12
65	Increased pulse wave velocity in patients with panic disorder: Independent vascular influence of panic disorder on arterial stiffness. <i>Journal of Psychosomatic Research</i> , 2012, 73, 145-148.	2.6	28
66	The Independent Relationship of Systemic Inflammation With Fragmented QRS Complexes in Patients With Acute Coronary Syndromes. <i>Korean Circulation Journal</i> , 2012, 42, 449.	1.9	15
67	Increased plasminogen activator inhibitor-1 levels in patients with isolated coronary artery ectasia. <i>Journal of Thrombosis and Thrombolysis</i> , 2012, 33, 120-123.	2.1	11
68	The importance of fragmented QRS complexes in prediction of myocardial infarction and reperfusion parameters in patients undergoing primary percutaneous coronary intervention. <i>Turk Kardiyoloji Dernegi Arsivi</i> , 2012, 40, 213-222.	0.5	24
69	A rare sign of ischemia during exercise ECG: PR interval lengthening in the recovery period. <i>Turk Kardiyoloji Dernegi Arsivi</i> , 2012, 40, 719-722.	0.5	0
70	Distributions of interleukin-6 (IL-6) promoter and metallothionein 2A (MT2A) core promoter region gene polymorphisms and their associations with aging in Turkish population. <i>Archives of Gerontology and Geriatrics</i> , 2011, 53, 354-358.	3.0	28
71	The Success of Clopidogrel Treatment for Recurrent Coronary Thrombosis in a Patient With Aortic and Mitral Valve Replacement. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2008, 14, 492-493.	1.7	0
72	Heart fatty acid binding protein and myoglobin after reperfusion of acute myocardial infarction. <i>Acta Cardiologica</i> , 2007, 62, 473-478.	0.9	6

#	ARTICLE	IF	CITATIONS
73	Effect of Metoprolol on Heart Rate Variability in Symptomatic Patients With Mitral Valve Prolapse. American Journal of Cardiology, 2007, 99, 1568-1570.	1.6	10
74	High Output Heart Failure 8 Months After an Acquired Arteriovenous Fistula. International Heart Journal, 2003, 44, 805-809.	0.6	11