Peng-Fan Yang

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1287337/publications.pdf

Version: 2024-02-01

	1307594	1372567
252	7	10
citations	h-index	g-index
1.1	1.1	401
11	11	401
docs citations	times ranked	citing authors
	citations 11	252 7 citations h-index 11 11

#	Article	IF	Citations
1	Effects of scorpion venom heat-resistant peptide on the hippocampal neurons of kainic acid-induced epileptic rats. Brazilian Journal of Medical and Biological Research, 2021, 54, e10717.	1.5	6
2	Neuronavigation-assisted surgical treatments for medically refractory epilepsy: Single-hospital experience with 4 surgical approaches. Clinical Neurology and Neurosurgery, 2019, 182, 148-151.	1.4	0
3	Surgical Treatment and Long-Term Outcome of Cerebral Cavernous Malformations-Related Epilepsy in Pediatric Patients. Neuropediatrics, 2018, 49, 173-179.	0.6	7
4	Surgical Management and Long-Term Seizure Outcome After Surgery for Temporal Lobe Epilepsy Associated with Cerebral Cavernous Malformations. World Neurosurgery, 2018, 110, e659-e670.	1.3	13
5	Three-dimensional intracranial EEG monitoring in presurgical assessment of MRI-negative frontal lobe epilepsy. Medicine (United States), 2016, 95, e5192.	1.0	4
6	Keyhole epilepsy surgery: corticoamygdalohippocampectomy for mesial temporal sclerosis. Neurosurgical Review, 2016, 39, 99-108.	2.4	38
7	Intractable occipital lobe epilepsy: clinical characteristics, surgical treatment, and a systematic review of the literature. Acta Neurochirurgica, 2015, 157, 63-75.	1.7	34
8	Disconnective surgery in posterior quadrantic epilepsy: a series of 12 paediatric patients. Epileptic Disorders, 2014, 16, 296-304.	1.3	28
9	Long-term epilepsy surgery outcomes in patients with PET-positive, MRI-negative temporal lobe epilepsy. Epilepsy and Behavior, 2014, 41, 91-97.	1.7	77
10	Intracranial electroencephalography with subdural and/or depth electrodes in children with epilepsy: Techniques, complications, and outcomes. Epilepsy Research, 2014, 108, 1662-1670.	1.6	31
11	Outcome after anterior callosal section that spares the splenium in pediatric patients with drop attacks. Epilepsy and Behavior, 2014, 36, 47-52.	1.7	14