

# Hans Gerd Nothwang

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1290775/publications.pdf>

Version: 2024-02-01

75  
papers

3,972  
citations

147801

31  
h-index

118850

62  
g-index

77  
all docs

77  
docs citations

77  
times ranked

4402  
citing authors

#	ARTICLE	IF	CITATIONS
1	Expression pattern of cochlear microRNAs in the mammalian auditory hindbrain. <i>Cell and Tissue Research</i> , 2021, 383, 655-666.	2.9	7
2	Structural changes in the extracellular loop 2 of the murine KCC2 potassium chloride cotransporter modulate ion transport. <i>Journal of Biological Chemistry</i> , 2021, 296, 100793.	3.4	5
3	The Lbx1 lineage differentially contributes to inhibitory cell types of the dorsal cochlear nucleus, a cerebellum-like structure, and the cerebellum. <i>Journal of Comparative Neurology</i> , 2021, 529, 3032-3045.	1.6	6
4	Loss of miR-183/96 Alters Synaptic Strength via Presynaptic and Postsynaptic Mechanisms at a Central Synapse. <i>Journal of Neuroscience</i> , 2021, 41, 6796-6811.	3.6	9
5	Differential expression of <scp>microRNAs</scp> in the developing avian auditory hindbrain. <i>Journal of Comparative Neurology</i> , 2021, 529, 3477-3496.	1.6	2
6	Staurosporine and NEM mainly impair WNK-SPAK/OSR1 mediated phosphorylation of KCC2 and NKCC1. <i>PLoS ONE</i> , 2020, 15, e0232967.	2.5	14
7	Loss of inner hair cell ribbon synapses and auditory nerve fiber regression in Cldn14 knockout mice. <i>Hearing Research</i> , 2020, 391, 107950.	2.0	0
8	miR-96 is required for normal development of the auditory hindbrain. <i>Human Molecular Genetics</i> , 2018, 27, 860-874.	2.9	31
9	Evolution of Endolymph Secretion and Endolymphatic Potential Generation in the Vertebrate Inner Ear. <i>Brain, Behavior and Evolution</i> , 2018, 92, 1-31.	1.7	29
10	Phosphoregulation of the intracellular termini of K <sup>+</sup> -Cl <sup>-</sup> cotransporter 2 (KCC2) enables flexible control of its activity. <i>Journal of Biological Chemistry</i> , 2018, 293, 16984-16993.	3.4	22
11	Differences in molecular mechanisms of K <sup>+</sup> clearance in the auditory sensory epithelium of birds and mammals. <i>Journal of Experimental Biology</i> , 2017, 220, 2701-2705.	1.7	3
12	Activity-dependent formation of a vesicular inhibitory amino acid transporter gradient in the superior olivary complex of NMRI mice. <i>BMC Neuroscience</i> , 2017, 18, 75.	1.9	0
13	Molecular cloning and biochemical characterization of two cation chloride cotransporter subfamily members of <i>Hydra vulgaris</i> . <i>PLoS ONE</i> , 2017, 12, e0179968.	2.5	9
14	Functional Role of $\beta$ -Crystallin N in the Auditory Hindbrain. <i>PLoS ONE</i> , 2016, 11, e0161140.	2.5	5
15	Comparative Analysis of Gene Regulatory Network Components in the Auditory Hindbrain of Mice and Chicken. <i>Brain, Behavior and Evolution</i> , 2016, 88, 161-176.	1.7	8
16	Molecular bases of K <sup>+</sup> secretory cells in the inner ear: shared and distinct features between birds and mammals. <i>Scientific Reports</i> , 2016, 6, 34203.	3.3	18
17	Evolution of mammalian sound localization circuits: A developmental perspective. <i>Progress in Neurobiology</i> , 2016, 141, 1-24.	5.7	44
18	Differential patterns of histone methylase EHMT2 and its catalyzed histone modifications H3K9me1 and H3K9me2 during maturation of central auditory system. <i>Cell and Tissue Research</i> , 2016, 365, 247-264.	2.9	13

#	ARTICLE	IF	CITATIONS
19	Neto2-null mice have impaired GABAergic inhibition and are susceptible to seizures. <i>Frontiers in Cellular Neuroscience</i> , 2015, 9, 368.	3.7	25
20	The emerging framework of mammalian auditory hindbrain development. <i>Cell and Tissue Research</i> , 2015, 361, 33-48.	2.9	20
21	The gene regulatory networks underlying formation of the auditory hindbrain. <i>Cellular and Molecular Life Sciences</i> , 2015, 72, 519-535.	5.4	17
22	KCC2 regulates actin dynamics in dendritic spines via interaction with $\hat{I}^2$ -PIX. <i>Journal of Cell Biology</i> , 2015, 209, 671-686.	5.2	97
23	L-type Calcium Channel Cav1.2 Is Required for Maintenance of Auditory Brainstem Nuclei. <i>Journal of Biological Chemistry</i> , 2015, 290, 23692-23710.	3.4	17
24	KCC2 regulates actin dynamics in dendritic spines via interaction with $\hat{I}^2$ -PIX. <i>Journal of Experimental Medicine</i> , 2015, 212, 2127OIA56.	8.5	0
25	L-Typ-Kalzium-Kanäle im Hörsystem. <i>E-Neuroforum</i> , 2014, 20, 250-257.	0.1	0
26	A Novel Regulatory Locus of Phosphorylation in the C Terminus of the Potassium Chloride Cotransporter KCC2 That Interferes with N-Ethylmaleimide or Staurosporine-mediated Activation <sup>TM</sup> . <i>Journal of Biological Chemistry</i> , 2014, 289, 18668-18679.	3.4	56
27	Central auditory function of deafness genes. <i>Hearing Research</i> , 2014, 312, 9-20.	2.0	24
28	Evolution of the Cation Chloride Cotransporter Family: Ancient Origins, Gene Losses, and Subfunctionalization through Duplication. <i>Molecular Biology and Evolution</i> , 2014, 31, 434-447.	8.9	54
29	Molecular and evolutionary insights into the structural organization of cation chloride cotransporters. <i>Frontiers in Cellular Neuroscience</i> , 2014, 8, 470.	3.7	43
30	Monitoring the native phosphorylation state of plasma membrane proteins from a single mouse cerebellum. <i>Journal of Neuroscience Methods</i> , 2013, 213, 153-164.	2.5	7
31	Evolutionary Conservation of Kv3.1 in the Barn Owl <i>Tyto alba</i> . <i>Brain, Behavior and Evolution</i> , 2013, 81, 187-193.	1.7	1
32	Time-dependent Gene Expression Analysis of the Developing Superior Olivary Complex. <i>Journal of Biological Chemistry</i> , 2013, 288, 25865-25879.	3.4	32
33	Retrocochlear function of the peripheral deafness gene <i>Cacna1d</i> . <i>Human Molecular Genetics</i> , 2012, 21, 3896-3909.	2.9	45
34	KCC2 transport activity requires the highly conserved L675 in the C-terminal $\hat{I}^2$ 1 strand. <i>Biochemical and Biophysical Research Communications</i> , 2012, 420, 492-497.	2.1	3
35	<i>Egr2::Cre</i> Mediated Conditional Ablation of <i>Dicer</i> Disrupts Histogenesis of Mammalian Central Auditory Nuclei. <i>PLoS ONE</i> , 2012, 7, e49503.	2.5	20
36	Opposite temperature effect on transport activity of KCC2/KCC4 and N(K)CCs in HEK-293 cells. <i>BMC Research Notes</i> , 2011, 4, 526.	1.4	11

#	ARTICLE	IF	CITATIONS
37	Ca <sup>v</sup> 1.3 Calcium Channels Are Required for Normal Development of the Auditory Brainstem. <i>Journal of Neuroscience</i> , 2011, 31, 8280-8294.	3.6	80
38	Differences in the Large Extracellular Loop between the K <sup>+</sup> -Cl <sup>-</sup> Cotransporters KCC2 and KCC4. <i>Journal of Biological Chemistry</i> , 2010, 285, 23994-24002.	3.4	36
39	Approaching clinical proteomics: current state and future fields of application in fluid proteomics. <i>Clinical Chemistry and Laboratory Medicine</i> , 2009, 47, 724-44.	2.3	112
40	Opposite effect of membrane raft perturbation on transport activity of KCC2 and NKCC1. <i>Journal of Neurochemistry</i> , 2009, 111, 321-331.	3.9	41
41	CIP1 is an activator of the K <sup>+</sup> -Cl <sup>-</sup> cotransporter KCC2. <i>Biochemical and Biophysical Research Communications</i> , 2009, 381, 388-392.	2.1	30
42	Subcellular Fractionation of Small Sample Amounts. <i>Springer Protocols</i> , 2009, , 165-170.	0.3	2
43	Enrichment of Brain Plasma Membranes by Affinity Two-Phase Partitioning. <i>Methods in Molecular Biology</i> , 2009, 528, 119-126.	0.9	2
44	Two-Dimensional Separation of Membrane Proteins by 16-BAC-SDS-PAGE. <i>Methods in Molecular Biology</i> , 2009, 528, 269-277.	0.9	12
45	Isolation of Plasma Membranes from the Nervous System by Countercurrent Distribution in Aqueous Polymer Two-Phase Systems. <i>Methods in Molecular Biology</i> , 2009, 564, 335-340.	0.9	1
46	Hypothyroidism impairs chloride homeostasis and onset of inhibitory neurotransmission in developing auditory brainstem and hippocampal neurons. <i>European Journal of Neuroscience</i> , 2008, 28, 2371-2380.	2.6	41
47	Minimal sex differences in gene expression in the rat superior olivary complex. <i>Hearing Research</i> , 2008, 245, 65-72.	2.0	6
48	Neuronale Chloridhomöostase: entwicklungs- und aktivitätsabhängige Regulation von Chloridtransportern. <i>E-Neuroforum</i> , 2008, 14, 148-158.	0.1	0
49	Aqueous polymer two-phase systems: Effective tools for plasma membrane proteomics. <i>Proteomics</i> , 2006, 6, 5409-5417.	2.2	96
50	Comparative gene expression analysis reveals a characteristic molecular profile of the superior olivary complex. <i>The Anatomical Record Part A: Discoveries in Molecular, Cellular, and Evolutionary Biology</i> , 2006, 288A, 409-423.	2.0	10
51	Neuroproteomics – the tasks lying ahead. <i>Electrophoresis</i> , 2006, 27, 2819-2829.	2.4	41
52	Proteomic Analysis of Brain Plasma Membranes Isolated by Affinity Two-phase Partitioning. <i>Molecular and Cellular Proteomics</i> , 2006, 5, 390-400.	3.8	96
53	Oligomerization of KCC2 Correlates with Development of Inhibitory Neurotransmission. <i>Journal of Neuroscience</i> , 2006, 26, 10407-10419.	3.6	223
54	Developmental pattern of three vesicular glutamate transporters in the rat superior olivary complex. <i>Cell and Tissue Research</i> , 2005, 320, 33-50.	2.9	67

#	ARTICLE	IF	CITATIONS
55	A subcellular prefractionation protocol for minute amounts of mammalian cell cultures and tissue. <i>Proteomics</i> , 2005, 5, 35-45.	2.2	95
56	Gene expression profiling of the rat superior olivary complex using serial analysis of gene expression. <i>European Journal of Neuroscience</i> , 2004, 20, 3244-3258.	2.6	15
57	Differential expression pattern of chloride transporters NCC, NKCC2, KCC1, KCC3, KCC4, and AE3 in the developing rat auditory brainstem. <i>Cell and Tissue Research</i> , 2003, 312, 155-165.	2.9	44
58	Protein analysis in the rat auditory brainstem by two-dimensional gel electrophoresis and mass spectrometry. <i>Molecular Brain Research</i> , 2003, 116, 59-69.	2.3	11
59	Efficient Cloning of SAGE Tags by Blunt-End Ligation of Polished Concatemers. <i>BioTechniques</i> , 2003, 34, 692-694.	1.8	9
60	Expression and Function of Chloride Transporters during Development of Inhibitory Neurotransmission in the Auditory Brainstem. <i>Journal of Neuroscience</i> , 2003, 23, 4134-4145.	3.6	173
61	Refinement and delineation of the breakpoint regions of a chromosome 1;22 translocation in a patient with Costello syndrome*. <i>American Journal of Medical Genetics Part A</i> , 2002, 109, 234-237.	2.4	9
62	Mutation of CDH23, encoding a new member of the cadherin gene family, causes Usher syndrome type 1D. <i>Nature Genetics</i> , 2001, 27, 108-112.	21.4	442
63	A translocation breakpoint cluster disrupts the newly defined 3' end of the SNURF-SNRPN transcription unit on chromosome 15. <i>Human Molecular Genetics</i> , 2001, 10, 201-210.	2.9	62
64	Mutations in ARHGEF6, encoding a guanine nucleotide exchange factor for Rho GTPases, in patients with X-linked mental retardation. <i>Nature Genetics</i> , 2000, 26, 247-250.	21.4	329
65	Molecular Cloning of the Critical Region for Glomerulopathy with Fibronectin Deposits (GFND) and Evaluation of Candidate Genes. <i>Genomics</i> , 2000, 68, 127-135.	2.9	14
66	300 million years of conserved synteny between chicken Z and human chromosome 9. <i>Nature Genetics</i> , 1999, 21, 258-259.	21.4	330
67	Cloning and Characterization of UXT, a Novel Gene in Human Xp11, which Is Widely and Abundantly Expressed in Tumor Tissue. <i>Genomics</i> , 1999, 56, 340-343.	2.9	50
68	Lack of large, homozygous deletions of the nephronophthisis 1 region in Joubert syndrome type B. <i>Pediatric Nephrology</i> , 1998, 12, 16-19.	1.7	29
69	Construction of a Gene Map of the Nephronophthisis Type 1 (NPHP1) Region on Human Chromosome 2q12-q13. <i>Genomics</i> , 1998, 47, 276-285.	2.9	26
70	Identification of a Novel Ran Binding Protein 2 Related Gene (RANBP2L1) and Detection of a Gene Cluster on Human Chromosome 2q11-q12. <i>Genomics</i> , 1998, 47, 383-392.	2.9	36
71	Molecular Cloning of the Interleukin-1 Gene Cluster: Construction of an Integrated YAC/PAC Contig and a Partial Transcriptional Map in the Region of Chromosome 2q13. <i>Genomics</i> , 1997, 41, 370-378.	2.9	45
72	A novel gene encoding an SH3 domain protein is mutated in nephronophthisis type 1. <i>Nature Genetics</i> , 1997, 17, 149-153.	21.4	327

#	ARTICLE	IF	CITATIONS
73	Molecular genetic identification of families with juvenile nephronophthisis type 1: Rate of progression to renal failure. <i>Kidney International</i> , 1997, 51, 261-269.	5.2	88
74	cDNA cloning and interferon gamma down-regulation of proteasomal subunits X and Y. <i>Science</i> , 1994, 265, 1231-1234.	12.6	194
75	Phylogenic relationships of the amino acid sequences of prosome (proteasome, MCP) subunits. <i>Molecular Genetics and Genomics</i> , 1994, 245, 769-780.	2.4	51