Thierry Facon

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1295602/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Bortezomib or High-Dose Dexamethasone for Relapsed Multiple Myeloma. New England Journal of Medicine, 2005, 352, 2487-2498.	13.9	2,356
2	Revised International Staging System for Multiple Myeloma: A Report From International Myeloma Working Group. Journal of Clinical Oncology, 2015, 33, 2863-2869.	0.8	1,525
3	Lenalidomide plus Dexamethasone for Relapsed or Refractory Multiple Myeloma. New England Journal of Medicine, 2007, 357, 2123-2132.	13.9	1,365
4	Lenalidomide Maintenance after Stem-Cell Transplantation for Multiple Myeloma. New England Journal of Medicine, 2012, 366, 1782-1791.	13.9	1,022
5	Consensus recommendations for the uniform reporting of clinical trials: report of the International Myeloma Workshop Consensus Panel 1. Blood, 2011, 117, 4691-4695.	0.6	849
6	Melphalan and prednisone plus thalidomide versus melphalan and prednisone alone or reduced-intensity autologous stem cell transplantation in elderly patients with multiple myeloma (IFM 99–06): a randomised trial. Lancet, The, 2007, 370, 1209-1218.	6.3	820
7	Heterogeneity of genomic evolution and mutational profiles in multiple myeloma. Nature Communications, 2014, 5, 2997.	5.8	741
8	Lenalidomide and Dexamethasone in Transplant-Ineligible Patients with Myeloma. New England Journal of Medicine, 2014, 371, 906-917.	13.9	697
9	Daratumumab plus Lenalidomide and Dexamethasone for Untreated Myeloma. New England Journal of Medicine, 2019, 380, 2104-2115.	13.9	684
10	Bortezomib, thalidomide, and dexamethasone with or without daratumumab before and after autologous stem-cell transplantation for newly diagnosed multiple myeloma (CASSIOPEIA): a randomised, open-label, phase 3 study. Lancet, The, 2019, 394, 29-38.	6.3	665
11	Geriatric assessment predicts survival and toxicities in elderly myeloma patients: an International Myeloma Working Group report. Blood, 2015, 125, 2068-2074.	0.6	586
12	Comparison of 200 mg/m2 melphalan and 8 Gy total body irradiation plus 140 mg/m2 melphalan as conditioning regimens for peripheral blood stem cell transplantation in patients with newly diagnosed multiple myeloma: final analysis of the Intergroupe Francophone du Myelome 9502 randomized trial. Blood, 2002, 99, 731-735.	0.6	531
13	Efficacy of venetoclax as targeted therapy for relapsed/refractory t(11;14) multiple myeloma. Blood, 2017, 130, 2401-2409.	0.6	403
14	Minimal residual disease negativity using deep sequencing is a major prognostic factor in multiple myeloma. Blood, 2018, 132, 2456-2464.	0.6	301
15	Prospective Evaluation of Magnetic Resonance Imaging and [¹⁸ F]Fluorodeoxyglucose Positron Emission Tomography-Computed Tomography at Diagnosis and Before Maintenance Therapy in Symptomatic Patients With Multiple Myeloma Included in the IFM/DFCI 2009 Trial: Results of the IMAJEM Study. Journal of Clinical Oncology. 2017. 35, 2911-2918.	0.8	247
16	Frontline therapy of multiple myeloma. Blood, 2015, 125, 3076-3084.	0.6	244
17	VTD is superior to VCD prior to intensive therapy in multiple myeloma: results of the prospective IFM2013-04 trial. Blood, 2016, 127, 2569-2574.	0.6	224
18	Final analysis of survival outcomes in the phase 3 FIRST trial of up-front treatment for multiple myeloma. Blood, 2018, 131, 301-310.	0.6	216

#	Article	IF	CITATIONS
19	Prognostic role of circulating exosomal miRNAs in multiple myeloma. Blood, 2017, 129, 2429-2436.	0.6	214
20	Once-per-week selinexor, bortezomib, and dexamethasone versus twice-per-week bortezomib and dexamethasone in patients with multiple myeloma (BOSTON): a randomised, open-label, phase 3 trial. Lancet, The, 2020, 396, 1563-1573.	6.3	188
21	Isatuximab, carfilzomib, and dexamethasone in relapsed multiple myeloma (IKEMA): a multicentre, open-label, randomised phase 3 trial. Lancet, The, 2021, 397, 2361-2371.	6.3	177
22	Combination of International Scoring System 3, High Lactate Dehydrogenase, and t(4;14) and/or del(17p) Identifies Patients With Multiple Myeloma (MM) Treated With Front-Line Autologous Stem-Cell Transplantation at High Risk of Early MM Progression–Related Death. Journal of Clinical Oncology, 2014, 32, 2173-2180.	0.8	150
23	Daratumumab, lenalidomide, and dexamethasone versus lenalidomide and dexamethasone alone in newly diagnosed multiple myeloma (MAIA): overall survival results from a randomised, open-label, phase 3 trial. Lancet Oncology, The, 2021, 22, 1582-1596.	5.1	141
24	Treatment of relapsed and refractory multiple myeloma: recommendations from the International Myeloma Working Group. Lancet Oncology, The, 2021, 22, e105-e118.	5.1	136
25	A simplified frailty scale predicts outcomes in transplant-ineligible patients with newly diagnosed multiple myeloma treated in the FIRST (MM-020) trial. Leukemia, 2020, 34, 224-233.	3.3	122
26	Labial salivary gland biopsy is a reliable test for the diagnosis of primary and secondary amyloidosis. a prospective clinical and immunohistologic study in 59 patients. Arthritis and Rheumatism, 1993, 36, 691-697.	6.7	96
27	Bortezomib, Doxorubicin, Cyclophosphamide, Dexamethasone Induction Followed by Stem Cell Transplantation for Primary Plasma Cell Leukemia: A Prospective Phase II Study of the Intergroupe Francophone du Myélome. Journal of Clinical Oncology, 2016, 34, 2125-2132.	0.8	91
28	High incidence of cryptic translocations involving the Ig heavy chain gene in multiple myeloma, as shown by fluorescence in situ hybridization. , 1999, 24, 9-15.		84
29	Role of additional chromosomal changes in the prognostic value of t(4;14) and del(17p) in multiple myeloma: the IFM experience. Blood, 2015, 125, 2095-2100.	0.6	82
30	Recommendations for vaccination in multiple myeloma: a consensus of the European Myeloma Network. Leukemia, 2021, 35, 31-44.	3.3	79
31	Health-related quality-of-life in patients with newly diagnosed multiple myeloma in the FIRST trial: lenalidomide plus low-dose dexamethasone versus melphalan, prednisone, thalidomide. Haematologica, 2015, 100, 826-833.	1.7	76
32	Updated Outcomes and Impact of Age With Lenalidomide and Low-Dose Dexamethasone or Melphalan, Prednisone, and Thalidomide in the Randomized, Phase III FIRST Trial. Journal of Clinical Oncology, 2016, 34, 3609-3617.	0.8	71
33	Prevention and management of adverse events of novel agents in multiple myeloma: a consensus of the European Myeloma Network. Leukemia, 2018, 32, 1542-1560.	3.3	68
34	Evaluation of Daratumumab for the Treatment of Multiple Myeloma in Patients With High-risk Cytogenetic Factors. JAMA Oncology, 2020, 6, 1759.	3.4	64
35	P53 deletion is not a frequent event in multiple myeloma. British Journal of Haematology, 1999, 106, 717-719.	1.2	62
36	The molecular make up of smoldering myeloma highlights the evolutionary pathways leading to multiple myeloma. Nature Communications, 2021, 12, 293.	5.8	54

#	Article	IF	CITATIONS
37	A predictive model for risk of early grade ≥ 3 infection in patients with multiple myeloma not eligible for transplant: analysis of the FIRST trial. Leukemia, 2018, 32, 1404-1413.	3.3	53
38	del(17p) without <i>TP53</i> mutation confers a poor prognosis in intensively treated newly diagnosed patients with multiple myeloma. Blood, 2021, 137, 1192-1195.	0.6	48
39	Phase 1/2 study of carfilzomib plus melphalan and prednisone in patients aged over 65 years with newly diagnosed multiple myeloma. Blood, 2015, 125, 3100-3104.	0.6	47
40	Deregulation and Targeting of TP53 Pathway in Multiple Myeloma. Frontiers in Oncology, 2018, 8, 665.	1.3	47
41	Defining the vulnerable patient with myeloma—a frailty position paper of the European Myeloma Network. Leukemia, 2020, 34, 2285-2294.	3.3	45
42	Multiple Myeloma: EHA-ESMO Clinical Practice Guidelines for Diagnosis, Treatment and Follow-up. HemaSphere, 2021, 5, e528.	1.2	45
43	Early relapse after autologous transplant for myeloma is associated with poor survival regardless of cytogenetic risk. Haematologica, 2020, 105, e480-483.	1.7	42
44	Daratumumab plus lenalidomide and dexamethasone in transplant-ineligible newly diagnosed multiple myeloma: frailty subgroup analysis of MAIA. Leukemia, 2022, 36, 1066-1077.	3.3	39
45	Current state and next-generation CAR-T cells in multiple myeloma. Blood Reviews, 2022, 54, 100929.	2.8	38
46	<i>BRAF</i> and <i>DIS3</i> Mutations Associate with Adverse Outcome in a Long-term Follow-up of Patients with Multiple Myeloma. Clinical Cancer Research, 2020, 26, 2422-2432.	3.2	37
47	Bendamustine is effective in T ell prolymphocytic leukaemia. British Journal of Haematology, 2015, 168, 916-919.	1.2	36
48	Frontline Therapy with Carfilzomib, Lenalidomide, and Dexamethasone (KRd) Induction Followed By Autologous Stem Cell Transplantation, Krd Consolidation and Lenalidomide Maintenance in Newly Diagnosed Multiple Myeloma (NDMM) Patients: Primary Results of the Intergroupe Francophone Du MyA©Lome (IFM) Krd Phase II Study. Blood, 2016, 128, 1142-1142.	0.6	36
49	Age is a prognostic factor even among patients with multiple myeloma younger than 66 years treated with high-dose melphalan: the IFM experience on 2316 patients. Haematologica, 2014, 99, 1236-1238.	1.7	35
50	Effective anti-BCMA retreatment in multiple myeloma. Blood Advances, 2021, 5, 3016-3020.	2.5	30
51	Updated Analysis of Daratumumab Plus Lenalidomide and Dexamethasone (D-Rd) Versus Lenalidomide and Dexamethasone (Rd) in Patients with Transplant-Ineligible Newly Diagnosed Multiple Myeloma (NDMM): The Phase 3 Maia Study. Blood, 2020, 136, 24-26.	0.6	29
52	Isatuximab plus pomalidomide and dexamethasone in elderly patients with relapsed/refractory multiple myeloma: ICARIA-MM subgroup analysis. Haematologica, 2021, 106, 1182-1187.	1.7	27
53	Subgroup analysis of ICARIAâ€MM study in relapsed/refractory multiple myeloma patients with highâ€risk cytogenetics. British Journal of Haematology, 2021, 194, 120-131.	1.2	27
54	Daratumumab is effective in the relapsed or refractory systemic lightâ€chain amyloidosis but associated with high infection burden in a frail realâ€life population. British Journal of Haematology, 2020, 188, e24-e27.	1.2	26

#	Article	IF	CITATIONS
55	Minimal Residual Disease in Myeloma: Application for Clinical Care and New Drug Registration. Clinical Cancer Research, 2021, 27, 5195-5212.	3.2	26
56	Daratumumab Plus Lenalidomide and Dexamethasone (D-Rd) Versus Lenalidomide and Dexamethasone (Rd) in Patients with Newly Diagnosed Multiple Myeloma (NDMM) Ineligible for Transplant: Updated Analysis of Maia. Blood, 2019, 134, 1875-1875.	0.6	26
57	Daratumumabâ€lenalidomideâ€dexamethasone vs standardâ€ofâ€care regimens: Efficacy in transplantâ€ineligibl untreated myeloma. American Journal of Hematology, 2020, 95, 1486-1494.	^e 2.0	25
58	The Prognostic Impact of Complete Remission (CR) Plus Very Good Partial Remission (VGPR) in a Double-Transplantation Program for Newly Diagnosed Multiple Myeloma (MM). Combined Results of the IFM 99 Trials Blood, 2006, 108, 3077-3077.	0.6	20
59	Phase I Study of AVE1642 Anti IGF-1R Monoclonal Antibody in Patients with Advanced Multiple Myeloma Blood, 2007, 110, 1166-1166.	0.6	19
60	Daratumumab and dexamethasone is safe and effective for triple refractory myeloma patients: final results of the IFM 2014â€04 (Etoile du Nord) trial. British Journal of Haematology, 2019, 187, 319-327.	1.2	18
61	Efficacy and safety of carfilzomib-based regimens in frail patients with relapsed and/or refractory multiple myeloma. Blood Advances, 2020, 4, 5449-5459.	2.5	17
62	Bortezomib, lenalidomide, and dexamethasone (VRd) ± daratumumab (DARA) in patients (pts) with newly diagnosed multiple myeloma (NDMM) for whom transplant is not planned as initial therapy: A multicenter, randomized, phase III study (CEPHEUS) Journal of Clinical Oncology, 2019, 37, TPS8056-TPS8056.	0.8	17
63	Onceâ€weekly (70 mg/m ²) vs twiceâ€weekly (56 mg/m ²) dosing of carfilzomib in patients with relapsed or refractory multiple myeloma: A post hoc analysis of the ENDEAVOR, A.R.R.O.W., and CHAMPIONâ€1 trials. Cancer Medicine, 2020, 9, 2989-2996.	1.3	16
64	Effect of age and frailty on the efficacy and tolerability of onceâ€weekly selinexor, bortezomib, and dexamethasone in previously treated multiple myeloma. American Journal of Hematology, 2021, 96, 708-718.	2.0	16
65	Continuous treatment with lenalidomide and lowâ€dose dexamethasone in transplantâ€ineligible patients with newly diagnosed multiple myeloma in Asia: subanalysis of the <scp>FIRST</scp> trial. British Journal of Haematology, 2017, 176, 743-749.	1.2	14
66	A Frailty Scale Predicts Outcomes of Patients with Newly Diagnosed Multiple Myeloma Who Are Ineligible for Transplant Treated with Continuous Lenalidomide Plus Low-Dose Dexamethasone on the First Trial. Blood, 2015, 126, 4239-4239.	0.6	13
67	The genomic landscape of plasma cells in systemic light chain amyloidosis. Blood, 2018, 132, 2775-2777.	0.6	12
68	Improved survival in multiple myeloma during the 2005–2009 and 2010–2014 periods. Leukemia, 2021, 35, 3600-3603.	3.3	11
69	Synergistic effects of low-dose belantamab mafodotin in combination with a gamma-secretase inhibitor (nirogacestat) in patients with relapsed/refractory multiple myeloma (RRMM): DREAMM-5 study Journal of Clinical Oncology, 2022, 40, 8019-8019.	0.8	11
70	Isatuximab plus pomalidomide and dexamethasone in frail patients with relapsed/refractory multiple myeloma: <scp>ICARIAâ€MM</scp> subgroup analysis. American Journal of Hematology, 2021, 96, E423-E427.	2.0	10
71	Efficacy of Daratumumab in Combination with Standard of Care Regimens in Lenalidomide-Exposed or -Refractory Patients with Relapsed/Refractory Multiple Myeloma (RRMM): Analysis of the Castor, Pollux, and MMY1001 Studies. Blood, 2018, 132, 3288-3288.	0.6	10
72	Treatment Regimens for Transplant-Ineligible Patients With Newly Diagnosed Multiple Myeloma: A Systematic Literature Review and Network Meta-analysis. Advances in Therapy, 2022, 39, 1976-1992.	1.3	10

#	Article	IF	CITATIONS
73	Quality of life analyses in patients with multiple myeloma: results from the Selinexor (KPT-330) Treatment of Refractory Myeloma (STORM) phase 2b study. BMC Cancer, 2021, 21, 993.	1.1	8
74	A Comprehensive Analysis of Cytogenetic Abnormalities in Myeloma: Results of the FISH Analysis of 1000 Patients Enrolled in the IFM99 Trials Blood, 2005, 106, 622-622.	0.6	8
75	Treatment approach for the older, unfit patient with myeloma from diagnosis to relapse: perspectives of a European hematologist. Hematology American Society of Hematology Education Program, 2018, 2018, 83-87.	0.9	7
76	Frontline Therapy for Patients with Multiple Myeloma not Eligible for Stem Cell Transplantation. Hematology/Oncology Clinics of North America, 2014, 28, 829-838.	0.9	6
77	Clinical Outcomes in Patients (Pts) with Dose Reduction of Selinexor in Combination with Bortezomib, and Dexamethasone (XVd) in Previously Treated Multiple Myeloma from the Boston Study. Blood, 2021, 138, 3793-3793.	0.6	6
78	Isatuximab plus carfilzomib and dexamethasone versus carfilzomib and dexamethasone in elderly patients with relapsed multiple myeloma: IKEMA subgroup analysis. Hematological Oncology, 2022, 40, 1020-1029.	0.8	6
79	Isatuximab plus carfilzomib and dexamethasone versus carfilzomib and dexamethasone in elderly patients with relapsed multiple myeloma: IKEMA subgroup analysis Journal of Clinical Oncology, 2021, 39, 8026-8026.	0.8	5
80	Isatuximab plus carfilzomib and dexamethasone in relapsed multiple myeloma patients with high-risk cytogenetics: IKEMA subgroup analysis Journal of Clinical Oncology, 2021, 39, 8042-8042.	0.8	5
81	Predictive biomarkers with isatuximab plus pomalidomide and dexamethasone in relapsed/refractory multiple myeloma. Blood Cancer Journal, 2021, 11, 55.	2.8	4
82	Association of Morbid Progression With Overall Survival Among Patients With Multiple Myeloma: Validation of the Progression-free Survival Endpoint. Clinical Lymphoma, Myeloma and Leukemia, 2021, 21, 345-354.e4.	0.2	4
83	First-Line Use of Daratumumab, Lenalidomide, and Dexamethasone Confers Survival Benefit Compared with Second-Line Use of Daratumumab-Based Regimens in Transplant-Ineligible Patients with Multiple Myeloma: Analysis of Different Clinical Scenarios. Blood, 2021, 138, 118-118.	0.6	4
84	Evaluation of the Relevance of Surgery in Patients with Multiple Myeloma Harboring Symptomatic Spinal Involvement: A Retrospective Case Series. World Neurosurgery, 2018, 114, e356-e365.	0.7	3
85	A matching-adjusted indirect treatment comparison (MAIC) of daratumumab–bortezomib–melphalan–prednisone (D-VMP) versus lenalidomide–dexamethasone continuous (Rd continuous), lenalidomide–dexamethasone 18 months (Rd 18), and melphalan–rrednisone–thalidomide (MPI) Leukemia and Lymphoma, 2020, 61, 714-720	0.6	3
86	Case Report: Two Cases of Cryptosporidiosis in Heavily Pretreated Patients With Myeloma. Clinical Lymphoma, Myeloma and Leukemia, 2021, 21, e545-e547.	0.2	3
87	Efficacy of Isatuximab with Pomalidomide and Dexamethasone in Elderly Patients with Relapsed/Refractory Multiple Myeloma: Icaria-MM Subgroup Analysis. Blood, 2019, 134, 1893-1893.	0.6	3
88	Isatuximab plus pomalidomide and dexamethasone in elderly patients with relapsed/refractory multiple myeloma: ICARIA-MM subgroup analysis. Haematologica, 2022, 107, 774-775.	1.7	2
89	A Matching-Adjusted Indirect Treatment Comparison of Daratumumab-Bortezomib-Melphalan-Prednisone Versus Lenalidomide-Dexamethasone Continuous, Lenalidomide-Dexamethasone 18 Months, and Melphalan-Prednisone-Thalidomide. Blood, 2018, 132, 3551-3551	0.6	1
90	Effectiveness of Daratumumab in Combination with Lenalidomide and Dexamethasone (DRd) Vs. Common Standard-of-Care Regimens in Patients with Non-Transplant Newly Diagnosed Multiple Myeloma (NDMM). Blood, 2019, 134, 1830-1830.	0.6	1

#	Article	IF	CITATIONS
91	The Relationship between Baseline Biomarkers and Efficacy of Isatuximab in Combination with Pomalidomide and Dexamethasone in RRMM: Insights from Phase 1 and Phase 3 Studies. Blood, 2019, 134, 3179-3179.	0.6	1
92	High-Resolution Genomic Profiles Identify Novel Genes and/or Chromosomal Regions with Prognostic and Oncogenic Significance in Myeloma Patients Blood, 2007, 110, 657-657.	0.6	0
93	Guérison du myélome multiple : un objectif envisageable à court terme ?. Bulletin De L'Academie Nationale De Medecine, 2018, 202, 953-961.	0.0	0
94	Mutations and Copy Number Changes Predict Progression from Smoldering Myeloma to Symptomatic Myeloma in the Era of Novel IMWG Criteria. Blood, 2018, 132, 4456-4456.	0.6	0