

# Jinghui Zhai

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1296549/publications.pdf>

Version: 2024-02-01

7  
papers

187  
citations

1478280  
6  
h-index

1719901  
7  
g-index

7  
all docs

7  
docs citations

7  
times ranked

233  
citing authors

#	ARTICLE	IF	CITATIONS
1	Calycosin ameliorates doxorubicin-induced cardiotoxicity by suppressing oxidative stress and inflammation via the sirtuin 1 "NOD-like receptor protein 3 pathway. <i>Phytotherapy Research</i> , 2020, 34, 649-659.	2.8	59
2	Dysregulation of BSEP and MRP2 May Play an Important Role in Isoniazid-Induced Liver Injury & via the SIRT1/FXR Pathway in Rats and HepG2 Cells. <i>Biological and Pharmaceutical Bulletin</i> , 2018, 41, 1211-1218.	0.6	47
3	Quercetin protected against isoniazide-induced HepG2 cell apoptosis by activating the SIRT1/ERK pathway. <i>Journal of Biochemical and Molecular Toxicology</i> , 2019, 33, e22369.	1.4	25
4	Quercetin attenuates NLRP3 inflammasome activation and apoptosis to protect INH-induced liver injury via regulating SIRT1 pathway. <i>International Immunopharmacology</i> , 2020, 85, 106634.	1.7	25
5	Ginsenoside Rg3 attenuates cisplatin-induced kidney injury through inhibition of apoptosis and autophagy-inhibited NLRP3. <i>Journal of Biochemical and Molecular Toxicology</i> , 2021, 35, e22896.	1.4	13
6	Salvianolic Acid B Attenuates Apoptosis of HUVEC Cells Treated with High Glucose or High Fat via Sirt1 Activation. <i>Evidence-based Complementary and Alternative Medicine</i> , 2019, 2019, 1-11.	0.5	12
7	Salvianolic acid inhibits the effects of high glucose on vascular endothelial dysfunction by modulating the Sirt1-NOS pathway. <i>Journal of Biochemical and Molecular Toxicology</i> , 2019, 33, e22245.	1.4	6