

# Ann-Charlotte Iversen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1305659/publications.pdf>

Version: 2024-02-01

55  
papers

1,826  
citations

257450

24  
h-index

276875

41  
g-index

64  
all docs

64  
docs citations

64  
times ranked

2860  
citing authors

#	ARTICLE	IF	CITATIONS
1	Changes in Serum Cytokines Throughout Pregnancy in Women With Polycystic Ovary Syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, 107, 39-52.	3.6	11
2	Divergent Regulation of Decidual Oxidative-Stress Response by NRF2 and KEAP1 in Preeclampsia with and without Fetal Growth Restriction. <i>International Journal of Molecular Sciences</i> , 2022, 23, 1966.	4.1	11
3	Circulating Levels of Anti-C1q and Anti-Factor H Autoantibodies and Their Targets in Normal Pregnancy and Preeclampsia. <i>Frontiers in Immunology</i> , 2022, 13, 842451.	4.8	5
4	TLR3 expression by maternal and fetal cells at the maternal-fetal interface in normal and preeclamptic pregnancies. <i>Journal of Leukocyte Biology</i> , 2021, 109, 173-183.	3.3	14
5	Decidual and placental NOD1 is associated with inflammation in normal and preeclamptic pregnancies. <i>Placenta</i> , 2021, 105, 23-31.	1.5	10
6	Systemic immunological perturbations and placental pathology in preeclampsia. <i>Placenta</i> , 2021, 112, e18-e19.	1.5	0
7	Cytokine Patterns in Maternal Serum From First Trimester to Term and Beyond. <i>Frontiers in Immunology</i> , 2021, 12, 752660.	4.8	40
8	Cholesterol Crystals and NLRP3 Mediated Inflammation in the Uterine Wall Decidua in Normal and Preeclamptic Pregnancies. <i>Frontiers in Immunology</i> , 2020, 11, 564712.	4.8	15
9	Genetic predisposition to hypertension is associated with preeclampsia in European and Central Asian women. <i>Nature Communications</i> , 2020, 11, 5976.	12.8	102
10	NLRP3 inflammasome expression by maternal and fetal cells in the decidua and its association with preeclampsia. <i>Placenta</i> , 2019, 83, e15.	1.5	1
11	Metabolomics Identifies Placental Dysfunction and Confirms Flt-1 (FMS-Like Tyrosine Kinase Receptor 1) Biomarker Specificity. <i>Hypertension</i> , 2019, 74, 1136-1143.	2.7	14
12	Hypertensive pregnancy disorders increase the risk of maternal cardiovascular disease after adjustment for cardiovascular risk factors. <i>International Journal of Cardiology</i> , 2019, 282, 81-87.	1.7	39
13	Serum cytokine patterns in first half of pregnancy. <i>Cytokine</i> , 2019, 119, 188-196.	3.2	29
14	Metabolomics identifies placental dysfunction and confirms Flt-1 biomarker specificity. <i>Pregnancy Hypertension</i> , 2019, 17, S5.	1.4	0
15	Placental inflammation in pre-eclampsia by Nod-like receptor protein (NLRP)3 inflammasome activation in trophoblasts. <i>Clinical and Experimental Immunology</i> , 2018, 193, 84-94.	2.6	75
16	Placental inflammation by HMGB1 activation of TLR4 at the syncytium. <i>Placenta</i> , 2018, 72-73, 53-61.	1.5	24
17	Association Between Gestational Hypertension and Risk of Cardiovascular Disease Among 617,589 Norwegian Women. <i>Journal of the American Heart Association</i> , 2018, 7, .	3.7	85
18	Incident Coronary Heart Disease After Preeclampsia: Role of Reduced Fetal Growth, Preterm Delivery, and Parity. <i>Journal of the American Heart Association</i> , 2017, 6, .	3.7	77

#	ARTICLE	IF	CITATIONS
19	Variants in the fetal genome near FLT1 are associated with risk of preeclampsia. <i>Nature Genetics</i> , 2017, 49, 1255-1260.	21.4	205
20	The antihypertensive MTHFR gene polymorphism rs17367504-G is a possible novel protective locus for preeclampsia. <i>Journal of Hypertension</i> , 2017, 35, 132-139.	0.5	15
21	NLRP3 inflammasome expression and activation at the maternal-fetal interface in preeclamptic and healthy pregnancies. <i>Placenta</i> , 2016, 45, 88.	1.5	0
22	Refined phenotyping identifies links between preeclampsia and related diseases in a Norwegian preeclampsia family cohort. <i>Journal of Hypertension</i> , 2015, 33, 2294-2302.	0.5	21
23	Meta-analysis of the human leukocyte antigen (HLA) 14 bp insertion/deletion polymorphism as a risk factor for preeclampsia. <i>Tissue Antigens</i> , 2015, 86, 186-194.	1.0	24
24	First Trimester Urine and Serum Metabolomics for Prediction of Preeclampsia and Gestational Hypertension: A Prospective Screening Study. <i>International Journal of Molecular Sciences</i> , 2015, 16, 21520-21538.	4.1	55
25	[278-POS]. <i>Pregnancy Hypertension</i> , 2015, 5, 138-139.	1.4	0
26	Distinct First Trimester Cytokine Profiles for Gestational Hypertension and Preeclampsia. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2015, 35, 2478-2485.	2.4	36
27	Toll-like receptor profiling of seven trophoblast cell lines warrants caution for translation to primary trophoblasts. <i>Placenta</i> , 2015, 36, 1246-1253.	1.5	30
28	Metabolic profiles of placenta in preeclampsia using HR-MAS MRS metabolomics. <i>Placenta</i> , 2015, 36, 1455-1462.	1.5	53
29	Metabolomic Biomarkers in Serum and Urine in Women with Preeclampsia. <i>PLoS ONE</i> , 2014, 9, e91923.	2.5	54
30	Functional Toll-like receptors in primary first-trimester trophoblasts. <i>Journal of Reproductive Immunology</i> , 2014, 106, 89-99.	1.9	45
31	Preeclampsia and cardiovascular disease share genetic risk factors on chromosome 2q22. <i>Pregnancy Hypertension</i> , 2014, 4, 178-185.	1.4	14
32	InterPregGen: genetic studies of pre-eclampsia in three continents. <i>Norsk Epidemiologi</i> , 2014, 24, 141-146.	0.3	12
33	PP042. Cell surface toll-like receptors in primary first trimester trophoblasts. <i>Pregnancy Hypertension</i> , 2013, 3, 81-82.	1.4	1
34	PP002. Metabolomic biomarkers in serum and urine of preeclamptic women. <i>Pregnancy Hypertension</i> , 2013, 3, 67-68.	1.4	0
35	OP004. A SNP associated with susceptibility to preeclampsia near the inhibin, beta B gene, is also associated with cardiovascular disease risk traits. <i>Pregnancy Hypertension</i> , 2013, 3, 63.	1.4	0
36	PP040. Activation of endosomal toll-like receptors in first trimester trophoblasts. <i>Pregnancy Hypertension</i> , 2013, 3, 81.	1.4	0

#	ARTICLE	IF	CITATIONS
37	Inflammatory mechanisms in preeclampsia. <i>Pregnancy Hypertension</i> , 2013, 3, 58.	1.4	8
38	Changing patterns of cytomegalovirus seroprevalence among pregnant women in Norway between 1995 and 2009 examined in the Norwegian Mother and Child Cohort Study and two cohorts from Sør-Trøndelag County: a cross-sectional study. <i>BMJ Open</i> , 2013, 3, e003066.	1.9	18
39	Mediators of the association between pre-eclampsia and cerebral palsy: population based cohort study. <i>BMJ</i> , 2013, 347, f4089-f4089.	6.0	59
40	Cytomegalovirus antibody status at 17-18 weeks of gestation and pre-eclampsia: a case-control study of pregnant women in Norway. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2012, 119, 1316-1323.	2.3	13
41	OS046. Genome-wide association scans identify novel maternal susceptibility loci for preeclampsia. <i>Pregnancy Hypertension</i> , 2012, 2, 202.	1.4	2
42	OS070. Shared genetic risk factors for preeclampsia and cardiovascular disease. <i>Pregnancy Hypertension</i> , 2012, 2, 214-215.	1.4	0
43	Genome-Wide Association Scan Identifies a Risk Locus for Preeclampsia on 2q14, Near the Inhibin, Beta B Gene. <i>PLoS ONE</i> , 2012, 7, e33666.	2.5	110
44	IL-10 Enhances MD-2 and CD14 Expression in Monocytes and the Proteins Are Increased and Correlated in HIV-Infected Patients. <i>Journal of Immunology</i> , 2009, 182, 588-595.	0.8	27
45	A Proviral Role for CpG in Cytomegalovirus Infection. <i>Journal of Immunology</i> , 2009, 182, 5672-5681.	0.8	31
46	The HLA-G 14bp gene polymorphism and decidual HLA-G 14bp gene expression in pre-eclamptic and normal pregnancies. <i>Journal of Reproductive Immunology</i> , 2008, 78, 158-165.	1.9	52
47	Fetal growth restriction is associated with reduced FasL expression by decidual cells. <i>Journal of Reproductive Immunology</i> , 2007, 74, 7-14.	1.9	22
48	A comparative study of immunomagnetic methods used for separation of human natural killer cells from peripheral blood. <i>Journal of Immunological Methods</i> , 2005, 303, 1-10.	1.4	19
49	Human NK Cells Inhibit Cytomegalovirus Replication through a Noncytolytic Mechanism Involving Lymphotoxin-Dependent Induction of IFN- $\gamma$ . <i>Journal of Immunology</i> , 2005, 175, 7568-7574.	0.8	37
50	The role of interleukin-2 in regulating the sensitivity of natural killer cells for Fas-mediated apoptosis. <i>Cancer Immunology, Immunotherapy</i> , 1999, 48, 139-146.	4.2	22
51	REGULATION OF APO-2 LIGAND/TRAIL EXPRESSION IN NK CELLS—INVOLVEMENT IN NK CELL-MEDIATED CYTOTOXICITY. <i>Cytokine</i> , 1999, 11, 664-672.	3.2	83
52	Apoptosis, proliferation and NF- $\kappa$ B activation induced by agonistic Fas antibodies in the human myeloma cell line OH-2: amplification of Fas-mediated apoptosis by tumor necrosis factor. <i>European Journal of Haematology</i> , 1999, 63, 345-353.	2.2	21
53	REGULATION OF FAS AND FAS-LIGAND EXPRESSION IN NK CELLS BY CYTOKINES AND THE INVOLVEMENT OF FAS-LIGAND IN NK/LAK CELL-MEDIATED CYTOTOXICITY. <i>Cytokine</i> , 1997, 9, 394-404.	3.2	75
54	Polymorphonuclear granulocytes enhance lipopolysaccharide-induced soluble p75 tumor necrosis factor receptor release from mononuclear cells. <i>European Journal of Immunology</i> , 1995, 25, 2714-2717.	2.9	32

#	ARTICLE	IF	CITATIONS
55	Gene expression and secretion of cytokines and cytokine receptors from highly purified CD56+ natural killer cells stimulated with interleukin-2, interleukin-7 and interleukin-12. European Journal of Immunology, 1993, 23, 1831-1838.	2.9	77