

Valeria Severino

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/131662/valeria-severino-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

44
papers

734
citations

15
h-index

24
g-index

44
ext. papers

858
ext. citations

4
avg, IF

3.35
L-index

#	Paper	IF	Citations
44	Extracellular Vesicles in Bile as Markers of Malignant Biliary Stenoses. <i>Gastroenterology</i> , 2017 , 153, 495-504	13.8	53
43	Trifluoroacetylated tyrosine-rich D-tetrapeptides have potent antioxidant activity. <i>Peptides</i> , 2017 , 89, 50-59	3.8	5
42	AMP18 interacts with the anion exchanger SLC26A3 and enhances its expression in gastric cancer cells. <i>Biochimie</i> , 2016 , 121, 151-60	4.6	10
41	De-regulated expression of the BRG1 chromatin remodeling factor in bone marrow mesenchymal stromal cells induces senescence associated with the silencing of NANOG and changes in the levels of chromatin proteins. <i>Cell Cycle</i> , 2015 , 14, 1315-26	4.7	20
40	Structural characterization of dioicin 1 from <i>Phytolacca dioica</i> L. gains novel insights into phylogenetic relationships of Phytolaccaceae type 1 RIPs. <i>Biochemical and Biophysical Research Communications</i> , 2015 , 463, 732-8	3.4	4
39	Structural Elucidation of Saponins: A Combined Approach Based on High-Resolution Spectroscopic Techniques. <i>Studies in Natural Products Chemistry</i> , 2015 , 45, 85-120	1.5	2
38	An improved UPLC method for the detection of undeclared horse meat addition by using myoglobin as molecular marker. <i>Food Chemistry</i> , 2015 , 169, 241-5	8.5	30
37	A targeted secretome profiling by multiplexed immunoassay revealed that secreted chemokine ligand 2 (MCP-1/CCL2) affects neural differentiation in mesencephalic neural progenitor cells. <i>Proteomics</i> , 2015 , 15, 714-24	4.8	9
36	Insight into the structural and functional features of myoglobin from <i>Hystrix cristata</i> L. and <i>Rangifer tarandus</i> L.. <i>RSC Advances</i> , 2015 , 5, 26388-26401	3.7	4
35	Ribosome-inactivating Proteins from Phytolaccaceae 2014 , 28-43		1
34	Purification and characterization of novel cationic peroxidases from <i>Asparagus acutifolius</i> L. with biotechnological applications. <i>Molecular Biotechnology</i> , 2014 , 56, 738-46	3	8
33	Two new acylated drimane-type sesquiterpene glucosides from <i>Petrohragia saxifraga</i> . <i>Phytochemistry Letters</i> , 2014 , 7, 46-51	1.9	1
32	Ultra-high performance liquid chromatography tandem mass spectrometry for the detection of durum wheat contamination or adulteration. <i>Journal of Mass Spectrometry</i> , 2014 , 49, 1239-46	2.2	15
31	Saliva from obese individuals suppresses the release of aroma compounds from wine. <i>PLoS ONE</i> , 2014 , 9, e85611	3.7	70
30	Comparison of binder compositions in Pompeian wall painting styles from Insula Occidentalis. <i>Chemistry Central Journal</i> , 2014 , 8, 65		13
29	Anti-amyloidogenic property of human gastrokine 1. <i>Biochimie</i> , 2014 , 106, 91-100	4.6	14
28	Molecular profiling of the <i>Phytophthora plurivora</i> secretome: a step towards understanding the cross-talk between plant pathogenic oomycetes and their hosts. <i>PLoS ONE</i> , 2014 , 9, e112317	3.7	12

27	A comparative structural and bioanalytical study of IVIG clinical lots. <i>Molecular Biotechnology</i> , 2013 , 54, 983-95	3	9
26	An IgE receptor mimetic peptide (PepE) protects mice from IgE mediated anaphylaxis. <i>Molecular BioSystems</i> , 2013 , 9, 2853-9		12
25	Raviscanina wild asparagus (<i>Asparagus acutifolius</i> L.): A nutritionally valuable crop with antioxidant and antiproliferative properties. <i>Food Research International</i> , 2013 , 53, 180-188	7	39
24	Spectroscopic identification and anti-biofilm properties of polar metabolites from the medicinal plant <i>Helichrysum italicum</i> against <i>Pseudomonas aeruginosa</i> . <i>Bioorganic and Medicinal Chemistry</i> , 2013 , 21, 7038-46	3.4	33
23	Secretome profiling of differentiated neural mes-c-myc A1 cell line endowed with stem cell properties. <i>Biochimica Et Biophysica Acta - Proteins and Proteomics</i> , 2013 , 1834, 2385-95	4	10
22	Analysis of secreted proteins. <i>Methods in Molecular Biology</i> , 2013 , 1002, 37-60	1.4	14
21	An integrated approach based on multiplexed protein array and iTRAQ labeling for in-depth identification of pathways associated to IVF outcome. <i>PLoS ONE</i> , 2013 , 8, e77303	3.7	15
20	Synthesis and biological properties of caffeic acid-PNA dimers containing guanine. <i>Molecules</i> , 2013 , 18, 9147-62	4.8	8
19	Polar and non-polar organic binder characterization in Pompeian wall paintings: comparison to a simulated painting mimicking an "a secco" technique. <i>Analytical and Bioanalytical Chemistry</i> , 2012 , 402, 3011-6	4.4	14
18	Detection of buffalo mozzarella adulteration by an ultra-high performance liquid chromatography tandem mass spectrometry methodology. <i>Journal of Mass Spectrometry</i> , 2012 , 47, 1407-14	2.2	35
17	A novel malate dehydrogenase from <i>Ceratonia siliqua</i> L. seeds with potential biotechnological applications. <i>Protein Journal</i> , 2012 , 31, 667-73	3.9	1
16	De novo sequencing and characterization of a novel Bowman-Birk inhibitor from <i>Lathyrus sativus</i> L. seeds by electrospray mass spectrometry. <i>Molecular BioSystems</i> , 2012 , 8, 3232-41		4
15	Structural and enzymatic properties of an in vivo proteolytic form of PD-S2, type 1 ribosome-inactivating protein from seeds of <i>Phytolacca dioica</i> L. <i>Biochemical and Biophysical Research Communications</i> , 2012 , 421, 514-20	3.4	8
14	Characterization of pigments and ligands in a wall painting fragment from Liternum archaeological park (Italy). <i>Journal of Separation Science</i> , 2012 , 35, 2986-93	3.4	9
13	Oleanane saponins from <i>Bellis sylvestris</i> Cyr. and evaluation of their phytotoxicity on <i>Aegilops geniculata</i> Roth. <i>Phytochemistry</i> , 2012 , 84, 125-34	4	27
12	Rapid primary structure determination of myoglobins by a complementary approach based on mass spectrometry and Edman degradation. <i>Food Chemistry</i> , 2012 , 133, 1646-1652	8.5	16
11	WCI, a novel wheat chymotrypsin inhibitor: purification, primary structure, inhibitory properties and heterologous expression. <i>Planta</i> , 2011 , 234, 723-35	4.7	10
10	Temporal proteomic profiling of embryonic stem cell secretome during cardiac and neural differentiation. <i>Proteomics</i> , 2011 , 11, 3972-82	4.8	27

9	Proteomic characterization of early changes induced by triiodothyronine in rat liver. <i>Journal of Proteome Research</i> , 2011 , 10, 3212-24	5.6	17
8	Proteomic analysis of human U937 cell line activation mediated by Haemophilus influenzae type b P2 porin and its surface-exposed loop 7. <i>Journal of Proteome Research</i> , 2010 , 9, 1050-62	5.6	21
7	The role of the glycan moiety on the structure-function relationships of PD-L1, type 1 ribosome-inactivating protein from <i>P. dioica</i> leaves. <i>Molecular BioSystems</i> , 2010 , 6, 570-9		11
6	Pathways affected by 3,5-diiodo-L-thyronine in liver of high fat-fed rats: evidence from two-dimensional electrophoresis, blue-native PAGE, and mass spectrometry. <i>Molecular BioSystems</i> , 2010 , 6, 2256-71		33
5	Quantification of thyrotropin-releasing hormone by liquid chromatography-electrospray mass spectrometry. <i>Amino Acids</i> , 2010 , 38, 1031-41	3.5	3
4	Crystal structure of PD-L1, a ribosome inactivating protein from <i>Phytolacca dioica</i> L. leaves with the property to induce DNA cleavage. <i>Biopolymers</i> , 2009 , 91, 1135-42	2.2	15
3	Improved procedure for protein binder analysis in mural painting by LC-ESI/Q-q-TOF mass spectrometry: detection of different milk species by casein proteotypic peptides. <i>Analytical and Bioanalytical Chemistry</i> , 2009 , 395, 2281-91	4.4	46
2	Expression of the brain transcription factor OTX1 occurs in a subset of normal germinal-center B cells and in aggressive Non-Hodgkin Lymphoma. <i>American Journal of Pathology</i> , 2009 , 175, 2609-17	5.8	20
1	Cloning and expression of the B chain of volkensin, type 2 ribosome inactivating protein from <i>Adenia volkensii</i> harms: co-folding with the A chain for heterodimer reconstitution. <i>Protein Expression and Purification</i> , 2007 , 51, 209-15	2	6