Hyungjin Rhee

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/1321434/hyungjin-rhee-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

19 1,177 49 33 h-index g-index citations papers 1,662 6.6 55 4.4 L-index avg, IF ext. citations ext. papers

#	Paper	IF	Citations
49	Molecular and radio-pathologic spectrum between hepatocellular carcinoma and intrahepatic cholangiocarcinoma <i>Hepatology</i> , 2022 ,	11.2	2
48	Ezetimibe combination therapy with statin for non-alcoholic fatty liver disease: an open-label randomized controlled trial (ESSENTIAL study) <i>BMC Medicine</i> , 2022 , 20, 93	11.4	4
47	MRI features of histologic subtypes of hepatocellular carcinoma: correlation with histologic, genetic, and molecular biologic classification <i>European Radiology</i> , 2022 , 1	8	1
46	Lymphoepithelioma-like neoplasm of the biliary tract with \probable low malignant potential\textit{V} Histopathology, 2021,	7.3	1
45	Direct-Acting Antivirals Improve Treatment Outcomes in Patients with Hepatitis C Virus-Related Hepatocellular Carcinoma Treated with Transarterial Chemoembolization: A Nationwide, Multi-center, Retrospective Cohort Study. <i>Digestive Diseases and Sciences</i> , 2021 , 66, 2427-2438	4	2
44	Update on Pathologic and Radiologic Diagnosis of Combined Hepatocellular-Cholangiocarcinoma. Journal of Liver Cancer, 2021 , 21, 12-24	0.7	0
43	Preoperative cephalhematoma size measured with computed tomography predicts intraoperative bleeding in pediatric patients undergoing cranioplasty. <i>Anesthesia and Pain Medicine</i> , 2021 , 16, 151-157	0.3	
42	Preoperative prediction of postsurgical outcomes in mass-forming intrahepatic cholangiocarcinoma based on clinical, radiologic, and radiomics features. <i>European Radiology</i> , 2021 , 31, 8638-8648	8	6
41	Radiomics analysis of contrast-enhanced CT for classification of hepatic focal lesions in colorectal cancer patients: its limitations compared to radiologists. <i>European Radiology</i> , 2021 , 31, 8786-8796	8	1
40	YAP inactivation in estrogen receptor alpha-positive hepatocellular carcinoma with less aggressive behavior. <i>Experimental and Molecular Medicine</i> , 2021 , 53, 1055-1067	12.8	1
39	Gadoxetic acid-enhanced MRI of macrotrabecular-massive hepatocellular carcinoma and its prognostic implications. <i>Journal of Hepatology</i> , 2021 , 74, 109-121	13.4	15
38	Clinical Staging of Mass-Forming Intrahepatic Cholangiocarcinoma: Computed Tomography Versus Magnetic Resonance Imaging. <i>Hepatology Communications</i> , 2021 , 5, 2009-2018	6	3
37	Lung and lymph node metastases from hepatocellular carcinoma: Comparison of pathological aspects. <i>Liver International</i> , 2021 ,	7.9	4
36	The Role of Imaging in Current Treatment Strategies for Pancreatic Adenocarcinoma. <i>Korean Journal of Radiology</i> , 2021 , 22, 23-40	6.9	12
35	Clinico-Radio-Pathological and Molecular Features of Hepatocellular Carcinomas with Keratin 19 Expression. <i>Liver Cancer</i> , 2020 , 9, 663-681	9.1	6
34	Diagnostic performance of Liver Imaging Reporting and Data System in patients at risk of both hepatocellular carcinoma and metastasis. <i>Abdominal Radiology</i> , 2020 , 45, 3789-3799	3	3
33	Gross type of hepatocellular carcinoma reflects the tumor hypoxia, fibrosis, and stemness-related marker expression. <i>Hepatology International</i> , 2020 , 14, 239-248	8.8	11

(2017-2020)

32	The Modified Response Evaluation Criteria in Solid Tumors (RECIST) Yield a More Accurate Prognoses Than the RECIST 1.1 in Hepatocellular Carcinoma Treated with Transarterial Radioembolization. <i>Gut and Liver</i> , 2020 , 14, 765-774	4.8	8	
31	Clinicopathological characteristics of intrahepatic cholangiocarcinoma according to gross morphologic type: cholangiolocellular differentiation traits and inflammation- and proliferation-phenotypes. <i>Hpb</i> , 2020 , 22, 864-873	3.8	8	
30	Vessels Encapsulating Tumor Clusters (VETC) Is a Powerful Predictor of Aggressive Hepatocellular Carcinoma. <i>Hepatology</i> , 2020 , 71, 183-195	11.2	44	
29	Dynamics of Genomic, Epigenomic, and Transcriptomic Aberrations during Stepwise Hepatocarcinogenesis. <i>Cancer Research</i> , 2019 , 79, 5500-5512	10.1	21	
28	Hepatic sarcomatoid carcinoma: magnetic resonance imaging evaluation by using the liver imaging reporting and data system. <i>European Radiology</i> , 2019 , 29, 3761-3771	8	14	
27	A dichotomous imaging classification for cholangiocarcinomas based on new histologic concepts. <i>European Journal of Radiology</i> , 2019 , 113, 182-187	4.7	3	
26	Hepatocellular Carcinoma with Irregular Rim-Like Arterial Phase Hyperenhancement: More Aggressive Pathologic Features. <i>Liver Cancer</i> , 2019 , 8, 24-40	9.1	27	
25	Alpha-Fetoprotein, Des-Gamma-Carboxy Prothrombin, and Modified RECIST Response as Predictors of Survival after Transarterial Radioembolization for Hepatocellular Carcinoma. <i>Journal of Vascular and Interventional Radiology</i> , 2019 , 30, 1194-1200.e1	2.4	10	
24	Increased Expression of the Matrix-Modifying Enzyme Lysyl Oxidase-Like 2 in Aggressive Hepatocellular Carcinoma with Poor Prognosis. <i>Gut and Liver</i> , 2019 , 13, 83-92	4.8	13	
23	A Comprehensive Review of Hepatocellular Carcinoma Enhancement Patterns in MRI: Emphasis on Gadoxetate-Enhanced Imaging. <i>Journal of the Korean Society of Radiology</i> , 2019 , 80, 374	0.2	1	
22	A proposal of imaging classification of intrahepatic mass-forming cholangiocarcinoma into ductal and parenchymal types: clinicopathologic significance. <i>European Radiology</i> , 2019 , 29, 3111-3121	8	8	
21	Keratin 19 Expression in Hepatocellular Carcinoma Is Regulated by Fibroblast-Derived HGF via a MET-ERK1/2-AP1 and SP1 Axis. <i>Cancer Research</i> , 2018 , 78, 1619-1631	10.1	38	
20	Transcriptomic and histopathological analysis of cholangiolocellular differentiation trait in intrahepatic cholangiocarcinoma. <i>Liver International</i> , 2018 , 38, 113-124	7.9	26	
19	Tumour epithelial and stromal characteristics of hepatocellular carcinomas with abundant fibrous stroma: fibrolamellar versus scirrhous hepatocellular carcinoma. <i>Histopathology</i> , 2017 , 71, 217-226	7.3	20	
18	Whole-Genome and Epigenomic Landscapes of Etiologically Distinct Subtypes of Cholangiocarcinoma. <i>Cancer Discovery</i> , 2017 , 7, 1116-1135	24.4	368	
17	Added value of smooth hypointense rim in the hepatobiliary phase of gadoxetic acid-enhanced MRI in identifying tumour capsule and diagnosing hepatocellular carcinoma. <i>European Radiology</i> , 2017 , 27, 2610-2618	8	33	
16	Progressive Enrichment of Stemness Features and Tumor Stromal Alterations in Multistep Hepatocarcinogenesis. <i>PLoS ONE</i> , 2017 , 12, e0170465	3.7	8	
15	Tumor stroma with senescence-associated secretory phenotype in steatohepatitic hepatocellular carcinoma. <i>PLoS ONE</i> , 2017 , 12, e0171922	3.7	21	

14	IMP2/IGF2BP2 expression, but not IMP1 and IMP3, predicts poor outcome in patients and high tumor growth rate in xenograft models of gallbladder cancer. <i>Oncotarget</i> , 2017 , 8, 89736-89745	3.3	23
13	Increased expression of stemness markers and altered tumor stroma in hepatocellular carcinoma under TACE-induced hypoxia: A biopsy and resection matched study. <i>Oncotarget</i> , 2017 , 8, 99359-99371	3.3	16
12	Transarterial Therapies for Hepatocellular Carcinoma. <i>Liver Cancer</i> , 2016 , 6, 27-33	9.1	34
11	Poor outcome of hepatocellular carcinoma with stemness marker under hypoxia: resistance to transarterial chemoembolization. <i>Modern Pathology</i> , 2016 , 29, 1038-49	9.8	26
10	Interaction of hepatitis B virus X protein with PARP1 results in inhibition of DNA repair in hepatocellular carcinoma. <i>Oncogene</i> , 2016 , 35, 5435-5445	9.2	32
9	Single Hepatocellular Carcinoma: Preoperative MR Imaging to Predict Early Recurrence after Curative Resection. <i>Radiology</i> , 2015 , 276, 433-43	20.5	91
8	Increased expression of stathmin and elongation factor 1 In precancerous nodules with telomere dysfunction in hepatitis B viral cirrhotic patients. <i>Journal of Translational Medicine</i> , 2014 , 12, 154	8.5	4
7	Increased expression of CCN2, epithelial membrane antigen, and fibroblast activation protein in hepatocellular carcinoma with fibrous stroma showing aggressive behavior. <i>PLoS ONE</i> , 2014 , 9, e10509.	4 ^{3.7}	28
6	ECatenin activated hepatocellular adenoma: a report of three cases in Korea. <i>Gut and Liver</i> , 2014 , 8, 452-8	4.8	8
5	Added value of subtraction imaging in detecting arterial enhancement in small (. <i>European Radiology</i> , 2013 , 23, 924-30	8	38
4	Histological characteristics of small hepatocellular carcinomas showing atypical enhancement patterns on gadoxetic acid-enhanced MR imaging. <i>Journal of Magnetic Resonance Imaging</i> , 2013 , 37, 138	8 4: 91	25
3	Gadoxetic acid-enhanced MRI findings of early hepatocellular carcinoma as defined by new histologic criteria. <i>Journal of Magnetic Resonance Imaging</i> , 2012 , 35, 393-8	5.6	36
2	Hyperintense lesions on gadoxetate disodium-enhanced hepatobiliary phase imaging. <i>American Journal of Roentgenology</i> , 2012 , 199, W575-86	5.4	17
1	Differentiation of early hepatocellular carcinoma from benign hepatocellular nodules on gadoxetic	3.4	55