

Takeshi Iwasaki

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/132733/publications.pdf>

Version: 2024-02-01

58
papers

1,089
citations

393982

19
h-index

454577

30
g-index

60
all docs

60
docs citations

60
times ranked

1318
citing authors

#	ARTICLE	IF	CITATIONS
1	Histological background of dedifferentiated solitary fibrous tumour. <i>Journal of Clinical Pathology</i> , 2022, 75, 397-403.	1.0	4
2	Merkel cell polyomavirus-negative Merkel cell carcinoma is associated with JAK-STAT and MEK-ERK pathway activation. <i>Cancer Science</i> , 2022, 113, 251-260.	1.7	9
3	Histological and immunohistochemical features and genetic alterations in the malignant progression of giant cell tumor of bone: a possible association with TP53 mutation and loss of H3K27 trimethylation. <i>Modern Pathology</i> , 2022, 35, 640-648.	2.9	13
4	TROP2 Expression in Sebaceous and Sweat Gland Carcinoma. <i>Journal of Clinical Medicine</i> , 2022, 11, 607.	1.0	2
5	Recurrent Massive Hemothorax of Unknown Etiology in an 85-Year-Old Man. <i>Chest</i> , 2022, 161, e103-e110.	0.4	2
6	Myxoid type and non-myxoid type of intimal sarcoma in large vessels and heart: review of histological and genetic profiles of 20 cases. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2022, 480, 919-925.	1.4	7
7	Approach for reclassification of collecting duct carcinoma and comparative histopathological analysis with SMARCB1/INI1-deficient renal cell carcinoma and fumarate hydratase-deficient renal cell carcinoma. <i>Human Pathology</i> , 2022, 124, 36-44.	1.1	3
8	Cyclin-dependent kinase 8 is an independent prognosticator in uterine leiomyosarcoma. <i>Pathology Research and Practice</i> , 2022, 235, 153920.	1.0	1
9	Association of PD-L1 and IDO1 expression with JAK-STAT pathway activation in soft-tissue leiomyosarcoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2021, 147, 1451-1463.	1.2	10
10	The association between the expression of PD-L1 and CMTM6 in undifferentiated pleomorphic sarcoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2021, 147, 2003-2011.	1.2	8
11	Morphological, immunohistochemical, and genomic analyses of papillary renal neoplasm with reverse polarity. <i>Human Pathology</i> , 2021, 112, 48-58.	1.1	24
12	Risk factors for excessive postoperative sliding of femoral trochanteric fracture in elderly patients: A retrospective multicenter study. <i>Injury</i> , 2021, 52, 3369-3376.	0.7	10
13	Clinicopathological features and immunohistochemical utility of NTRK-, ALK-, and ROS1-rearranged papillary thyroid carcinomas and anaplastic thyroid carcinomas. <i>Human Pathology</i> , 2020, 106, 82-92.	1.1	18
14	Frequent MN1 Gene Mutations in Malignant Peripheral Nerve Sheath Tumor. <i>Anticancer Research</i> , 2020, 40, 6221-6228.	0.5	1
15	PD-L1 and IDO1 expression and tumor-infiltrating lymphocytes in osteosarcoma patients: comparative study of primary and metastatic lesions. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020, 146, 2607-2620.	1.2	21
16	First case of pyrin-associated autoinflammation with neutrophilic dermatosis complicated by amyloidosis. <i>Rheumatology</i> , 2020, 59, e41-e43.	0.9	11
17	Establishment and Characterization of a Novel Primitive Yolk Sac Tumour Cell Line, TC587. <i>Anticancer Research</i> , 2020, 40, 759-766.	0.5	2
18	Clinicopathological review of solitary fibrous tumors: dedifferentiation is a major cause of patient death. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2019, 475, 467-477.	1.4	40

#	ARTICLE	IF	CITATIONS
19	Decreased H3K27me3 Expression Is Associated With Merkel Cell Polyomavirus-negative Merkel Cell Carcinoma, Especially Combined With Cutaneous Squamous Cell Carcinoma. <i>Anticancer Research</i> , 2019, 39, 5573-5579.	0.5	15
20	Current Update on the Molecular Biology of Cutaneous Sarcoma: Dermatofibrosarcoma Protuberans. <i>Current Treatment Options in Oncology</i> , 2019, 20, 29.	1.3	25
21	Expression of the IDO1/TDO2-AhR pathway in tumor cells or the tumor microenvironment is associated with Merkel cell polyomavirus status and prognosis in Merkel cell carcinoma. <i>Human Pathology</i> , 2019, 84, 52-61.	1.1	23
22	Histone H3.3 sub-variant H3mm7 is required for normal skeletal muscle regeneration. <i>Nature Communications</i> , 2018, 9, 1400.	5.8	23
23	Insulin-like growth factor messenger RNA-binding protein is an independent prognostic factor in uterine leiomyosarcoma. <i>Histopathology</i> , 2018, 72, 739-748.	1.6	10
24	Histopathological and genetic review of phosphaturic mesenchymal tumours, mixed connective tissue variant. <i>Histopathology</i> , 2018, 72, 460-471.	1.6	33
25	Diagnostic utility of histone H3.3 G34W, G34R, and G34V mutant-specific antibodies for giant cell tumors of bone. <i>Human Pathology</i> , 2018, 73, 41-50.	1.1	81
26	Merkel cell polyomavirus and Langerhans cell neoplasm. <i>Cell Communication and Signaling</i> , 2018, 16, 49.	2.7	10
27	Sensitive detection of fluorescence in western blotting by merging images. <i>PLoS ONE</i> , 2018, 13, e0191532.	1.1	13
28	Claudin 6 expression is useful to distinguish myxofibrosarcomas from other myxoid soft tissue tumors. <i>Pathology Research and Practice</i> , 2017, 213, 674-679.	1.0	6
29	Coexpression of SALL4 with HDAC1 and/or HDAC2 is associated with underexpression of PTEN and poor prognosis in patients with hepatocellular carcinoma. <i>Human Pathology</i> , 2017, 64, 69-75.	1.1	18
30	Crystal Structure and Characterization of Novel Human Histone H3 Variants, H3.6, H3.7, and H3.8. <i>Biochemistry</i> , 2017, 56, 2184-2196.	1.2	20
31	Association of expression of the hedgehog signal with Merkel cell polyomavirus infection and prognosis of Merkel cell carcinoma. <i>Human Pathology</i> , 2017, 69, 8-14.	1.1	20
32	Higher Expression of Activation-induced Cytidine Deaminase Is Significantly Associated with Merkel Cell Polyomavirus-negative Merkel Cell Carcinomas. <i>Yonago Acta Medica</i> , 2017, 60, 145-153.	0.3	8
33	Higher Expression of Activation-induced Cytidine Deaminase Is Significantly Associated with Merkel Cell Polyomavirus-negative Merkel Cell Carcinomas. <i>Yonago Acta Medica</i> , 2017, 60, 145-153.	0.3	4
34	Prognostic Significance of Forkhead Box M1 (FOXM1) Expression and Antitumor Effect of FOXM1 Inhibition in Angiosarcoma. <i>Journal of Cancer</i> , 2016, 7, 823-830.	1.2	18
35	Identification of Immunoglobulin Gene Sequences from a Small Read Number of mRNA-Seq Using Hybridomas. <i>PLoS ONE</i> , 2016, 11, e0165473.	1.1	11
36	Lower expression of CADM1 and higher expression of MAL in Merkel cell carcinomas are associated with Merkel cell polyomavirus infection and better prognosis. <i>Human Pathology</i> , 2016, 48, 1-8.	1.1	20

#	ARTICLE	IF	CITATIONS
37	Comment on "Cytokeratin 20-negative Merkel cell carcinoma is infrequently associated with the Merkel cell polyomavirus". <i>Modern Pathology</i> , 2016, 29, 89-90.	2.9	2
38	Acute-phase ITIH4 levels distinguish multi-system from single-system Langerhans cell histiocytosis via plasma peptidomics. <i>Clinical Proteomics</i> , 2015, 12, 16.	1.1	8
39	Severe post-transplant lymphoproliferative disorder after living donor liver transplantation. <i>Hepatology Research</i> , 2015, 45, 356-362.	1.8	1
40	Phosphohistone-H3 (PHH3) is prognostic relevant in Merkel cell carcinomas but Merkel cell polyomavirus is a more powerful prognostic factor than AJCC clinical stage, PHH3, Ki-67 or mitotic indices. <i>Pathology International</i> , 2015, 65, 404-409.	0.6	10
41	Comparison of Akt/mTOR/4E-BP1 pathway signal activation and mutations of PIK3CA in Merkel cell polyomavirus-positive and Merkel cell polyomavirus-negative carcinomas. <i>Human Pathology</i> , 2015, 46, 210-216.	1.1	28
42	Interleukin-1 loop model for pathogenesis of Langerhans cell histiocytosis. <i>Cell Communication and Signaling</i> , 2015, 13, 13.	2.7	30
43	Reactivation of persistent Epstein-Barr virus (EBV) causes secretion of thyrotropin receptor antibodies (TRAbs) in EBV-infected B lymphocytes with TRAbs on their surface. <i>Autoimmunity</i> , 2015, 48, 328-335.	1.2	23
44	Multiple Skin Cancers in a Renal Transplant Recipient: A Patient Report with Analyses of Human Papillomavirus and Human Polyomavirus Infection. <i>Yonago Acta Medica</i> , 2015, 58, 145-50.	0.3	1
45	Immunoglobulin Expressions Are Only Associated With MCPyV-positive Merkel Cell Carcinomas But Not With MCPyV-negative Ones. <i>American Journal of Surgical Pathology</i> , 2014, 38, 1627-1635.	2.1	21
46	Merkel cell polyomavirus (MCPyV) strains in Japanese merkel cell carcinomas (MCC) are distinct from Caucasian type MCPyVs: genetic variability and phylogeny of MCPyV genomes obtained from Japanese MCPyV-infected MCCs. <i>Virus Genes</i> , 2014, 48, 233-242.	0.7	20
47	Presence of Epstein-Barr virus-infected B lymphocytes with thyrotropin receptor antibodies on their surface in Graves' disease patients and in healthy individuals. <i>Autoimmunity</i> , 2014, 47, 193-200.	1.2	15
48	A new in situ hybridization and immunohistochemistry with a novel antibody to detect small T-antigen expressions of Merkel cell polyomavirus (MCPyV). <i>Diagnostic Pathology</i> , 2014, 9, 65.	0.9	17
49	High viral load of Merkel cell polyomavirus DNA sequences in Langerhans cell sarcoma tissues. <i>Infectious Agents and Cancer</i> , 2014, 9, 15.	1.2	12
50	Merkel cell polyomavirus DNA sequences in peripheral blood and tissues from patients with Langerhans cell histiocytosis. <i>Human Pathology</i> , 2014, 45, 119-126.	1.1	24
51	Usefulness of significant morphologic characteristics in distinguishing between Merkel cell polyomavirus-positive and Merkel cell polyomavirus-negative Merkel cell carcinomas. <i>Human Pathology</i> , 2013, 44, 1912-1917.	1.1	45
52	Merkel cell polyomavirus infection in both components of a combined Merkel cell carcinoma and basal cell carcinoma with ductal differentiation; each component had a similar but different novel Merkel cell polyomavirus large T antigen truncating mutation. <i>Human Pathology</i> , 2013, 44, 442-447.	1.1	15
53	Detection of Merkel Cell Polyomavirus in the Human Tissues from 41 Japanese Autopsy Cases Using Polymerase Chain Reaction. <i>Intervirology</i> , 2013, 56, 1-5.	1.2	28
54	Association of Merkel cell polyomavirus infection with clinicopathological differences in Merkel cell carcinoma. <i>Human Pathology</i> , 2012, 43, 2282-2291.	1.1	111

#	ARTICLE	IF	CITATIONS
55	Association of Merkel cell polyomavirus infection with morphologic differences in Merkel cell carcinoma. <i>Human Pathology</i> , 2011, 42, 632-640.	1.1	117
56	A pediatric intramedullary spinal cord tumor with unusual solid-cystic and papillary features: A case report. <i>Neuropathology</i> , 2011, 31, 632-638.	0.7	2
57	Effect of Sodium Thiosulfate on Cisplatin Removal With Complete Hepatic Venous Isolation and Extracorporeal Charcoal Hemoperfusion: A Pharmacokinetic Evaluation. <i>Annals of Surgical Oncology</i> , 2001, 8, 449-457.	0.7	11
58	Pancreatic hamartoma: detection of harbouring <i>NAB2::STAT6</i> fusion gene. <i>Histopathology</i> , 0, , .	1.6	3