Norbert Schmitz

List of Publications by Citations

Source: https://exaly.com/author-pdf/1330252/norbert-schmitz-publications-by-citations.pdf

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

 252
 7,581
 46
 76

 papers
 citations
 h-index
 g-index

 279
 8,843
 4.2
 6.06

 ext. papers
 ext. citations
 avg, IF
 L-index

#	Paper	IF	Citations
252	The association between obesity and anxiety disorders in the population: a systematic review and meta-analysis. <i>International Journal of Obesity</i> , 2010 , 34, 407-19	5.5	388
251	Project Ice Storm: prenatal maternal stress affects cognitive and linguistic functioning in 5 1/2-year-old children. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2008 , 47, 10	63 ⁷ -72	288
250	The Symptom Check-List-90-R (SCL-90-R): a German validation study. <i>Quality of Life Research</i> , 2000 , 9, 185-93	3.7	265
249	Association of diabetes with anxiety: a systematic review and meta-analysis. <i>Journal of Psychosomatic Research</i> , 2013 , 74, 89-99	4.1	228
248	Disability in depression and back pain: evaluation of the World Health Organization Disability Assessment Schedule (WHO DAS II) in a primary care setting. <i>Journal of Clinical Epidemiology</i> , 2003 , 56, 507-14	5.7	228
247	Diagnosing mental disorders in primary care: the General Health Questionnaire (GHQ) and the Symptom Check List (SCL-90-R) as screening instruments. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 1999 , 34, 360-6	4.5	180
246	Predictors of rate and time to remission in first-episode psychosis: a two-year outcome study. <i>Psychological Medicine</i> , 2006 , 36, 649-58	6.9	168
245	Psychometric properties of the General Health Questionnaire (GHQ-12) in a German primary care sample. <i>Acta Psychiatrica Scandinavica</i> , 1999 , 100, 462-8	6.5	146
244	Epidemiological associations between gambling behavior, substance use & mood and anxiety disorders. <i>Journal of Gambling Studies</i> , 2006 , 22, 275-87	3	123
243	Stress, burnout and locus of control in German nurses. <i>International Journal of Nursing Studies</i> , 2000 , 37, 95-9	5.8	114
242	The relationship between work stress and mental disorders in men and women: findings from a population-based study. <i>Journal of Epidemiology and Community Health</i> , 2008 , 62, 42-7	5.1	103
241	Disabilities, quality of life, and mental disorders associated with smoking and nicotine dependence. <i>American Journal of Psychiatry</i> , 2003 , 160, 1670-6	11.9	100
240	Changes in perceived job strain and the risk of major depression: results from a population-based longitudinal study. <i>American Journal of Epidemiology</i> , 2009 , 169, 1085-91	3.8	99
239	On the association between diabetes and mental disorders in a community sample: results from the German National Health Interview and Examination Survey. <i>Diabetes Care</i> , 2003 , 26, 1841-6	14.6	98
238	Testing definitions of symptom remission in first-episode psychosis for prediction of functional outcome at 2 years. <i>Schizophrenia Bulletin</i> , 2010 , 36, 1001-8	1.3	92
237	Association of functional variants in the dopamine D2-like receptors with risk for gambling behaviour in healthy Caucasian subjects. <i>Biological Psychology</i> , 2010 , 85, 33-7	3.2	92
236	The association between C-reactive protein, Interleukin-6 and depression among older adults in the community: A systematic review and meta-analysis. <i>Experimental Gerontology</i> , 2018 , 102, 109-132	4.5	90

(2010-2007)

Joint effect of depression and chronic conditions on disability: results from a population-based study. <i>Psychosomatic Medicine</i> , 2007 , 69, 332-8	3.7	80	
Prenatal exposure to a natural disaster increases risk for obesity in 5 ^[] -year-old children. <i>Pediatric Research</i> , 2012 , 71, 126-31	3.2	78	
On the relation between neuroticism, self-esteem, and depression: results from the National Comorbidity Survey. <i>Comprehensive Psychiatry</i> , 2003 , 44, 169-76	7-3	78	
Fatigue and Personality Pathology in the Mentally Ill. <i>Journal of General Internal Medicine</i> , 2000 , 15, 1-1	4	78	
A population-based longitudinal study on work environmental factors and the risk of major depressive disorder. <i>American Journal of Epidemiology</i> , 2012 , 176, 52-9	3.8	74	
Socioeconomic status and the risk of major depression: the Canadian National Population Health Survey. <i>Journal of Epidemiology and Community Health</i> , 2010 , 64, 447-52	5.1	72	
Assessing clinically significant change: application to the SCL-90-R. <i>Psychological Reports</i> , 2000 , 86, 263-	- 7:4 6	72	
Determinants of help-seeking and system related components of delay in the treatment of first-episode psychosis. <i>Schizophrenia Research</i> , 2007 , 96, 206-14	3.6	70	
How do neighborhoods affect depression outcomes? A realist review and a call for the examination of causal pathways. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2014 , 49, 873-87	4.5	69	
Determinants of negative pathways to care and their impact on service disengagement in first-episode psychosis. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2013 , 48, 125-36	4.5	67	
The prevalence of mental disorders in the working population over the period of global economic crisis. <i>Canadian Journal of Psychiatry</i> , 2010 , 55, 598-605	4.8	66	
Validation of the English and French versions of the Community Assessment of Psychic Experiences (CAPE) with a Montreal community sample. <i>Schizophrenia Research</i> , 2007 , 95, 86-95	3.6	64	
The association between physical exercises and health-related quality of life in subjects with mental disorders: results from a cross-sectional survey. <i>Preventive Medicine</i> , 2004 , 39, 1200-7	4.3	62	
Psychological distress and socioeconomic status in single mothers and their children in a German city. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2003 , 38, 59-68	4.5	61	
Dopamine transporter 3'-UTR VNTR genotype and ADHD: a pharmaco-behavioural genetic study with methylphenidate. <i>Neuropsychopharmacology</i> , 2007 , 32, 1370-6	8.7	58	
Early predictors of nonadherence to antipsychotic therapy in first-episode psychosis. <i>Canadian Journal of Psychiatry</i> , 2009 , 54, 28-35	4.8	57	
Adherence to Disease Modifying Drugs among Patients with Multiple Sclerosis in Germany: A Retrospective Cohort Study. <i>PLoS ONE</i> , 2015 , 10, e0133279	3.7	57	
Workplace characteristics, depression, and health-related presenteeism in a general population sample. <i>Journal of Occupational and Environmental Medicine</i> , 2010 , 52, 836-42	2	56	
	Prenatal exposure to a natural disaster increases risk for obesity in 5fl -year-old children. <i>Pediatric Research</i> , 2012, 71, 126-31 On the relation between neuroticism, self-esteem, and depression: results from the National Comorbidity Survey. <i>Comprehensive Psychiatry</i> , 2003, 44, 169-76 Fatigue and Personality Pathology in the Mentally Ill. <i>Journal of General Internal Medicine</i> , 2000, 15, 1-1 A population-based longitudinal study on work environmental factors and the risk of major depressive disorder. <i>American Journal of Epidemiology</i> , 2012, 176, 52-9 Socioeconomic status and the risk of major depression: the Canadian National Population Health Survey. <i>Journal of Epidemiology and Community Health</i> , 2010, 64, 447-52 Assessing clinically significant change: application to the SCL-90-R. <i>Psychological Reports</i> , 2000, 86, 263- Determinants of help-seeking and system related components of delay in the treatment of first-episode psychosis. <i>Schizophrenia Research</i> , 2007, 96, 206-14 How do neighborhoods affect depression outcomes? A realist review and a call for the examination of causal pathways. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2014, 49, 873-87 Determinants of negative pathways to care and their impact on service disengagement in first-episode psychosis. <i>Schizophrenia Psychiatry and Psychiatric Epidemiology</i> , 2013, 48, 125-36 The prevalence of mental disorders in the working population over the period of global economic crisis. <i>Canadian Journal of Psychiatry</i> , 2010, 55, 598-605 Validation of the English and French versions of the Community Assessment of Psychic Experiences (CAPE) with a Montreal community sample. <i>Schizophrenia Research</i> , 2007, 95, 86-95 The association between physical exercises and health-related quality of life in subjects with mental disorders: results from a cross-sectional survey. <i>Preventive Medicine</i> , 2004, 39, 1200-7 Psychological distress and socioeconomic status in single mothers and their children in a German city. <i>Social Psychiatry</i> , 2009, 54, 28-35	Prenatal exposure to a natural disaster increases risk for obesity in 50 -year-old children. <i>Pediatric Research</i> , 2012, 71, 126-31 On the relation between neuroticism, self-esteem, and depression: results from the National Comorbidity Survey. <i>Comprehensive Psychiatry</i> , 2003, 44, 169-76 73 Fatigue and Personality Pathology in the Mentally Ill. <i>Journal of General Internal Medicine</i> , 2000, 15, 1-1 A population-based longitudinal study on work environmental factors and the risk of major depressive disorder. <i>American Journal of Epidemiology</i> , 2012, 176, 52-9 Socioeconomic status and the risk of major depression: the Canadian National Population Health Survey. <i>Journal of Epidemiology and Community Health</i> , 2010, 64, 447-52 Assessing clinically significant change: application to the SCL-90-R. <i>Psychological Reports</i> , 2000, 86, 263-746 Determinants of help-seeking and system related components of delay in the treatment of first-episode psychosis. <i>Schizophrenia Research</i> , 2007, 96, 206-14 How do neighborhoods affect depression outcomes? A realist review and a call for the examination of causal pathways. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2014, 49, 873-87 Determinants of negative pathways to care and their impact on service disengagement in first-episode psychosis. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2013, 48, 125-36 4.5 The prevalence of mental disorders in the working population over the period of global economic crisis. <i>Canadian Journal of Psychiatry</i> , 2010, 55, 598-605 Validation of the English and French versions of the Community Assessment of Psychic Experiences (CAPE) with a Montreal community sample. <i>Schizophrenia Research</i> , 2007, 95, 86-95 The association between physical exercises and health-related quality of life in subjects with mental disorders; results from a cross-sectional survey. <i>Preventive Medicine</i> , 2004, 39, 1200-7 Psychological distress and socioeconomic status in single mothers and their children in a German city. <i>Social Psychiatry and Psychiatri</i>	Prenatal exposure to a natural disaster increases risk for obesity in 50 -year-old children. Pediatric Research, 2012, 71, 126-31 On the relation between neuroticism, self-esteem, and depression: results from the National Comorbidity Survey. Comprehensive Psychiatry, 2003, 44, 169-76 Fatigue and Personality Pathology in the Mentally III. Journal of General Internal Medicine, 2000, 15, 1-1 A population-based longitudinal study on work environmental factors and the risk of major depressive disorder. American Journal of Epidemiology, 2012, 176, 52-9 Socioeconomic status and the risk of major depression: the Canadian National Population Health Survey. Journal of Epidemiology and Community Health, 2010, 64, 447-52 Assessing clinically significant change: application to the SCL-90-R. Psychological Reports, 2000, 86, 263-746 Determinants of help-seeking and system related components of delay in the treatment of first-episode psychosis. Schizophrenia Research, 2007, 96, 206-14 How do neighborhoods affect depression outcomes? A realist review and a call for the examination of causal pathways. Social Psychiatry and Psychiatric Epidemiology, 2014, 49, 873-87 Determinants of negative pathways to care and their impact on service disengagement in first-episode psychosis. Social Psychiatry and Psychiatric Epidemiology, 2013, 48, 125-36 The prevalence of mental disorders in the working population over the period of global economic disist. Canadian Journal of Psychiatry, 2010, 55, 598-605 The association between physical exercises and health-related quality of life in subjects with mental disorders: results from a cross-sectional survey. Preventive Medicine, 2004, 39, 1200-7 Psychological distress and socioeconomic status in single mothers and their children in a German dty. Social Psychiatry, 2009, 54, 28-35 Adherence to Disease Modifying Drugs among Patients with Multiple Sclerosis in Germany: A Retrospective Cohort Study. PLoS ONE, 2015, 10, e0133279 Workplace characteristics, depression, and health-related pres

217	Determinants of participation in leisure activities among adolescents with cerebral palsy. <i>Research in Developmental Disabilities</i> , 2013 , 34, 2621-34	2.7	55
216	Factors influencing relapse during a 2-year follow-up of first-episode psychosis in a specialized early intervention service. <i>Psychological Medicine</i> , 2008 , 38, 1585-93	6.9	54
215	Association between obesity-related gene FTO and ADHD. <i>Obesity</i> , 2013 , 21, E738-44	8	51
214	The longitudinal association from obesity to depression: results from the 12-year National Population Health Survey. <i>Obesity</i> , 2010 , 18, 1033-8	8	50
213	Three job-related stress models and depression: a population-based study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2012 , 47, 185-93	4.5	49
212	Recurrent subthreshold depression in type 2 diabetes: an important risk factor for poor health outcomes. <i>Diabetes Care</i> , 2014 , 37, 970-8	14.6	49
211	Comparing three-year extension of early intervention service to regular care following two years of early intervention service in first-episode psychosis: a randomized single blind clinical trial. <i>World Psychiatry</i> , 2017 , 16, 278-286	14.4	48
210	Effects of trauma-focused psychotherapy upon war refugees. <i>Journal of Traumatic Stress</i> , 2009 , 22, 585	-928	48
209	The longitudinal associations between C-reactive protein and depressive symptoms: evidence from the English Longitudinal Study of Ageing (ELSA). <i>International Journal of Geriatric Psychiatry</i> , 2015 , 30, 976-84	3.9	46
208	Predictors of quality of life in a longitudinal study of users with severe mental disorders. <i>Health and Quality of Life Outcomes</i> , 2013 , 11, 92	3	46
207	A comparison study of multiple measures of adherence to antipsychotic medication in first-episode psychosis. <i>Journal of Clinical Psychopharmacology</i> , 2010 , 30, 64-7	1.7	46
206	Interpersonal Problems as Predictors of Symptom Related Treatment Outcome in Longterm Psychotherapy. <i>Psychotherapy Research</i> , 1996 , 6, 164-176	3.6	45
205	Investigating the longitudinal association between diabetes and anxiety: a systematic review and meta-analysis. <i>Diabetic Medicine</i> , 2018 , 35, 677-693	3.5	44
204	Rehabilitation service utilization in children and youth with cerebral palsy. <i>Child: Care, Health and Development</i> , 2014 , 40, 275-82	2.8	42
203	Impact of a specialized early intervention service for psychotic disorders on patient characteristics, service use, and hospital costs in a defined catchment area. <i>Canadian Journal of Psychiatry</i> , 2006 , 51, 895-903	4.8	41
202	Problem gambling in bipolar disorder: results from the Canadian Community Health Survey. <i>Journal of Affective Disorders</i> , 2007 , 102, 27-34	6.6	40
201	QF2011: a protocol to study the effects of the Queensland flood on pregnant women, their pregnancies, and their children's early development. <i>BMC Pregnancy and Childbirth</i> , 2015 , 15, 109	3.2	39
200	Depression and risk of type 2 diabetes: the potential role of metabolic factors. <i>Molecular Psychiatry</i> , 2016 , 21, 1726-1732	15.1	39

(2013-2009)

199	Association between neighborhood-level deprivation and disability in a community sample of people with diabetes. <i>Diabetes Care</i> , 2009 , 32, 1998-2004	14.6	39	
198	From symptom relief to interpersonal change: Treatment outcome and effectiveness in inpatient psychotherapy. <i>Psychotherapy Research</i> , 2008 , 18, 615-24	3.6	39	
197	Association of trait-defined, eating-disorder sub-phenotypes with (biallelic and triallelic) 5HTTLPR variations. <i>Journal of Psychiatric Research</i> , 2009 , 43, 1086-94	5.2	38	
196	A multisite Canadian study of outcome of first-episode psychosis treated in publicly funded early intervention services. <i>Canadian Journal of Psychiatry</i> , 2007 , 52, 563-71	4.8	37	
195	Maternal smoking during pregnancy and ADHD: a comprehensive clinical and neurocognitive characterization. <i>Nicotine and Tobacco Research</i> , 2013 , 15, 149-57	4.9	36	
194	Associations between diabetes, major depressive disorder and generalized anxiety disorder comorbidity, and disability: findings from the 2012 Canadian Community Health SurveyMental Health (CCHS-MH). <i>Journal of Psychosomatic Research</i> , 2015 , 78, 137-42	4.1	36	
193	Psychometric evaluation of the Problem Gambling Severity Index-Chinese version (PGSI-C). <i>Journal of Gambling Studies</i> , 2011 , 27, 453-66	3	36	
192	The four Es of problem gambling: a psychological measure of risk. <i>Journal of Gambling Studies</i> , 2006 , 22, 101-20	3	36	
191	Relevance of the 5-HTTLPR polymorphism and childhood abuse to increased psychiatric comorbidity in women with bulimia-spectrum disorders. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, 981-90	4.6	36	
190	Associations between depression, chronic physical health conditions, and disability in a community sample: A focus on the persistence of depression. <i>Journal of Affective Disorders</i> , 2015 , 179, 6-13	6.6	35	
189	Picture me playing-a portrait of participation and enjoyment of leisure activities in adolescents with cerebral palsy. <i>Research in Developmental Disabilities</i> , 2013 , 34, 1001-10	2.7	35	
188	Posttraumatic stress disorder and treatment seeking in a nationally representative Canadian military sample. <i>Journal of Traumatic Stress</i> , 2006 , 19, 847-58	3.8	35	
187	Association between Depressive Symptoms and Cognitive Function in Persons with Diabetes Mellitus: A Systematic Review. <i>PLoS ONE</i> , 2016 , 11, e0160809	3.7	35	
186	Indicators of self-rated health in the Canadian population with diabetes. <i>Diabetic Medicine</i> , 2012 , 29, 1021-8	3.5	34	
185	A prediction algorithm for first onset of major depression in the general population: development and validation. <i>Journal of Epidemiology and Community Health</i> , 2014 , 68, 418-24	5.1	34	
184	Validation of the alcohol use disorders identification test and the drug abuse screening test in first episode psychosis. <i>Canadian Journal of Psychiatry</i> , 2008 , 53, 26-33	4.8	34	
183	Effects of Short-Term Dynamic Psychotherapy for Neurotic, Somatoform, and Personality Disorders: a Prospective 1-Year Follow-Up Study. <i>Psychotherapy Research</i> , 2001 , 11, 187-200	3.6	34	
182	Behavioral difficulties in adolescents with cerebral palsy. <i>Journal of Child Neurology</i> , 2013 , 28, 27-33	2.5	33	

181	COMT Val108/158Met gene variant, birth weight, and conduct disorder in children with ADHD. Journal of the American Academy of Child and Adolescent Psychiatry, 2006 , 45, 1363-1369	7.2	33
180	A time-limited, problem-orientated psychotherapeutic intervention in Type 1 diabetic patients with complications: a randomized controlled trial. <i>Diabetic Medicine</i> , 2002 , 19, 814-21	3.5	32
179	Application of stratum-specific likelihood ratios in mental health screening. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2000 , 35, 375-9	4.5	32
178	A five-year randomized parallel and blinded clinical trial of an extended specialized early intervention vs. regular care in the early phase of psychotic disorders: study protocol. <i>BMC Psychiatry</i> , 2015 , 15, 22	4.2	30
177	The incidence of major depression in Canada: the National Population Health Survey. <i>Journal of Affective Disorders</i> , 2010 , 123, 158-63	6.6	30
176	Self-rated health: a predictor for the three year incidence of major depression in individuals with Type II diabetes. <i>Journal of Affective Disorders</i> , 2013 , 145, 100-5	6.6	28
175	Perceived needs for and use of workplace accommodations by individuals with a depressive and/or anxiety disorder. <i>Journal of Occupational and Environmental Medicine</i> , 2011 , 53, 1268-72	2	28
174	Development and validation of prediction algorithms for major depressive episode in the general population. <i>Journal of Affective Disorders</i> , 2013 , 151, 39-45	6.6	27
173	Cyclical relationship between depressive symptoms and diabetes distress in people with Type 2 diabetes mellitus: results from the Montreal Evaluation of Diabetes Treatment Cohort Study. <i>Diabetic Medicine</i> , 2015 , 32, 1272-8	3.5	27
172	Trait-defined eating-disorder subtypes and history of childhood abuse. <i>International Journal of Eating Disorders</i> , 2010 , 43, 428-32	6.3	27
171	Psychometric properties of the German version of the NEO-FFI in psychosomatic outpatients. <i>Personality and Individual Differences</i> , 2001 , 31, 713-722	3.3	27
170	Development and validation of a prediction algorithm for use by health professionals in prediction of recurrence of major depression. <i>Depression and Anxiety</i> , 2014 , 31, 451-7	8.4	26
169	C-reactive protein, depressive symptoms, and risk of diabetes: results from the English Longitudinal Study of Ageing (ELSA). <i>Journal of Psychosomatic Research</i> , 2014 , 77, 180-6	4.1	26
168	The Canadian Problem Gambling Index: an evaluation of the scale and its accompanying profiler software in a clinical setting. <i>Journal of Gambling Studies</i> , 2011 , 27, 467-85	3	26
167	COMT Val108/158Met polymorphism and the modulation of task-oriented behavior in children with ADHD. <i>Neuropsychopharmacology</i> , 2008 , 33, 3069-77	8.7	25
166	Improving screening for mental disorders in the primary care setting by combining the GHQ-12 and SCL-90-R subscales. <i>Comprehensive Psychiatry</i> , 2001 , 42, 166-73	7.3	25
165	Cardiovascular reactions induced by unpredictable, predictable, and controllable painful stimuli during sphygmomanometry. <i>International Journal of Psychophysiology</i> , 2001 , 40, 161-5	2.9	25
164	Should psychological distress screening in the community account for self-perceived health status?. <i>Canadian Journal of Psychiatry</i> , 2009 , 54, 526-33	4.8	24

(2011-2015)

163	Stability of leisure participation from school-age to adolescence in individuals with cerebral palsy. <i>Research in Developmental Disabilities</i> , 2015 , 47, 73-9	2.7	23
162	The 5-HTTLPR polymorphism of the serotonin transporter gene and short term behavioral response to methylphenidate in children with ADHD. <i>BMC Psychiatry</i> , 2010 , 10, 50	4.2	23
161	Prediabetes, depressive and anxiety symptoms, and risk of type 2 diabetes: A community-based cohort study. <i>Journal of Psychosomatic Research</i> , 2016 , 89, 85-90	4.1	23
160	Type 2 Diabetes Self-Management: Role of Diet Self-Efficacy. <i>Canadian Journal of Diabetes</i> , 2012 , 36, 337-344	2.1	22
159	The influence of perceived social support on medication adherence in first-episode psychosis. <i>Canadian Journal of Psychiatry</i> , 2013 , 58, 59-65	4.8	22
158	The relationship between mental disorders and medical service utilization in a representative community sample. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2002 , 37, 380-6	4.5	22
157	Measures of depression and risk of type 2 diabetes: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2020 , 265, 224-232	6.6	21
156	Adverse Childhood Experiences and the Risk of Diabetes: Examining the Roles of Depressive Symptoms and Cardiometabolic Dysregulations in the Whitehall II Cohort Study. <i>Diabetes Care</i> , 2018 , 41, 2120-2126	14.6	21
155	Inconsistency in the relationship between duration of untreated psychosis (DUP) and negative symptoms: sorting out the problem of heterogeneity. <i>Schizophrenia Research</i> , 2007 , 93, 152-9	3.6	21
154	Comparison of the standard and the computerized versions of the Symptom Check List (SCL-90-R): a randomized trial. <i>Acta Psychiatrica Scandinavica</i> , 2000 , 102, 147-52	6.5	21
153	Associations between coping strategies and mental health in individuals with type 2 diabetes: Prospective analyses. <i>Health Psychology</i> , 2016 , 35, 78-86	5	20
152	Diabetes distress and neighborhood characteristics in people with type 2 diabetes. <i>Journal of Psychosomatic Research</i> , 2013 , 75, 147-52	4.1	20
151	Neighbourhood characteristics and 10-year risk of depression in Canadian adults with and without a chronic illness. <i>Health and Place</i> , 2014 , 30, 279-86	4.6	20
150	Body weight and ADHD: examining the role of self-regulation. <i>PLoS ONE</i> , 2013 , 8, e55351	3.7	20
149	Exploring trajectories of diabetes distress in adults with type 2 diabetes; a latent class growth modeling approach. <i>Journal of Affective Disorders</i> , 2015 , 188, 160-6	6.6	19
148	Are you doing what you want to do? Leisure preferences of adolescents with cerebral palsy. <i>Developmental Neurorehabilitation</i> , 2015 , 18, 234-40	1.8	19
147	Exploring the association of psychological status with self-rated diabetes control: results from the Montreal evaluation of diabetes treatment study. <i>Psychosomatics</i> , 2013 , 54, 35-43	2.6	19
146	Factors associated with disability and depressive symptoms among individuals with diabetes: a community study in Quebec. <i>Psychosomatics</i> , 2011 , 52, 167-77	2.6	19

145	Case manager- and patient-rated alliance as a predictor of medication adherence in first-episode psychosis. <i>Journal of Clinical Psychopharmacology</i> , 2012 , 32, 465-9	1.7	19
144	Apomorphine effects on episodic memory in young healthy volunteers. <i>Neuropsychologia</i> , 2008 , 46, 292	-3.00	19
143	Short duration of untreated psychosis enhances negative symptom remission in extended early intervention service for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2019 , 140, 65-76	6.5	18
142	The Neighbourhood Built Environment and Trajectories of Depression Symptom Episodes in Adults: A Latent Class Growth Analysis. <i>PLoS ONE</i> , 2015 , 10, e0133603	3.7	18
141	Associations between Depressive Symptoms and Social Support in Adults with Diabetes: Comparing Directionality Hypotheses with a Longitudinal Cohort. <i>Annals of Behavioral Medicine</i> , 2016 , 50, 348-57	4.5	18
140	Mastery motivation in adolescents with cerebral palsy. <i>Research in Developmental Disabilities</i> , 2013 , 34, 3384-92	2.7	17
139	Predictors of 1-year outcomes of major depressive disorder among individuals with a lifetime diagnosis: a population-based study. <i>Psychological Medicine</i> , 2012 , 42, 327-34	6.9	17
138	Self-management of oral anticoagulation: long-term results. <i>Journal of Internal Medicine</i> , 2003 , 254, 515	5 -6 0.8	17
137	Preferred Features of E-Mental Health Programs for Prevention of Major Depression in Male Workers: Results From a Canadian National Survey. <i>Journal of Medical Internet Research</i> , 2016 , 18, e132	7.6	17
136	Cardiometabolic dysregulation and cognitive decline: potential role of depressive symptoms. British Journal of Psychiatry, 2018 , 212, 96-102	5.4	16
135	Does job strain interact with psychosocial factors outside of the workplace in relation to the risk of major depression? The Canadian National Population Health Survey. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2011 , 46, 577-84	4.5	16
134	The impact of psychological distress on functional disability in asthma: results from the Canadian community health survey. <i>Psychosomatics</i> , 2009 , 50, 42-9	2.6	16
133	Subjective distress in first-episode psychosis: role of symptoms and self-esteem. <i>Microbial Biotechnology</i> , 2007 , 1, 251-258	3.3	16
132	Computerized administration of the Symptom Checklist (SCL-90-R) and the Inventory of Interpersonal Problems (IIP-C) in psychosomatic outpatients. <i>Psychiatry Research</i> , 1999 , 87, 217-21	9.9	16
131	Gender differences in the relationship between anxiety symptoms and physical inactivity in a community-based sample of adults with type 2 diabetes. <i>Canadian Journal of Diabetes</i> , 2014 , 38, 444-50	2.1	15
130	Place and health in diabetes: the neighbourhood environment and risk of depression in adults with type 2 diabetes. <i>Diabetic Medicine</i> , 2015 , 32, 944-50	3.5	15
129	Investigation of anxiety and depression symptom co-morbidity in a community sample with type 2 diabetes: Associations with indicators of self-care. <i>Canadian Journal of Public Health</i> , 2016 , 106, e496-50) ^{3.2}	15
128	Long-term effects of a community intervention for early identification of first-episode psychosis. Acta Psychiatrica Scandinavica, 2008, 117, 440-8	6.5	15

(2013-2013)

127	Trajectories of self-rated health in people with diabetes: associations with functioning in a prospective community sample. <i>PLoS ONE</i> , 2013 , 8, e83088	3.7	15	
126	Dimensions of insight in schizophrenia: Exploratory factor analysis of items from multiple self- and interviewer-rated measures of insight. <i>Schizophrenia Research</i> , 2018 , 199, 319-325	3.6	14	
125	A controlled evaluation of a targeted early case detection intervention for reducing delay in treatment of first episode psychosis. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2014 , 49, 1711-8	4.5	14	
124	The pattern of depressive symptoms in people with type 2 diabetes: a prospective community study. <i>Journal of Psychosomatic Research</i> , 2013 , 74, 128-34	4.1	14	
123	Receiving workplace mental health accommodations and the outcome of mental disorders in employees with a depressive and/or anxiety disorder. <i>Journal of Occupational and Environmental Medicine</i> , 2013 , 55, 1293-9	2	14	
122	Obesity and the risk of disability in a 12-year cohort study: the role of psychological distress. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2011 , 46, 1173-9	4.5	14	
121	Duration of active psychosis and first-episode psychosis negative symptoms. <i>Microbial Biotechnology</i> , 2017 , 11, 63-71	3.3	13	
120	The association between smoking and depression in a Canadian community-based sample with type 2 diabetes. <i>Canadian Journal of Diabetes</i> , 2013 , 37, 150-5	2.1	13	
119	Measuring psychological outcomes following pediatric intensive care unit hospitalization: psychometric analysis of the Children's Critical Illness Impact Scale. <i>Pediatric Critical Care Medicine</i> , 2011 , 12, 635-42	3	13	
118	The SF-36 summary scores and their relation to mental disorders: physical functioning may affect performance of the summary scores. <i>Journal of Clinical Epidemiology</i> , 2007 , 60, 163-70	5.7	13	
117	Rise in systolic blood pressure during sphygmomanometry depends on the maximum inflation pressure of the arm cuff. <i>Journal of Hypertension</i> , 1994 , 12, 825???830	1.9	13	
116	Comorbid depressive and anxiety symptoms and the risk of type 2 diabetes: Findings from the Lifelines Cohort Study. <i>Journal of Affective Disorders</i> , 2018 , 238, 24-31	6.6	13	
115	Adequacy of Help Received by Individuals with Severe Mental Disorders After a Major Healthcare Reform in Quebec: Predictors and Changes at 5-Year Follow-Up. <i>Administration and Policy in Mental Health and Mental Health Services Research</i> , 2016 , 43, 799-812	3.1	12	
114	Longitudinal relationships between depression and functioning in people with type 2 diabetes. <i>Annals of Behavioral Medicine</i> , 2014 , 47, 172-9	4.5	12	
113	Association of depression and anxiety symptoms with functional disability and disability days in a community sample with type 2 diabetes. <i>Psychosomatics</i> , 2014 , 55, 659-67	2.6	12	
112	A Comparison of Developmental Outcomes of Adolescent Neonatal Intensive Care Unit Survivors Born with a Congenital Heart Defect or Born Preterm. <i>Journal of Pediatrics</i> , 2019 , 207, 34-41.e2	3.6	12	
111	The association between sleep and diabetes outcomes - A systematic review. <i>Diabetes Research and Clinical Practice</i> , 2020 , 161, 108035	7:4	11	
110	Pursuit of Complementary and Alternative Medicine Treatments in Adolescents With Cerebral Palsy. <i>Journal of Child Neurology</i> , 2013 , 28, 1443-1447	2.5	11	

109	Smoking and its association with disability in chronic conditions: results from the Canadian Community and Health Survey 2.1. <i>Nicotine and Tobacco Research</i> , 2007 , 9, 959-64	4.9	11
108	Gender differences in the association between lifestyle behaviors and diabetes distress in a community sample of adults with type 2 diabetes. <i>Journal of Diabetes</i> , 2016 , 8, 269-78	3.8	11
107	Diabetes Complications and Depressive Symptoms: Prospective Results From the Montreal Diabetes Health and Well-Being Study. <i>Psychosomatic Medicine</i> , 2017 , 79, 603-612	3.7	10
106	Trajectories of anxiety symptoms and associations with incident cardiovascular disease in adults with type 2 diabetes. <i>Journal of Psychosomatic Research</i> , 2018 , 104, 95-100	4.1	10
105	Changes in depressive symptoms and changes in lifestyle-related indicators: a 1-year follow-up study among adults with type 2 diabetes in Quebec. <i>Canadian Journal of Diabetes</i> , 2013 , 37, 243-248	2.1	10
104	Self-rated diabetes control in a Canadian population with type 2 diabetes: associations with health behaviours and outcomes. <i>Diabetes Research and Clinical Practice</i> , 2012 , 95, 162-8	7.4	10
103	The interaction of obesity and psychological distress on disability. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2010 , 45, 531-40	4.5	10
102	A decade of spontaneous long-term course of psychogenic impairment in a community population sample. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 1999 , 34, 651-6	4.5	10
101	COVID-19 and Depressive Symptoms: A Community-based Study in Quebec, Canada. <i>Canadian Journal of Psychiatry</i> , 2020 , 65, 733-735	4.8	10
100	Anxiety symptoms and functioning in a community sample of individuals with type 2 diabetes: A longitudinal study. <i>Journal of Diabetes</i> , 2016 , 8, 854-862	3.8	10
99	Comparison of clinical outcomes following 2 years of treatment of first-episode psychosis in urban early intervention services in Canada and India. <i>British Journal of Psychiatry</i> , 2020 , 217, 514-520	5.4	9
98	Anxiety moderates cardiovascular responses to painful stimuli during sphygmomanometry. <i>International Journal of Psychophysiology</i> , 1999 , 33, 253-7	2.9	9
97	The Effects of an E-Mental Health Program and Job Coaching on the Risk of Major Depression and Productivity in Canadian Male Workers: Protocol for a Randomized Controlled Trial. <i>JMIR Research Protocols</i> , 2016 , 5, e218	2	9
96	Depressive symptoms and glycated hemoglobin A1c: a reciprocal relationship in a prospective cohort study. <i>Psychological Medicine</i> , 2016 , 46, 945-55	6.9	9
95	Context and contact: a comparison of patient and family engagement with early intervention services for psychosis in India and Canada. <i>Psychological Medicine</i> , 2020 , 1-10	6.9	8
94	Alcohol consumption, depressive symptoms, and the incidence of diabetes-related complications. <i>Journal of Diabetes</i> , 2019 , 11, 14-22	3.8	8
93	Good vs. poor self-rated diabetes control: differences in cardiovascular risk and self-care activities. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2014 , 122, 236-9	2.3	8
92	The relationship between manual ability and ambulation in adolescents with cerebral palsy. <i>Physical and Occupational Therapy in Pediatrics</i> , 2013 , 33, 243-52	2.1	8

91	Development and psychometric evaluation of a 10-item short form inventory of gambling situations. <i>Journal of Gambling Studies</i> , 2011 , 27, 115-28	3	8
90	Business mergers and acquisitions and the risk of mental disorders: a population-based study. <i>Occupational and Environmental Medicine</i> , 2012 , 69, 569-73	2.1	8
89	Properties of the Symptom Check List (SCL-90-R) in a Psychosomatic Consultation-Liaison Setting. <i>Psychological Reports</i> , 2002 , 90, 1201-1207	1.6	8
88	Risk of Diabetes in Older Adults with Co-Occurring Depressive Symptoms and Cardiometabolic Abnormalities: Prospective Analysis from the English Longitudinal Study of Ageing. <i>PLoS ONE</i> , 2016 , 11, e0155741	3.7	8
87	Progress of negative symptoms over the initial 5 years of a first episode of psychosis. <i>Psychological Medicine</i> , 2019 , 49, 66-74	6.9	8
86	Does Social Support Moderate the Association Among Major Depression, Generalized Anxiety Disorder, and Functional Disability in Adults With Diabetes?. <i>Psychosomatics</i> , 2017 , 58, 364-374	2.6	7
85	A randomized controlled trial to examine the impacts of disclosing personalized depression risk information on the outcomes of individuals who are at high risk of developing major depression: a research protocol. <i>BMC Psychiatry</i> , 2019 , 19, 285	4.2	7
84	Sex Differences in Clinical and Functional Outcomes among Patients Treated in an Early Intervention Service for Psychotic Disorders: An Observational Study. <i>Canadian Journal of Psychiatry</i> , 2019 , 64, 708-717	4.8	7
83	The longitudinal effects of neighbourhood social and material deprivation change on psychological distress in urban, community-dwelling Canadian adults. <i>Public Health</i> , 2015 , 129, 932-40	4	7
82	Psychological distress and short-term disability in people with diabetes: results from the Canadian Community Health Survey. <i>Journal of Psychosomatic Research</i> , 2008 , 65, 165-72	4.1	7
81	Depressive symptoms and sleep problems as risk factors for heart disease: a prospective community study. <i>Epidemiology and Psychiatric Sciences</i> , 2019 , 29, e50	5.1	7
80	Depressive symptoms, prediabetes, and incident diabetes in older English adults. <i>International Journal of Geriatric Psychiatry</i> , 2017 , 32, 1450-1458	3.9	6
79	Catechol-o-methyltransferase gene and executive function in children with ADHD. <i>Journal of Attention Disorders</i> , 2014 , 18, 202-11	3.7	6
78	Educational and rehabilitation service utilization in adolescents born preterm or with a congenital heart defect and at high risk for disability. <i>Developmental Medicine and Child Neurology</i> , 2017 , 59, 1056-	1062	6
77	A Longitudinal Investigation of Anxiety and Depressive Symptomatology and Exercise Behaviour Among Adults With Type 2 Diabetes Mellitus. <i>Canadian Journal of Diabetes</i> , 2017 , 41, 73-81	2.1	6
76	Mental disorders and hypertension: factors associated with awareness and treatment of hypertension in the general population of Germany. <i>Psychosomatic Medicine</i> , 2006 , 68, 246-52	3.7	6
75	Properties of the Symptom Check List (SCL-90-R) in a psychosomatic consultation-liaison setting. <i>Psychological Reports</i> , 2002 , 90, 1201-7	1.6	6
74	Was bewirken Therapieempfehlungen bei psychogen erkrankten Patienten einer psychosomatischen Konsiliarambulanz?. <i>Zeitschrift Fur Psychosomatische Medizin Und Psychotherapie</i> , 1999 , 45, 95-112	1.3	6

73	. Journal of Clinical Monitoring and Computing, 1997 , 14, 251-254		6
72	Adverse Childhood Experiences and the Risk of Coronary Heart Disease in Adulthood: Examining Potential Psychological, Biological, and Behavioral Mediators in the Whitehall II Cohort Study. <i>Journal of the American Heart Association</i> , 2021 , 10, e019013	6	6
71	Habit strength as a moderator of the association between symptoms of poor mental health and unintentional non-adherence to oral hypoglycemic medication in adults with type 2 diabetes. Journal of Health Psychology, 2019 , 24, 321-326	3.1	6
70	Does providing personalized depression risk information lead to increased psychological distress and functional impairment? Results from a mixed-methods randomized controlled trial. <i>Psychological Medicine</i> , 2020 , 1-9	6.9	5
69	The accuracy of depression risk perception in high risk Canadians. <i>Journal of Affective Disorders</i> , 2020 , 265, 410-415	6.6	5
68	Anxiety and Depression Symptom Comorbidity and the Risk of Heart Disease: A Prospective Community-Based Cohort Study. <i>Psychosomatic Medicine</i> , 2020 , 82, 296-304	3.7	5
67	The Relationship between Diabetes and Mental Health Conditions in an Aging Population. <i>Canadian Journal of Diabetes</i> , 2016 , 40, 4-5	2.1	5
66	Types of smokers in a community sample of individuals with Type 2 diabetes: a latent class analysis. <i>Diabetic Medicine</i> , 2012 , 29, 586-92	3.5	5
65	Redevelopment of tertiary psychiatric services in British Columbia: a prospective study of clinical, social, and residential outcomes of former long-stay inpatients. <i>Schizophrenia Research</i> , 2013 , 149, 96-1	03 ⁶	5
64	Weight cycling and depressive symptoms in diabetes: a community-based study of adults with type 2 diabetes mellitus in Quebec. <i>Canadian Journal of Diabetes</i> , 2014 , 38, 456-60	2.1	5
63	Tinnitus als soziale Infektion?. <i>Psychotherapeut</i> , 2005 , 50, 318-327	0.5	5
62	Assessing Clinically Significant Change: Application to the SCL-90 $f R$		5
61	Evaluating lifestyle and health-related characteristics of older adults with co-occurring depressive symptoms and cardiometabolic abnormalities. <i>International Journal of Geriatric Psychiatry</i> , 2016 , 31, 66-	. 75 9	5
60	Development of a Tool to Measure Women Agency in India. <i>Journal of Human Development and Capabilities</i> , 2019 , 20, 26-53	1.6	5
59	Trajectories of social support in adults with type 2 diabetes: Associations with depressive symptoms and functional disability. <i>International Journal of Geriatric Psychiatry</i> , 2019 , 34, 480-487	3.9	5
58	Adipose tissue dysregulation at the onset of psychosis: Adipokines and social determinants of health. <i>Psychoneuroendocrinology</i> , 2021 , 123, 104915	5	5
57	Economic Evaluation of Extended Early Intervention Service vs Regular Care Following 2 Years of Early Intervention: Secondary Analysis of a Randomized Controlled Trial. <i>Schizophrenia Bulletin</i> , 2021 , 47, 465-473	1.3	5
56	Medication adherence in first episode psychosis: the role of pre-onset subthreshold symptoms. <i>Acta Psychiatrica Scandinavica</i> , 2019 , 139, 336-347	6.5	4

(2022-2015)

55	Demographic, lifestyle, and health characteristics of older adults with prediabetes in England. <i>Preventive Medicine</i> , 2015 , 77, 74-9	4.3	4
54	Assessing the longitudinal associations and stability of smoking and depression syndrome over a 4-year period in a community sample with type 2 diabetes 24. <i>Journal of Diabetes</i> , 2015 , 7, 95-101	3.8	4
53	Self-rated health in diabetes: should the question be the first administered?. <i>Diabetes Research and Clinical Practice</i> , 2012 , 97, e27-30	7.4	4
52	Epidemiology of antidepressant medication use in the Canadian diabetes population. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2010 , 45, 911-9	4.5	4
51	Research in Psychotherapeutic Diagnosis and the Concept of Socio-Empirical Markers. <i>Psychotherapy Research</i> , 1997 , 7, 71-82	3.6	4
50	Is olanzapine a brain-sparing medication?. Archives of General Psychiatry, 2006, 63, 1292		4
49	A Bootstrap Method To Test If Study Dropouts Are Missing Randomly. <i>Quality and Quantity</i> , 2002 , 36, 1-16	2.4	4
48	Participation in leisure activities in adolescents with congenital heart defects. <i>Developmental Medicine and Child Neurology</i> , 2020 , 62, 946-953	3.3	4
47	Self-reported use of diabetes healthcare services in a Quebec community-based sample: impact of depression status. <i>Public Health</i> , 2014 , 128, 63-9	4	3
46	Lifestyle Care Indicators in Individuals with Major, Minor No Depression: A Community-Based Diabetes Study in Quebec. <i>Canadian Journal of Diabetes</i> , 2011 , 35, 22-30	2.1	3
45	Promoting measured genes and measured environments: on the importance of careful statistical analyses and biological relevance. <i>Archives of General Psychiatry</i> , 2007 , 64, 377-8; author reply 378-9		3
44	Self-concept and Engagement in LiFe (SELF): A waitlist-controlled pilot study of a novel psychological intervention to target illness engulfment in enduring schizophrenia and related psychoses. <i>Schizophrenia Research</i> , 2021 , 228, 567-574	3.6	3
43	Measures of depression and incident type 2 diabetes in a community sample. <i>Annals of Epidemiology</i> , 2021 , 55, 4-9	6.4	3
42	Understanding Components of Duration of Untreated Psychosis and Relevance for Early Intervention Services in the Canadian Context: Comprendre les Composantes de la Durê de la Psychose Non Traité et la Pertinence de Services D'intervention Prêoce Dans le Contexte	4.8	3
41	Associations between cognitive function, metabolic factors and depression: A prospective study in Quebec, Canada. <i>Journal of Affective Disorders</i> , 2021 , 283, 77-83	6.6	3
40	The relationship between depression risk perception and self-help behaviours in high risk Canadians: a cross-sectional study. <i>BMC Public Health</i> , 2020 , 20, 876	4.1	2
39	Effects of pain-incompatible imagery on tolerance of pain, heart rate, and skin resistance. <i>Perceptual and Motor Skills</i> , 1997 , 84, 939-43	2.2	2
38	Whose responsibility? 'Part 1 of 2: A'scale to assess how stakeholders apportion responsibilities for addressing the needs of persons with mental health problems <i>International Journal of Mental Health Systems</i> , 2022 , 16, 1	3.6	2

37	Unfinished business: Functional outcomes in a randomized controlled trial of a three-year extension of early intervention versus regular care following two years of early intervention for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2021 ,	6.5	2
36	Gender Differences in Cardiovascular Responses to Sphygmomanometric Measurements. <i>Journal of Psychophysiology</i> , 2001 , 15, 43-46	1	2
35	Association between depressive symptoms, metabolic risk factors, and cognitive function: cross-sectional results from a community study in Quebec, Canada. <i>Aging and Mental Health</i> , 2021 , 25, 2003-2010	3.5	2
34	The effect of affordable daycare on women's mental health: Evidence from a cluster randomized trial in rural India. <i>Social Science and Medicine</i> , 2018 , 217, 32-41	5.1	2
33	Associations Between Depressive Symptoms and Indices of Obesity in Adults With Prediabetes and Normal Blood Glucose Levels: Results From the Emotional Health and Wellbeing Study. <i>Canadian Journal of Diabetes</i> , 2018 , 42, 626-631	2.1	2
32	Longitudinal associations between number of cigarettes per day and depressive symptoms in adult smokers with type 2 diabetes: A path analysis approach. <i>Journal of Psychosomatic Research</i> , 2019 , 125, 109737	4.1	1
31	Response to Kawada: C-reactive protein, depressive symptoms and incident diabetes mellitus with special emphasis on physical activity. <i>Journal of Psychosomatic Research</i> , 2015 , 78, 407-8	4.1	1
30	The association between job strain, depressive symptoms, and cardiovascular disease risk: results from a cross-sectional population-based study in QuBec, Canada. <i>International Archives of Occupational and Environmental Health</i> , 2020 , 93, 1013-1021	3.2	1
29	Response to Kawada: Self-rated health and depression of patients with diabetes: how to handle with wide range of ageing and diabetic status?. <i>Diabetic Medicine</i> , 2013 , 30, 755-6	3.5	1
28	THREE AUTHORS REPLY. American Journal of Epidemiology, 2009, 170, 132-133	3.8	1
27	An observational study of antipsychotic medication discontinuation in first-episode psychosis: clinical and functional outcomes <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2022 , 1	4.5	1
26	Symptomatik und Lebensereignisse bei ost- und westdeutschen Psychotherapiepatienten/innen. Zeitschrift Fur Psychosomatische Medizin Und Psychotherapie, 1999 , 45, 157-169	1.3	1
25	Trajectories of Depressive Symptoms and Incident Diabetes: A Prospective Study. <i>Annals of Behavioral Medicine</i> , 2021 ,	4.5	1
24	PROPERTIES OF THE SYMPTOM CHECK LIST (SCL-90-R) IN A PSYCHOSOMATIC CONSULTATION-LIAISON SETTING. <i>Psychological Reports</i> , 2002 , 90, 1201	1.6	1
23	Job strain and the incidence of heart diseases: A prospective community study in Quebec, Canada. Journal of Psychosomatic Research, 2020 , 139, 110268	4.1	1
22	Depression-related weight change and incident diabetes in a community sample. <i>Scientific Reports</i> , 2021 , 11, 13575	4.9	1
21	Dyadic associations between physical activity and body mass index in couples in which one partner has diabetes: results from the Lifelines cohort study. <i>Journal of Behavioral Medicine</i> , 2020 , 43, 143-149	3.6	1
20	Patient satisfaction with random assignment to extended early intervention for psychosis vs regular care: Relationship with service engagement. <i>Microbial Biotechnology</i> , 2021 , 15, 746-750	3.3	1

19	Do mental disorders moderate the association between diabetes status and alcohol consumption?. <i>Psychology, Health and Medicine</i> , 2018 , 23, 277-284	2.1	1
18	Whose responsibility? Part 2 of 2: views of patients, families, and clinicians about responsibilities for addressing the needs of persons with mental health problems in Chennai, India and Montreal, Canada International Journal of Mental Health Systems, 2022, 16, 2	3.6	O
17	Manualized group cognitive behavioral therapy for social anxiety in first-episode psychosis: a randomized controlled trial <i>Psychological Medicine</i> , 2022 , 1-10	6.9	O
16	Incident psychopharmacological treatment and psychiatric hospital contact in individuals with newly developed type 2 diabetes - a register-based cohort study. <i>Acta Neuropsychiatrica</i> , 2021 , 33, 72-8	34 ^{3.9}	О
15	Measurement invariance testing of the patient health questionnaire-9 (PHQ-9) across people with and without diabetes mellitus from the NHANES, EMHS and UK Biobank datasets. <i>Journal of Affective Disorders</i> , 2021 , 292, 311-318	6.6	O
14	Response to Kawada: Anxiety symptoms and functioning in patients with type 2 diabetes. <i>Journal of Diabetes</i> , 2017 , 9, 635-636	3.8	
13	DEPRESSIVE SYMPTOMS, C-REACTIVE PROTEIN, AND INCIDENCE OF DIABETES. <i>Journal of Epidemiology and Community Health</i> , 2015 , 69, A2.3-A3	5.1	
12	Association between smoking and depression in patients with type 2 diabetes: A Response. <i>Journal of Diabetes</i> , 2015 , 7, 590	3.8	
11	Response to comment on Schmitz et al. Recurrent subthreshold depression in type 2 diabetes: an important risk factor for poor health outcomes. Diabetes care 2014;37:970-978. <i>Diabetes Care</i> , 2014 , 37, e146-7	14.6	
10	Letter to the editor: smoking and mental health in young women - challenges in interpretation. <i>Psychological Medicine</i> , 2013 , 43, 445	6.9	
9	P2-94 Types of smokers, depression and disability in type 2 diabetes: a latent class analysis. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, A246-A246	5.1	
8	The Relationship Between of Depression, Chronic Diseases and Short-Term Disability: Results from the Canadian Community and Health Survey. <i>American Journal of Epidemiology</i> , 2006 , 163, S220-S220	3.8	
7	494: Hypertension and Mental Disorders: Factors Associated with Treatment of Hypertension in a Community Sample. <i>American Journal of Epidemiology</i> , 2005 , 161, S124-S124	3.8	
6	Underestimation of sensorimotor task-induced blood-pressure changes by posttask sphygmomanometry. <i>Perceptual and Motor Skills</i> , 1995 , 81, 483-90	2.2	
5	Conditional associations between childhood cat ownership and psychotic experiences in adulthood: A retrospective study <i>Journal of Psychiatric Research</i> , 2022 , 148, 197-203	5.2	
4	Users' perceptions about receiving personalized depression risk information: findings from a qualitative study. <i>BMC Psychiatry</i> , 2021 , 21, 581	4.2	
3	The impact of providing personalized depression risk information on self-help and help-seeking behaviors: Results from a mixed methods randomized controlled trial. <i>Depression and Anxiety</i> , 2021 , 38, 917-924	8.4	
2	An approach to sample size calculation for research on duration of untreated psychosis. <i>Schizophrenia Research</i> , 2019 , 206, 466-467	3.6	

Self-help behaviors partially mediate the relationship between personalized depression risk disclosure and psychological distress: A mediation analysis using data from a randomized controlled trial. *Journal of Psychiatric Research*, **2021**, 140, 7-14

5.2