
## Laurent L Coquet

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1330785/publications.pdf Version: 2024-02-01



#	Article	lF	CITATIONS
1	Peptidomic Analysis of Skin Secretions of the Caribbean Frogs Leptodactylus insularum and Leptodactylus nesiotus (Leptodactylidae) Identifies an Ocellatin with Broad Spectrum Antimicrobial Activity. Antibiotics, 2020, 9, 718.	1.5	10
2	Membrane Proteocomplexome of Campylobacter jejuni Using 2-D Blue Native/SDS-PAGE Combined to Bioinformatics Analysis. Frontiers in Microbiology, 2020, 11, 530906.	1.5	2
3	Ethanol-Induced Alterations in Placental and Fetal Cerebrocortical Annexin-A4 and Cerebral Cavernous Malformation Protein 3 Are Associated With Reductions in Fetal Cortical VEGF Receptor Binding and Microvascular Density. Frontiers in Neuroscience, 2020, 14, 519.	1.4	6
4	Venom Peptide Repertoire of the European Myrmicine Ant <i>Manica rubida</i> : Identification of Insecticidal Toxins. Journal of Proteome Research, 2020, 19, 1800-1811.	1.8	30
5	Processing and Maturation of Cathepsin C Zymogen: A Biochemical and Molecular Modeling Analysis. International Journal of Molecular Sciences, 2019, 20, 4747.	1.8	12
6	Peptidomic analysis of the host-defense peptides in skin secretions of Rana graeca provides insight into phylogenetic relationships among Eurasian Rana species. Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 2019, 29, 228-234.	0.4	8
7	Dihydroquinoline Carbamate DQS1-02 as a Prodrug of a Potent Acetylcholinesterase Inhibitor for Alzheimer's Disease Therapy: Multigram-Scale Synthesis, Mechanism Investigations, in Vitro Safety Pharmacology, and Preliminary in Vivo Toxicology Profile. ACS Omega, 2018, 3, 18387-18397.	1.6	7
8	SAG12, a Major Cysteine Protease Involved in Nitrogen Allocation during Senescence for Seed Production in Arabidopsis thaliana. Plant and Cell Physiology, 2018, 59, 2052-2063.	1.5	66
9	InhA1-Mediated Cleavage of the Metalloprotease NprA Allows Bacillus cereus to Escape From Macrophages. Frontiers in Microbiology, 2018, 9, 1063.	1.5	19
10	Peptidomic analysis of the host-defense peptides in skin secretions of the Trinidadian leaf frog Phyllomedusa trinitatis (Phyllomedusidae). Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 2018, 28, 72-79.	0.4	7
11	Peptidomic analysis of skin secretions of the Mexican burrowing toad Rhinophrynus dorsalis (Rhinophrynidae): Insight into the origin of host-defense peptides within the Pipidae and characterization of a proline-arginine-rich peptide. Peptides, 2017, 97, 22-28.	1.2	5
12	Antioxidant, antityrosinase and antibiofilm activities of synthesized peptides derived from Vicia faba protein hydrolysate: A powerful agents in cosmetic application. Industrial Crops and Products, 2017, 109, 310-319.	2.5	60
13	Cytotoxic peptides with insulinâ€releasing activities from skin secretions of the Italian stream frog <scp><i>Rana italica</i></scp> (Ranidae). Journal of Peptide Science, 2017, 23, 769-776.	0.8	13
14	Proteomic Investigations of Proteases Involved in Cotyledon Senescence: A Model to Explore the Genotypic Variability of Proteolysis Machinery Associated with Nitrogen Remobilization Efficiency during the Leaf Senescence of Oilseed Rape. Proteomes, 2017, 5, 29.	1.7	10
15	Impact of chlorhexidine digluconate and temperature on curli production in <em>Escherichia coli</em> —consequence on its adhesion ability. AIMS Microbiology, 2017, 3, 915-937.	1.0	6
16	Proteomic Changes in Pseudomonas aeruginosa Biofilm Cells after Adaptive Resistance Development. Journal of Proteomics and Bioinformatics, 2016, 09, .	0.4	5
17	The Unusual Resistance of Avian Defensin AvBD7 to Proteolytic Enzymes Preserves Its Antibacterial Activity. PLoS ONE, 2016, 11, e0161573.	1.1	7
18	Mg deficiency affects leaf Mg remobilization and the proteome in Brassica napus. Plant Physiology and Biochemistry, 2016, 107, 337-343.	2.8	25

#	Article	IF	CITATIONS
19	Purification, Conformational Analysis, and Properties of a Family of Tigerinin Peptides from Skin Secretions of the Crowned Bullfrog <i>Hoplobatrachus occipitalis</i> . Journal of Natural Products, 2016, 79, 2350-2356.	1.5	12
20	MIB–MIP is a mycoplasma system that captures and cleaves immunoglobulin G. Proceedings of the National Academy of Sciences of the United States of America, 2016, 113, 5406-5411.	3.3	97
21	Peptidomic analysis of the extensive array of host-defense peptides in skin secretions of the dodecaploid frog Xenopus ruwenzoriensis (Pipidae). Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 2016, 19, 18-24.	0.4	4
22	Unraveling the effects of static magnetic field stress on cytosolic proteins of Salmonella by using a proteomic approach. Canadian Journal of Microbiology, 2016, 62, 338-348.	0.8	5
23	Host-defense and trefoil factor family peptides in skin secretions of the Mawa clawed frog Xenopus boumbaensis (Pipidae). Peptides, 2015, 72, 44-49.	1.2	5
24	Antimicrobial Peptide LL-37 Is Both a Substrate of Cathepsins S and K and a Selective Inhibitor of Cathepsin L. Biochemistry, 2015, 54, 2785-2798.	1.2	38
25	Determination of Multimodal Isotopic Distributions: The Case of a <sup>15</sup> N Labeled Protein Produced into Hairy Roots. Analytical Chemistry, 2015, 87, 5938-5946.	3.2	3
26	Evidence from peptidomic analysis of skin secretions that allopatric populations of Xenopus gilli (Anura:Pipidae) constitute distinct lineages. Peptides, 2015, 63, 118-125.	1.2	11
27	Copper-Deficiency in Brassica napus Induces Copper Remobilization, Molybdenum Accumulation and Modification of the Expression of Chloroplastic Proteins. PLoS ONE, 2014, 9, e109889.	1.1	41
28	Proteomic profile of pre - B2 lymphoblasts from children with acute lymphoblastic leukemia (ALL) in relation with the translocation (12; 21). Clinical Proteomics, 2014, 11, 31.	1.1	5
29	Host defense peptides from Lithobates forreri, Hylarana luctuosa, and Hylarana signata (Ranidae): Phylogenetic relationships inferred from primary structures of ranatuerin-2 and brevinin-2 peptides. Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 2014, 9, 49-57.	0.4	18
30	Host-defense peptides from skin secretions of Fraser's clawed frog Xenopus fraseri (Pipidae): Further insight into the evolutionary history of the Xenopodinae. Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 2014, 12, 45-52.	0.4	5
31	Host-defense peptides from skin secretions of the octoploid frogs Xenopus vestitus and Xenopus wittei (Pipidae): Insights into evolutionary relationships. Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 2014, 11, 20-28.	0.4	4
32	Escherichia coli Response to Uranyl Exposure at Low pH and Associated Protein Regulations. PLoS ONE, 2014, 9, e89863.	1.1	20
33	Evaluation of the Skin Peptide Defenses of the Oregon Spotted Frog Rana pretiosa Against Infection by the Chytrid Fungus Batrachochytrium dendrobatidis. Journal of Chemical Ecology, 2013, 39, 797-805.	0.9	11
34	Proteomic approach to Pseudomonas aeruginosa adaptive resistance to benzalkonium chloride. Journal of Proteomics, 2013, 89, 273-279.	1.2	23
35	Characterization of the host-defense peptides from skin secretions of Merlin's clawed frog Pseudhymenochirus merlini: Insights into phylogenetic relationships among the Pipidae. Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 2013, 8, 352-357.	0.4	15
36	VBNC Legionella pneumophila cells are still able to produce virulence proteins. Water Research, 2013, 47, 6606-6617.	5.3	77

#	Article	IF	CITATIONS
37	An immunomodulatory peptide related to frenatin 2 from skin secretions of the Tyrrhenian painted frog Discoglossus sardus (Alytidae). Peptides, 2013, 40, 65-71.	1.2	25
38	Chromogranin A Induces the Biogenesis of Granules with Calcium- and Actin-Dependent Dynamics and Exocytosis in Constitutively Secreting Cells. Endocrinology, 2012, 153, 4444-4456.	1.4	24
39	Enhanced Adhesion of Campylobacter jejuni to Abiotic Surfaces Is Mediated by Membrane Proteins in Oxygen-Enriched Conditions. PLoS ONE, 2012, 7, e46402.	1.1	60
40	Host-defense peptides from skin secretions of the tetraploid frogs Xenopus petersii and Xenopus pygmaeus, and the octoploid frog Xenopus lenduensis (Pipidae). Peptides, 2012, 33, 35-43.	1.2	24
41	The hymenochirins: A family of host-defense peptides from the Congo dwarf clawed frog Hymenochirus boettgeri (Pipidae). Peptides, 2012, 35, 269-275.	1.2	31
42	Host-defense peptides in skin secretions of the tetraploid frog Silurana epitropicalis with potent activity against methicillin-resistant Staphylococcus aureus (MRSA). Peptides, 2012, 37, 113-119.	1.2	30
43	Combined Proteomic and Molecular Approaches for Cloning and Characterization of Copper–Zinc Superoxide dismutase (Cu, Zn-SOD2) from Garlic (Allium sativum). Molecular Biotechnology, 2012, 52, 49-58.	1.3	6
44	Adaptation of Salmonella enterica Hadar under static magnetic field: effects on outer membrane protein pattern. Proteome Science, 2012, 10, 6.	0.7	15
45	Growth of Acinetobacter baumannii in Pellicle Enhanced the Expression of Potential Virulence Factors. PLoS ONE, 2011, 6, e26030.	1.1	80
46	Genome duplications within the Xenopodinae do not increase the multiplicity of antimicrobial peptides in Silurana paratropicalis and Xenopus andrei skin secretions. Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 2011, 6, 206-212.	0.4	12
47	Peptidomic analysis of skin secretions from the bullfrog Lithobates catesbeianus (Ranidae) identifies multiple peptides with potent insulin-releasing activity. Peptides, 2011, 32, 203-208.	1.2	34
48	Characterization of antimicrobial peptides in skin secretions from discrete populations of Lithobates chiricahuensis (Ranidae) from central and southern Arizona. Peptides, 2011, 32, 664-669.	1.2	25
49	Peptidomic analysis of skin secretions demonstrates that the allopatric populations of Xenopus muelleri (Pipidae) are not conspecific. Peptides, 2011, 32, 1502-1508.	1.2	29
50	Host defense peptides in skin secretions of the Oregon spotted frog Rana pretiosa: Implications for species resistance to chytridiomycosis. Developmental and Comparative Immunology, 2011, 35, 644-649.	1.0	24
51	Structure-Function Analysis of Grass Clip Serine Protease Involved in Drosophila Toll Pathway Activation. Journal of Biological Chemistry, 2011, 286, 12300-12307.	1.6	29
52	Primary structures of skin antimicrobial peptides indicate a close, but not conspecific, phylogenetic relationship between the leopard frogs Lithobates onca and Lithobates yavapaiensis (Ranidae). Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology, 2010, 151, 313-317.	1.3	5
53	Antimicrobial peptides with therapeutic potential from skin secretions of the Marsabit clawed frog Xenopus borealis (Pipidae). Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology, 2010, 152, 467-472.	1.3	34
54	Involvement of stathmin 1 in the neurotrophic effects of PACAP in PC12 cells. Journal of Neurochemistry, 2010, 114, 1498-1510.	2.1	12

#	Article	IF	CITATIONS
55	Evidence from the primary structures of dermal antimicrobial peptides that Rana tagoi okiensis and Rana tagoi tagoi (Ranidae) are not conspecific subspecies. Toxicon, 2010, 55, 430-435.	0.8	13
56	Antimicrobial peptides from the skin secretions of the South-East Asian frog Hylarana erythraea (Ranidae). Peptides, 2010, 31, 548-554.	1.2	31
57	EtpB Is a Pore-Forming Outer Membrane Protein Showing TpsB Protein Features Involved in the Two-Partner Secretion System. Journal of Membrane Biology, 2009, 230, 143-154.	1.0	4
58	A combined <sup>15</sup> N tracing/proteomics study in <i>Brassica napus</i> reveals the chronology of proteomics events associated with N remobilisation during leaf senescence induced by nitrate limitation or starvation. Proteomics, 2009, 9, 3580-3608.	1.3	78
59	Purification of peptides with differential cytolytic activities from the skin secretions of the Central American frog, Lithobates vaillanti (Ranidae). Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology, 2009, 150, 150-154.	1.3	17
60	Peptides with potent cytolytic activity from the skin secretions of the North American leopard frogs, Lithobates blairi and Lithobates yavapaiensis. Toxicon, 2009, 53, 699-705.	0.8	16
61	Antimicrobial peptides from the skin secretions of the New World frogs Lithobates capito and Lithobates warszewitschii (Ranidae). Peptides, 2009, 30, 1775-1781.	1.2	20
62	A potent, non-toxic insulin-releasing peptide isolated from an extract of the skin of the Asian frog, Hylarana guntheri (Anura:Ranidae). Regulatory Peptides, 2008, 151, 153-159.	1.9	48
63	Characterization of antimicrobial peptides from the skin secretions of the Malaysian frogs, Odorrana hosii and Hylarana picturata (Anura:Ranidae). Toxicon, 2008, 52, 465-473.	0.8	49
64	Purification and characterization of antimicrobial peptides from the Caribbean frog, Leptodactylus validus (Anura: Leptodactylidae). Peptides, 2008, 29, 1287-1292.	1.2	16
65	Peptidomic analysis of skin secretions from Rana heckscheri and Rana okaloosae provides insight into phylogenetic relationships among frogs of the Aquarana species group. Regulatory Peptides, 2007, 138, 87-93.	1.9	24
66	Peptides with differential cytolytic activity from skin secretions of the lemur leaf frog Hylomantis lemur (Hylidae: Phyllomedusinae). Toxicon, 2007, 50, 498-506.	0.8	60
67	Cytolytic peptides belonging to the brevinin-1 and brevinin-2 families isolated from the skin of the Japanese brown frog, Rana dybowskii. Toxicon, 2007, 50, 746-756.	0.8	46
68	Antimicrobial peptides from the skin of the Japanese mountain brown frog Rana ornativentris: Evidence for polymorphism among preprotemporin mRNAs. Peptides, 2007, 28, 524-532.	1.2	31
69	Expression of genes encoding antimicrobial and bradykinin-related peptides in skin of the stream brown frog Rana sakuraii. Peptides, 2007, 28, 505-514.	1.2	51
70	Peptidomic analysis of skin secretions supports separate species status for the tailed frogs, Ascaphus truei and Ascaphus montanus. Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 2007, 2, 121-125.	0.4	5
71	Antimicrobial peptides from the skin of the Tsushima brown frog Rana tsushimensis. Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology, 2006, 143, 42-49.	1.3	25
72	Functional characterization of Pseudomonas fluorescens OprE and OprQ membrane proteins. Biochemical and Biophysical Research Communications, 2006, 346, 1048-1052.	1.0	14

#	Article	IF	CITATIONS
73	Evidence from peptidomic analysis of skin secretions that the red-legged frogs, Rana aurora draytonii and Rana aurora aurora, are distinct species. Peptides, 2006, 27, 1305-1312.	1.2	44
74	Identification of Proteins Regulated by PACAP in PC12 Cells by 2D Gel Electrophoresis Coupled to Mass Spectrometry. Annals of the New York Academy of Sciences, 2006, 1070, 380-387.	1.8	11
75	A Proteomic Approach to Biofilm Cell Physiology. Methods in Biotechnology, 2006, , 403-414.	0.2	2
76	Immobilization Induces Alterations in the Outer Membrane Protein Pattern ofYersiniaruckeri. Journal of Proteome Research, 2005, 4, 1988-1998.	1.8	18
77	A family of acyclic brevinin-1 peptides from the skin of the Ryukyu brown frog Rana okinavana. Peptides, 2005, 26, 185-190.	1.2	31
78	Substitution of <i>Pichia pastoris</i> -Derived Recombinant Proteins with Mannose Containing O- and N-Linked Glycans Decreases Specificity of Diagnostic Tests. International Archives of Allergy and Immunology, 2004, 135, 187-195.	0.9	13
79	Adhesion of Yersinia ruckeri to fish farm materials: influence of cell and material surface properties. Colloids and Surfaces B: Biointerfaces, 2002, 26, 373-378.	2.5	49
80	Immobilized-cell physiology: current data and the potentialities of proteomics. Enzyme and Microbial Technology, 2002, 31, 201-212.	1.6	60